



Australian Government

Department of Sustainability, Environment,  
Water, Population and Communities



# Species group report card – dugongs

Supporting the marine bioregional plan  
for the North-west Marine Region

prepared under the *Environment Protection and Biodiversity Conservation Act 1999*


#### **Disclaimer**

© Commonwealth of Australia 2012

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without prior written permission from the Commonwealth. Requests and enquiries concerning reproduction and rights should be addressed to Department of Sustainability, Environment, Water, Population and Communities, Public Affairs, GPO Box 787 Canberra ACT 2601 or email [public.affairs@environment.gov.au](mailto:public.affairs@environment.gov.au)

#### **Images:**

Striped Nudibranch – C.Zwick and DSEWPaC, Raccoon butterfly fish – N.Wolfe, Display of colourful coral – Tourism WA, Red and yellow feather star (crinoids) – Tourism WA, Whale tail – Tourism WA, Snorkelling in Ningaloo Marine Park – Tourism WA, Green Turtle – Tourism WA, Black tip reef shark – N.Wolfe, Whale Shark – GBRMPA, Sea Grass Meadow – Lochman Transparencies



# CONTENTS

<b>Species group report card – dugongs .....</b>	<b>1</b>
1. Dugong of the North-west Marine Region.....	3
2. Vulnerabilities and pressures .....	4
3. Current protection measures.....	12
<b>References .....</b>	<b>13</b>



# SPECIES GROUP REPORT CARD – DUGONG

Supporting the marine bioregional plan for the North-west Marine Region prepared under the *Environment Protection and Biodiversity Conservation Act 1999*

## Report cards

The primary objective of the report cards is to provide accessible information on the conservation values found in Commonwealth marine regions. This information is maintained by the Department of Sustainability, Environment, Water, Population and Communities and is available online through the department's website ([www.environment.gov.au](http://www.environment.gov.au)). A glossary of terms relevant to marine bioregional planning is located at [www.environment.gov.au/marineplans](http://www.environment.gov.au/marineplans).

Reflecting the categories of conservation values, there are three types of report cards:

- species group report cards
- marine environment report cards
- heritage places report cards.

While the focus of these report cards is the Commonwealth marine environment, in some instances pressures and ecological processes occurring in state waters are referred to where there is connectivity between pressures and ecological processes in state and Commonwealth waters.





## Species group report cards

Species group report cards are prepared for large taxonomic groups that include species identified as conservation values in a region; that is, species that are listed under Part 13 of the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) and live in the Commonwealth marine area for all or part of their lifecycle. All listed threatened, migratory and marine species and all cetaceans occurring in Commonwealth waters are protected under the EPBC Act and are identified in the relevant marine bioregional plans as conservation values.

Species group report cards focus on species for which the region is important from a conservation perspective; for example, species of which a significant proportion of the population or an important life stage occurs in the region's waters.

For these species, the report cards:

- outline the conservation status of the species and the current state of knowledge about its ecology in the region
- define biologically important areas; that is, areas where aggregations of individuals of a species display biologically important behaviours
- assess the level of concern in relation to different pressures.



## 1. Dugong of the North-west Marine Region

The dugong (*Dugong dugon*) is the only living member of the family Dugongidae and is one of only four living species of the order Sirenia. A significant proportion of the world's dugong population occurs in coastal waters from Shark Bay in Western Australia to Moreton Bay in Queensland (Marsh et al. 2011). Current dugong distributions are believed to represent relict populations separated by large areas where they are either extinct or close to extinction.

Some of the coastal waters adjacent to the North-west Marine Region support significant populations of dugongs, including Shark Bay, which has an estimated population of around 10 000 individuals. Dugongs also occur in Exmouth Gulf and offshore on the North West Shelf, in and adjacent to Ningaloo Reef, in coastal waters close to Broome and along the Kimberley coast, and on the edge of the continental shelf at Ashmore Reef (DEWHA 2008).

Dugongs inhabit seagrass meadows in coastal waters, estuarine creeks and streams. They are migratory, which is believed to be related to their search for suitable seagrass beds or warmer waters. Seagrass is the preferred food of dugongs, but they are also known to eat algae and macroinvertebrates. Dugongs can live for more than 70 years, have a low reproduction rate and long generation time. Females reach sexual maturity between 6 and 17 years old, have a gestation period of 13 months and usually give birth to one calf.

The dugong is listed as migratory and marine under section 248 of the EPBC Act.

### Biologically important areas

Biologically important areas are areas that are particularly important for the conservation of the protected species and where aggregations of individuals display biologically important behaviour such as breeding, foraging, resting or migration. The presence of the observed behaviour is assumed to indicate that the habitat required for the behaviour is also present. Biologically important areas have been identified for some EPBC Act listed species found in the North-west Marine Region, using expert scientific knowledge about species' distribution, abundance and behaviour in the region. The selection of species was informed by the availability of scientific information, the conservation status of listed species and the importance of the region for the species. The range of species for which biologically important areas are identified will continue to expand as reliable spatial and scientific information becomes available.

Biologically important areas have been identified for dugongs. Behaviours used to identify biologically important areas for dugongs include foraging and nursing.

Biologically important areas are included in the North-west Marine Region Conservation Values Atlas ([www.environment.gov.au/cva](http://www.environment.gov.au/cva)).



## 2. Vulnerabilities and pressures

### Vulnerabilities

The population biology of dugongs renders them particularly vulnerable to mortality as adults (Marsh et al. 2011). Unexploited dugong populations are characterised by considerable longevity (a lifespan of more than 70 years), long gestation (12–14 months), litter sizes of one, long intervals between births (more than 2.5 years), prolonged periods until sexual maturity (6–17 years), and high and temporally stable adult survival (Marsh et al. 1984). Adult survival is the most important determinant of population growth. The maximum rate of population increase under optimum conditions when natural mortality is low is approximately 5 per cent per annum. The maximum sustainable mortality rate of adult females killed by human activities is around 1–2 per cent (Heinsohn et al. 2004; Marsh et al. 1997; Marsh et al. 2004); lower in areas where, or at times when, food supplies are low (Marsh & Kwan 2008; Marsh et al. 2002).

As dugongs are dependent on seagrass for food, any loss or degradation of seagrass due to anthropogenic activities (e.g. dredging, port construction, oil pollution), could adversely affect this species. The effect of seagrass loss or dieback on dugongs is twofold. Some dugongs remain in the affected area but lose body condition, reduce breeding and suffer increased mortality, while others move hundreds of kilometres with uncertain consequences (Preen & Marsh 1995; Marsh & Kwan 2008). Some animals that move die (Preen & Marsh 1995); others eventually migrate back to the affected area when it recovers (Marsh et al. 2011). Marsh and Kwan (2008) found that seagrass dieback events significantly decrease dugong reproductive rates.

### Analysis of pressures

On the basis of current information, pressures have been analysed for dugong in this report card. A summary of the pressure analysis is provided in Table 1. Only those pressures identified as *of concern* or *of potential concern* are discussed in further detail below.

An explanation of the pressure analysis process, including the definition of substantial impact used in this analysis, is provided in Part 3 and Section 1.1 of Schedule 1 of the plan.

**Table 1: Outputs of the dugong pressure analysis for the North-west Marine Region**

Pressure	Source	Rating
Sea level rise	Climate change	of potential concern
Changes in sea temperature	Climate change	of potential concern
Ocean acidification	Climate change	not of concern
Chemical pollution/contaminants	Shipping	not of concern
	Vessels (other)	not of concern
	Urban development	not of concern
	Agricultural activities	not of concern
Nutrient pollution	Onshore and offshore mining operations	of less concern
	Urban development	not of concern
Marine debris	Agricultural activities	not of concern
	Shipping	of potential concern
Noise pollution	Vessels (other)	of potential concern
	Fishing boats	of potential concern
	Land-based activities	of potential concern
	Seismic exploration	of less concern
Physical habitat modification	Shipping	of less concern
	Vessels	of less concern
	Onshore and offshore construction	of less concern
	Defence/surveillance activities	of less concern
Physical habitat modification	Dredging	of potential concern
	Dredge spoil	of potential concern
	Urban/coastal development	of less concern

**Legend**  of concern  of potential concern  of less concern  not of concern  data deficient or not assessed






**Table 1 continued: Outputs of the dugong pressure analysis for the North-west Marine Region**

Pressure	Source	Rating
Physical habitat modification	Fishing gear (active and derelict)	of less concern
	Storm events	of potential concern
Human presence at sensitive sites	Tourism	of less concern
	Recreation and charter fishing (burleying)	of less concern
	Research	of less concern
Extraction of living resources	Indigenous harvest	of potential concern
Bycatch	Commercial fishing (domestic)	of less concern
Oil pollution	Shipping	of less concern
	Vessels (other)	of less concern
	Oil rigs	of potential concern
Collision with vessels	Shipping	of potential concern
	Tourism	of potential concern
	Fishing	of potential concern
Disease	Shipping	data deficient or not assessed
	Fishing vessels	data deficient or not assessed
	Other vessels	data deficient or not assessed
Invasive species	Shipping	of potential concern
	Fishing vessels	of potential concern
	Vessels (other)	of potential concern

**Legend**       of concern       of potential concern       of less concern       not of concern       data deficient or not assessed



There is little information on the impacts of many of the pressures on dugongs in the North-west Marine Region, although there is information about impacts of pressures on dugongs elsewhere in Australia. In addition, the cumulative impacts of several pressures acting simultaneously is likely to be considerable but is poorly understood. The pressures identified as *of potential concern* are considered in two groups: pressures associated with habitat loss due to climate change and industrial developments; and pressures associated with human-induced mortality of dugongs.

### Pressures associated with habitat loss

Dugong habitat loss is likely to increase in the North-west Marine Region due to the large-scale industrial development occurring in the region and the impacts of climate change. The implications of climate change on seagrass distribution in the region are uncertain but could result in a decline in the extent and/or health of seagrass meadows. Thus pressures associated with both industrial development and climate change (and their cumulative impacts) are *of potential concern* for dugongs in the North-west Marine Region because they are likely to lead to the loss of dugong habitat and, consequently, population decline. Local seagrass loss can be very important for two reasons: there is some evidence that dugongs are faithful to specific areas learned from their mothers and are slow to recolonise other areas (Marsh et al. 2011), and the distribution of the dugong is typically fragmented. There is anecdotal evidence that some local areas may be staging posts between key habitats.

### Loss of dugong habitat associated with climate change

#### Sea level rise—climate change

Sea level has been rising at approximately 7.1 millimetres per year in the North-west Marine Region since the 1990s, the largest increase in Australia (National Tidal Centre 2010). Global sea levels have risen by 20 centimetres between 1870 and 2004 and predictions estimate a further rise of 5–15 centimetres by 2030, relative to 1990 levels (Church et al. 2009). Longer term predictions estimate increases of 0.5 to 1.0 metre by 2100, relative to 2000 levels (Climate Commission 2011). The main concern with sea level rise is the potential consequences when sea level rise is combined with increasing cyclone frequency (Climate Commission 2011; DCC 2009). Predictions for the North-west Marine Region in 2100 are for a hundred-fold increase in extreme sea level events in the Pilbara and up to a thousand-fold increase in the Kimberley (DCC 2009).

The likelihood of seagrass impacts due to climate change–induced sea level rise is linked to the location-specific interaction between climate, geomorphology and hydrology. The meso-tidal Pilbara Region may be more vulnerable to this pressure than the macro-tidal Kimberley (Semeniuk 1994; Woodroffe 1995), but this hypothesis remains untested. The resultant decrease in available light for seagrass meadows may lead to a reduction in growth and productivity of seagrass and the loss of seagrass in deeper waters. Sea level rise is also likely to lead to erosion of coastlines, which will increase the turbidity of coastal waters and affect survival of seagrasses. Although it is possible that new seagrass habitats may develop as low-lying coastal areas become intertidal, the overall effect of sea level rise on dugong habitats in the North-west Marine Region is uncertain and thus *of potential concern*.



### **Changes in sea temperature—climate change**

The likely increase in sea temperature associated with climate change is of *potential concern* for dugongs in the North-west Marine Region. Waycott et al. (2007) argue that elevated temperature will result in the greatest climate change impact on seagrasses, particularly in shallow habitats. Sea temperatures have warmed by 0.7 °C between 1910–1929 and 1989–2008, and current projections estimate ocean temperatures will be 1 °C warmer by 2030 (Lough 2009). Campbell et al. (2006) conducted temperature experiments on tropical seagrass species, including species eaten by dugongs, and demonstrated that the photosynthetic condition of all seagrass species tested was likely to suffer irreparable effects from short-term or episodic changes in seawater temperature (with temperatures reaching 40–45 °C). These acute stress responses of seagrasses to elevated seawater temperature are consistent with observed reductions in above-ground biomass during a recent El Niño event. In the Gulf of Carpentaria in 2002 seagrass loss and subsequent reports of ‘water fat’ (starving) dugongs were associated with elevated seawater temperature, although data are lacking to prove a causal link between these phenomena (Kwan & Bell 2003).

### **Physical habitat modification—storm events**

The likely increase in the intensity of storm events associated with climate change is of *potential concern* for dugongs and their habitats in the North-west Marine Region. Modelling predicts that climate change will result in increased intensity of storms and storm surges (Connolly 2009; Hyder Consulting 2008). Present indications are that modest to moderate increases (0–20 per cent) in average and maximum cyclone intensities are expected by the end of the century in some regions (Walsh & Ryan 2000). Evidence from various parts of northern Australia outside the North-west Marine Region points to episodic losses of hundreds of square kilometres of seagrass associated with extreme weather events such as cyclones and floods (Poiner & Peterkin 1996; Preen & Marsh 1995; Preen et al. 1995). Light availability for seagrass is typically significantly reduced after extreme weather events and deposited sediments can physically smother seagrass surfaces (Cabaço et al. 2008). In addition, storm surges can lead to dugongs being stranded above the high tide level (Marsh 1989).

## **Loss of dugong habitat associated with industrial development**

### **Physical habitat modification—dredging; dredge spoil**

The huge expansion in the offshore oil and gas and land-based mining industries in the North-west Marine Region and associated port and coastal development have the potential to impact dugong habitats because dredging and related activities, including spoil dumping, may degrade seagrass beds due to smothering and lack of light. There is little evidence of current substantial impact in the region but there is an established link between smothering, absence of light and seagrass decline (Cabaço et al. 2008).



### **Oil pollution—oil rigs**

Australia has a strong system for regulating industry activity that is the potential source of oil spills and this system has been strengthened further in response to the Montara oil spill. While oil spills are unpredictable events and their likelihood is low based on past experience, their consequences, especially for threatened species at important areas, could be severe. The coincidence of dugong habitats and extensive oil and gas exploration and production in the North-west Marine Region has the potential to adversely affect dugongs should a spill occur. Oil pollution may result in seagrass loss, although there is little evidence of a substantial impact on dugongs or their seagrass habitats from oil pollution from within the North-west Marine Region as yet. In addition, the dramatic effects of oil spills on seagrasses may not persist for long periods (Kenworthy et al. 1993).

### **Invasive species**


Asian bag or date mussel (*Musculista senhousia*) is a medium priority marine pest; that is, it has a reasonably high impact and/or invasion potential (Hayes et al. 2005). A review by Aquenal (2008) suggests that the potential for this species to become established in the North-west Marine Region is high. *Musculista* is transported in ballast water and as biofouling on vessel hulls. It tolerates low salinity and a wide range of temperatures and is present in parts of temperate southern Australia. It is present in the Leeuwin–Naturaliste coast of south-west Western Australia, including in Fremantle Harbour. *Musculista* invasion in the northern hemisphere has been linked with fragmentation of seagrass beds (Aquenal 2008). Shipping between the ports of Fremantle, north-western Australia and Asia is likely to increase in the future, thus increasing the potential for the mussel to be introduced into the North-west Marine Region (DEWHA 2008). There are reasonable grounds to predict that, if the pest is introduced, it has the potential to impact on seagrass habitats on which dugongs depend.

### **Pressures associated with human-induced mortality of dugongs**

Because of their life history characteristics and vulnerabilities, pressures that cause dugong mortality are *of potential concern* if such pressures occur over a wide geographic area, even if the magnitude of the pressures is uncertain.

### **Marine debris**

Debris harmful to marine wildlife includes plastics washed or blown from land into the sea, fishing gear abandoned by recreational and commercial fishers (ghost nets), and solid non-biodegradable floating materials (such as plastics) disposed of by ships at sea. The North-west Marine Region has increasing population, increasing shipping activity, fishing activities (commercial; recreational; traditional Indonesian; and illegal, unregulated and



unreported) and currents that bring debris from the populous Indonesian archipelago. All these factors increase the potential for marine debris to enter the region. Large amounts of fishing net are discarded or lost from the fisheries of the Arafura Sea (Limpus 2009). The amounts, characteristics and impacts of debris disposed of or lost overboard in the Arafura Sea are largely unknown (Kieessling 2003) and it is not known what proportion of such debris enters the North-west Marine Region. The established depositor beaches are mainly in the Northern Territory and the Gulf of Carpentaria coast, Queensland. This pressure is *of potential concern* because it is likely to cause injury or death to individual dugongs and there is inconclusive evidence about the adequacy of management measures to minimise the impact of marine debris on dugongs.

### ***Extraction of living resources—Indigenous harvest***

Indigenous communities and individuals with a native title right (determined or common law) to hunt dugongs may do so for their personal, domestic or non-commercial communal needs without a permit or licence. Indigenous harvest occurs in many communities adjacent to the North-west Marine Region. However, the level of harvest, and thus its sustainability, is unknown. Any impact is likely to be spatially variable. Dugongs are vulnerable to this pressure despite the recent investment in community ranger groups that are assuming responsibility for hunting management, and the Dugong and Marine Turtle Project coordinated by the North Australian Indigenous Land and Sea Management Alliance (NAILSMA 2010).

### ***Oil pollution—oil rigs***

Australia has a strong system for regulating industry activity that is the potential source of oil spills and this system is being strengthened further in response to the Montara oil spill. While oil spills are unpredictable events and their likelihood is low based on past experience, their consequences, especially for threatened species at important areas, could be severe. In the North-west Marine Region, large dugong populations coincide with oil and gas exploration and production activities. Oil pollution arising from these activities has the potential to result in dugong mortality. About 150 dugongs were estimated to have died in the Arabian Gulf during the Nowruz oil spill in 1983–1984 when more than one million barrels of oil flowed from seven wells damaged during the Iran–Iraq war (Preen et al. in press). There is little evidence of a substantial impact on dugongs in the North-west Marine Region to date. However, the species is vulnerable to oil pollution, the industry is expanding and there is potential for accidents to occur in areas used by dugong.



### ***Collisions with vessels—shipping; fishing; tourism***

The North-west Marine Region is experiencing significant growth in shipping and vessel traffic associated with increases in industrial development due to the resources boom. As a result, an increase in the human population with one of the highest levels of boat ownership per capita in Australia is occurring (Marsh et al. 2002). The Region and adjacent waters has a high level of shore-based tourist boat activity, including in the vicinity of seagrass beds in Ningaloo Marine Park, Shark Bay and Exmouth Gulf. All these changes have the potential to increase the risk to dugong of vessel strikes. Dugongs are killed accidentally when struck by boats and propellers when feeding in shallow inshore waters, particularly in areas where fast boats are used (Marsh et al. 2002). The relative contribution of vessels of different types to dugong mortality is not known and is likely to be area specific. The greatest danger appears to be in narrow channels used by boats and dugongs at low tide (Groom et al. 2004). Dugongs can become habituated to boat traffic, especially traffic concentrated around large seagrass meadows on which they feed. They have been known to continue to use an area even if they are continually disturbed by vessel traffic. There are anecdotal reports of dugongs being killed by vessel strike in the North-west Marine Region but evidence of impacts in the region is limited. The pressure is *of potential concern* because the dugong is vulnerable to the pressure, the pressure is increasing, and there is inadequate evidence of the effectiveness of management measures.



### 3. Relevant protection measures

The dugong is listed as a migratory and marine species under the EPBC Act. Under the Act, it is generally an offence to kill, injure, take, trade, keep or move listed marine, migratory or threatened species on Australian Government land or in Commonwealth waters without a permit.

Alongside the EPBC Act, a broad range of sector-specific management measures to address environmental issues and mitigate impacts apply to activities that take place in Commonwealth marine areas. These measures give effect to regulatory and administrative requirements under Commonwealth and state legislation for activities such as commercial and recreational fishing, oil and gas exploration and production, ports activities and maritime transport. In some instances, as in the case of shipping, these measures also fulfil Australia's obligations under a number of international conventions for the protection of the marine environment from pollution and environmental harm.

Australia has implemented a number of measures to protect dugongs, including spatial and seasonal commercial fisheries closures (aimed at reducing the bycatch of dugongs in gillnets and the physical damage of trawling on seagrass beds), and Indigenous management strategies for the protection of dugong habitat and populations. Evidence of the effectiveness of these management measures however, has not been documented. In addition, Australia is a signatory to several international agreements to protect dugongs

#### International measures

Australia is a signatory to the following international agreements for the conservation of dugongs:

- Convention on International Trade in Endangered Species (CITES)—[www.cites.org](http://www.cites.org)
- The Bonn Convention: Conservation of Migratory Species (CMS)—[www.cms.int](http://www.cms.int)
- CMS Memorandum of Understanding on the Conservation and Management of Dugongs and their Habitats Throughout their Range—[www.cms.int/species/dugong/index.htm](http://www.cms.int/species/dugong/index.htm).

For more information on conservation listings under the EPBC Act and related management objectives and protection measures, visit:

- [www.environment.gov.au/epbc/protect/migratory.html](http://www.environment.gov.au/epbc/protect/migratory.html)  
(listed migratory species)
- [www.environment.gov.au/coasts/species/marine-species-list.html](http://www.environment.gov.au/coasts/species/marine-species-list.html)  
(listed marine species)



# REFERENCES

Aquenal Pty Ltd 2008, *National control plan for the Asian bag or date mussel Musculista senhousia*, report prepared for the Australian Government.

Cabaço, S, Santos, R & Duarte, CM 2008, 'The impact of sediment burial and erosion on seagrasses: a review', *Estuarine Coastal and Shelf Science*, vol. 79, no. 3, pp. 354–366.

Campbell, SJ, McKenzie, LJ & Kerville, SP 2006, 'Photosynthetic responses of seven tropical seagrasses to elevated seawater temperature', *Journal of Experimental Marine Biology and Ecology*, vol. 330, pp. 455–468.

Church JA, White, NJ, Hunter, JR, McInnes, K & Mitchell, W 2009, 'Sea level', in ES Poloczanska, AJ Hobday & AJ Richardson (eds), *A marine climate change impacts and adaptation report card for Australia 2009*, National Climate Change Adaptation Research Facility.

Climate Commission 2011, *The critical decade: climate science, risks and responses*, Australian Government Department of Climate Change and Energy Efficiency, Canberra.

Connolly, R 2009, 'Seagrass', in ES Poloczanska, AJ Hobday & AJ Richardson (eds), *A marine climate change impacts and adaptation report card for Australia 2009*, National Climate Change Adaptation Research Facility publication 05/09, viewed 15 June 2011, <[www.oceanclimatechange.org.au/content/index.php/site/report\\_card\\_extended/category/seagrass/](http://www.oceanclimatechange.org.au/content/index.php/site/report_card_extended/category/seagrass/)>.


DCC (Australian Government Department of Climate Change) 2009, *Climate change risks to Australia's coasts: a first pass national assessment*, DCC, Canberra.

DEWHA (Australian Government Department of the Environment, Water, Heritage and the Arts) 2008, *The north-west marine bioregional plan: bioregional profile. A description of the ecosystems, conservation values and uses of the north-west marine bioregion*, DEWHA, Canberra.

Groom, R, Lawler, IR & Marsh, H 2004, 'The risk to dugongs of vessel strike in the southern bay island area of Moreton Bay', unpublished report to the Queensland Government Department of Environment by James Cook University.

Hayes, K, Sliwa, C, Migus, S, McEnnulty, F & Dunstan, P 2005, *National priority pests: part II – ranking of Australian marine pests*, an independent report undertaken for the Australian Government Department of Environment and Heritage by CSIRO Marine Research, viewed 20 July 2011, <[www.environment.gov.au/coasts/publications/imps/pubs/priority2.pdf](http://www.environment.gov.au/coasts/publications/imps/pubs/priority2.pdf)>.





Heinsohn, R, Lacy, RC, Lindenmayer, DB, Marsh, H, Kwan, D & Lawler, IR 2004, 'Unsustainable harvest of dugongs in Torres Strait and Cape York (Australia) waters: two case studies using population viability analysis', *Animal Conservation*, vol. 7, pp. 417–425.

Hyder Consulting Pty Ltd 2008, 'The impacts and management implications of climate change for the Australian Government's protected areas', unpublished report to the Australian Government Department of the Environment, Water, Heritage and the Arts and the Department of Climate Change, Canberra.

Kenworthy, WJ, Durako, MJ, Fatemy, SMR, Valavi, H & Thayer, GW 1993, 'Ecology of seagrasses in northeastern Saudi Arabia one year after the Gulf War oil spill', *Marine Pollution Bulletin*, vol. 27, pp. 213–222.

Kiessling, I 2003, *Finding solutions: derelict fishing gear and other marine debris in northern Australia*, National Oceans Office, Hobart, viewed 17 June 2011, <[www.environment.gov.au/coasts/mbp/publications/north/pubs/marine-debris-report.pdf](http://www.environment.gov.au/coasts/mbp/publications/north/pubs/marine-debris-report.pdf)>.

Kwan, D & Bell, I 2003, *Response to community concerns about green turtle, Chelonia mydas, and dugong, Dugong dugon, in waters adjacent to the Wellesley group of islands in the Gulf of Carpentaria*, report to the Mornington Shire Council, April 2003.

Limpus, CJ 2009, *A biological review of Australian marine turtles: 1. Loggerhead turtle Caretta caretta (Linnaeus)*, Queensland Environment Protection Agency.

Marsh, H 1989, 'Mass stranding of dugong by a tropical cyclone in northern Australia', *Marine Mammal Science*, vol. 5, no. 1, pp. 78–84.

Marsh, H & Kwan, D 2008, 'Temporal variability in the life history and reproductive biology of female dugongs in Torres Strait: the likely role of sea grass dieback', *Continental Shelf Research*, vol. 28, pp. 2152–2159.

Marsh, H, Heinsohn, GE & Marsh, LM 1984, 'Breeding cycle, life history and population dynamics of the dugong *Dugong dugon* (Sirenia: Dugongidae)', *Australian Journal of Zoology*, vol. 32, pp. 767–788.

Marsh H, Harris, ANM & Lawler, IR 1997, 'The sustainability of the Indigenous dugong fishery in Torres Strait, Australia/Papua New Guinea', *Conservation Biology*, vol. 11, pp. 1375–1386.

Marsh, H, Penrose, H, Eros, C & Hugues, J 2002, *Dugong status report and action plans for countries and territories*, United Nations Environment Programme, Nairobi, pp. 1–162.

Marsh, H, Lawler, I, Kwan, D, Delean, S, Pollock, K & Alldredge, M 2004, 'Aerial surveys and the potential biological removal technique indicate that the Torres Strait dugong fishery is unsustainable', *Animal Conservation*, vol. 7, pp. 1–9.



Marsh, H, O'Shea, TJ & Reynolds, JE III 2011, *The ecology and conservation of Sirenia: dugongs and manatees*, Cambridge University Press, United Kingdom, pp. 1–520.

NAILSMA (North Australian Indigenous Land and Sea Management Alliance) 2010, *Dugong and marine turtle project*, National Heritage Trust regional competitive component project final report, viewed 15 June 2011, <[www.nailsma.org.au/nailsma/publications/downloads/Final-Report-web.pdf](http://www.nailsma.org.au/nailsma/publications/downloads/Final-Report-web.pdf)>.

National Tidal Centre 2010, *The Australian baseline sea level monitoring project: annual sea level data summary report, July 2009–June 2010*, Australian Bureau of Meteorology, Canberra, viewed 20 June 2011, <[www.bom.gov.au/ntc/IDO60202/IDO60202.2010.pdf](http://www.bom.gov.au/ntc/IDO60202/IDO60202.2010.pdf)>.

Poiner, IR & Peterkin, C 1996, 'Seagrasses', in LP Zann & P Kailola (eds), *The state of the marine environment report for Australia, technical annex: 1*, Great Barrier Reef Marine Park Authority, Townsville, pp. 40–45.

Preen, AR & Marsh, H 1995, 'Response of dugongs to large scale loss of seagrass from Hervey Bay, Queensland', *Wildlife Research*, vol. 22, pp. 507–519.

Preen, AR, Lee Long, WJ & Coles, RG 1995, 'Flood and cyclone related loss, and partial recovery, of more than 1000 km<sup>2</sup> of seagrass in Hervey Bay, Queensland, Australia', *Aquatic Botany*, vol. 52, pp. 3–17.

Preen, A, Das, H, Al-Rumaidh, M & Hodgson, A in press, 'Dugongs in Arabia', in E Hines, J Reynolds, A Mignucci-Giannoni, L Aragones & M Marmontel (eds), *International strategies for manatee and dugong conservation*, University Press of Florida.

Semeniuk, V 1994, 'Predicting the effects of sea-level rise on mangroves in northwestern Australia', *Journal of Coastal Research*, vol. 10, pp. 1050–1076.

Walsh, KJE & Ryan, BF 2000, 'Tropical cyclone intensity increase near Australia as a result of climate change', *Journal of Climate*, vol. 13, pp. 3029–3036.

Waycott, M, Collier, C, McMahon, K, Ralph, P, McKenzie, L, Udy, J & Grech, A 2007, 'Vulnerability of seagrasses in the Great Barrier Reef to climate change', in JE Johnson & PA Marshall (eds), *Climate change and the Great Barrier Reef*, Great Barrier Reef Marine Park Authority and Australian Greenhouse Office, Australia, pp. 193–299.

Woodroffe, CD 1995, 'Response of tide-dominated mangrove shorelines in northern Australia to anticipated sea-level rise', *Earth Surface Processes and Landforms*, vol. 20, pp. 65–85.

