



Biodiversity Summary for NRM Regions Species List

What is the summary for and where does it come from?

This list has been produced by the Department of Sustainability, Environment, Water, Population and Communities (SEWPC) for the Natural Resource Management Spatial Information System.

The list was produced using the **Australian Natural Heritage Assessment Tool (ANHAT)**, which analyses data from a range of plant and animal surveys and collections from across Australia to automatically generate a report for each NRM region. Data sources (Appendix 2) include national and state herbaria, museums, state governments, CSIRO, Birds Australia and a range of surveys conducted by or for DEWHA.

For each family of plant and animal covered by ANHAT (Appendix 1), this document gives the number of species in the country and how many of them are found in the region. It also identifies species listed as Vulnerable, Critically Endangered, Endangered or Conservation Dependent under the EPBC Act.

A biodiversity summary for this region is also available.

For more information please see: www.environment.gov.au/heritage/anhat/index.html

Limitations

- ANHAT currently contains information on the distribution of over 30,000 Australian taxa. This includes all mammals, birds, reptiles, frogs and fish, 137 families of vascular plants (over 15,000 species) and a range of invertebrate groups. **Groups not yet covered in ANHAT are not included in the list.**
- The data used come from authoritative sources, but they are not perfect. All species names have been confirmed as valid species names, but it is not possible to confirm all species locations. The list summarises the input data, so errors in the data received would be reflected in this document.
- The scientific names and taxonomic concepts used in this report reflect an ANHAT view of the data and not necessarily that found in government censuses, databases or other authoritative lists.



Species List for NRM Region South, Tasmania

This list was produced by the Department of Sustainability, Environment, Water, Population and Communities using the Australian Natural Heritage Assessment Tool (ANHAT). ANHAT is a database and decision support tool which uses over 33 million specimen and survey records for vertebrates, selected invertebrates and 137 vascular plant families (over 15,000 species). This species list does not include exotic or extinct species.

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.

| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|---|-------------|---|--------------------|
| Vertebrate fauna | | | | |
| Actinopterygii | | | | |
| Atherinidae | | | | |
| | Hardyheads, Silversides, Whitebait | | (1 out of 20 species in Australia) | |
| Atherinosoma microstoma | Smallmouth Hardyhead | | 2 | 2010 |
| Brachionichthyidae | | | | |
| | Handfishes | | (1 out of 2 species in Australia) | |
| Brachionichthys hirsutus | Spotted Handfish | | 100 | 1997 |
| Eleotridae | | | | |
| | Gudgeons, Sleepers | | (1 out of 39 species in Australia) | |
| Philypnodon grandiceps | Flathead Gudgeon | | <1 | 1999 |
| Galaxiidae | | | | |
| | Galaxias, Salamanderfish | | (15 out of 23 species in Australia) | |
| Galaxias auratus | Golden Galaxias | | 100 | 2000 |
| Galaxias brevipinnis | Climbing Galaxias | | 17 | 2004 |
| Galaxias fontanus | Swan Galaxias | | 55 | 1993 |
| Galaxias johnstoni | Clarence Galaxias | | 100 | 2001 |
| Galaxias maculatus | Common Galaxias | | 6 | 2000 |
| Galaxias parvus | Swamp Galaxias | | 100 | 2004 |
| Galaxias pedderensis | Pedder Galaxias | | 100 | 1998 |
| Galaxias tanycephalus | Saddled Galaxias | | 100 | 1998 |
| Galaxias truttaceus | Trout Galaxias | | 21 | 2003 |
| Lovettia sealii | Tasmanian Whitebait | | 27 | 1991 |
| Neochanna cleaveri | Tasmanian Mudfish | | 33 | 1995 |
| Paragalaxias dissimilis | Shannon Galaxias | | 100 | 2004 |
| Paragalaxias eleotroides | Great Lake Galaxias | | 100 | 2004 |
| Paragalaxias julianus | Julian Galaxias | | 100 | 1995 |
| Paragalaxias mesotes | Arthurs Galaxias | | 100 | 1980 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------|---|---|---|--------------------|
| Gobiidae | Gobies | (2 out of 99 species in Australia) | | |
| Afurcagobius tamarensis | Tamar Goby | | 3 | 2000 |
| Tasmanogobius gloveri | Glover's Tasmangoby | | 33 | 1974 |
| Labridae | Wrasses, Parrotfishes, Rockwhitings | (1 out of 2 species in Australia) | | |
| Neoodax balteatus | Little Weed Whiting | | 4 | 1985 |
| Mugilidae | Mulletts | (1 out of 18 species in Australia) | | |
| Aldrichetta forsteri | Yelloweye Mullet | | <1 | 2010 |
| Percichthyidae | Australian Freshwater Basses, Australian Freshwater Blackfish, Australian Freshwater Cods, Temperate Seabasses | (1 out of 17 species in Australia) | | |
| Gadopsis marmoratus | River Blackfish | | 2 | 1996 |
| Pleuronectidae | Right Handed Flatfishes, Righteye Flounders | (1 out of 3 species in Australia) | | |
| Ammotretis rostratus | Longsnout Flounder | | 2 | 1978 |
| Retropinnidae | Southern Graylings, Southern Smelts | (2 out of 3 species in Australia) | | |
| Prototroctes maraena | Australian Grayling | | 8 | 1988 |
| Retropinna tasmanica | Tasmanian Smelt | | 54 | 1997 |
| Sparidae | Breams, Porgies, Seabreams | (1 out of 5 species in Australia) | | |
| Acanthopagrus butcheri | Black Bream | | 2 | 1997 |
| Syngnathidae | Pipefishes, Seahorses | (1 out of 9 species in Australia) | | |
| Stigmatopora nigra | Widebody Pipefish | | 2 | 2006 |
| Tetraodontidae | Blowfishes, Pufferfishes, Puffers, Toadfishes, Tobys | (1 out of 18 species in Australia) | | |
| Tetractenos glaber | Smooth Toadfish | | 1 | 1982 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|---------------------------------------|-------------|---|--------------------|
| Amphibia | | | | |
| Hylidae | | | | |
| | Tree-Frogs | | (3 out of 81 species in Australia) | |
| Litoria burrowsae | Tasmanian Tree Frog | | 53 | 2002 |
| Litoria ewingii | Brown Tree Frog | | 10 | 2008 |
| Litoria raniformis | Southern Bell-frog | Vulnerable | 2 | 2009 |
| Myobatrachidae | | | | |
| | Myobatrachid or Southern Frogs | | (9 out of 121 species in Australia) | |
| Crinia nimbus | Moss Frog, Moss Froglet | | 100 | 2000 |
| Crinia signifera | Common Froglet | | 5 | 2009 |
| Crinia tasmaniensis | Tasmanian Froglet | | 28 | 2009 |
| Geocrinia laevis | Smooth Frog | | 8 | 1994 |
| Geocrinia victoriana | Eastern Smooth Frog | | <1 | |
| Limnodynastes dumerilii | Banjo Frog, Eastern Pobblebonk | | 3 | 2008 |
| Limnodynastes tasmaniensis | Spotted Marsh Frog | | 2 | 1997 |
| Pseudophryne bibroni | Bibron's Toadlet, Brown Toadlet | | <1 | |
| Pseudophryne semimarmorata | Southern Toadlet | | 10 | 1994 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|---------------------------|-------------|---|--------------------|
| Aves | | | | |
| Acanthizidae | | | | |
| Scrubwrens, Thornbills (14 out of 43 species in Australia) and Allies | | | | |
| Acanthiza apicalis | Inland Thornbill | | <1 | 1993 |
| Acanthiza chrysorrhoa | Yellow-rumped Thornbill | | 1 | 2002 |
| Acanthiza ewingii | Tasmanian Thornbill | | 37 | 2007 |
| Acanthiza iredalei | Slender-billed Thornbill | | <1 | 1989 |
| Acanthiza lineata | Striated Thornbill | | <1 | 1994 |
| Acanthiza nana | Yellow Thornbill | | <1 | 1966 |
| Acanthiza pusilla | Brown Thornbill | | 4 | 2003 |
| Acanthornis magnus | Scrubtit | | 39 | 2002 |
| Calamanthus fuliginosus | Striated Fieldwren | | 14 | 2002 |
| Dasyornis brachypterus | Eastern Bristlebird | Endangered | 1 | 1994 |
| Gerygone fusca | Western Gerygone | | <1 | 1991 |
| Gerygone levigaster | Mangrove Gerygone | | <1 | 1994 |
| Sericornis frontalis | White-browed Scrubwren | | 3 | 2004 |
| Smicronis brevirostris | Weebill | | <1 | 1991 |
| Accipitridae | | | | |
| Eagles, Hawks and Kites (8 out of 19 species in Australia) | | | | |
| Accipiter cirrhocephalus | Collared Sparrowhawk | | 1 | 2002 |
| Accipiter fasciatus | Brown Goshawk | | 2 | 2008 |
| Accipiter novaehollandiae | Grey Goshawk | | 5 | 2009 |
| Aquila audax | Wedge-tailed Eagle | | 2 | 2009 |
| Circus approximans | Swamp Harrier | | 5 | 2008 |
| Haliaeetus leucogaster | White-bellied Sea-Eagle | | 3 | 2009 |
| Haliastur sphenurus | Whistling Kite | | <1 | 1992 |
| Milvus migrans | Black Kite | | <1 | 1995 |
| Aegothelidae | | | | |
| Owlet Nightjars (1 out of 1 species in Australia) | | | | |
| Aegothales cristatus | Australian Owlet-nightjar | | <1 | 2008 |
| Alaudidae | | | | |
| Larks (2 out of 2 species in Australia) | | | | |
| Alauda arvensis | Eurasian Skylark | | 7 | 2002 |
| Mirafra javanica | Horsfield's Bushlark | | <1 | 1990 |
| Alcedinidae | | | | |
| Kingfishers and Kookaburras (4 out of 10 species in Australia) | | | | |
| Alcedo azurea | Azure Kingfisher | | <1 | 2006 |
| Dacelo leachii | Blue-winged Kookaburra | | <1 | 1991 |
| Dacelo novaeguineae | Laughing Kookaburra | | 2 | 2007 |
| Todiramphus sanctus | Sacred Kingfisher | | <1 | 2001 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|--|--|---|--------------------|
| Anatidae | Swans, Ducks and Geese | (13 out of 20 species in Australia) | | |
| <i>Anas castanea</i> | Chestnut Teal | | 5 | 2002 |
| <i>Anas gracilis</i> | Grey Teal | | <1 | 2002 |
| <i>Anas rhynchotis</i> | Australasian Shoveler | | 2 | 2002 |
| <i>Anas superciliosa</i> | Pacific Black Duck | | 2 | 2004 |
| <i>Aythya australis</i> | Hardhead | | <1 | 2002 |
| <i>Biziura lobata</i> | Musk Duck | | 3 | 2002 |
| <i>Cereopsis novaehollandiae</i> | Cape Barren Goose | | 5 | 2007 |
| <i>Chenonetta jubata</i> | Australian Wood Duck | | <1 | 2008 |
| <i>Cygnus atratus</i> | Black Swan | | 3 | 2008 |
| <i>Dendrocygna eytoni</i> | Plumed Whistling-Duck | | <1 | 1972 |
| <i>Oxyura australis</i> | Blue-billed Duck | | 2 | 2002 |
| <i>Stictonetta naevosa</i> | Freckled Duck | | <1 | 1998 |
| <i>Tadorna tadornoides</i> | Australian Shelduck | | 2 | 2002 |
| Anhingidae | Darter | (1 out of 1 species in Australia) | | |
| <i>Anhinga melanogaster</i> | Darter | | <1 | 1955 |
| Apodidae | Swifts and Swiftlets | (2 out of 4 species in Australia) | | |
| <i>Apus pacificus</i> | Fork-tailed Swift | | <1 | 1981 |
| <i>Hirundapus caudacutus</i> | White-throated Needletail | | 2 | 2004 |
| Ardeidae | Hérons, Egrets and Bitterns | (9 out of 14 species in Australia) | | |
| <i>Ardea alba</i> | Great Egret | | 1 | 2002 |
| <i>Ardea intermedia</i> | Intermediate Egret | | <1 | 1980 |
| <i>Ardea pacifica</i> | White-necked Heron | | <1 | 2001 |
| <i>Botaurus poiciloptilus</i> | Australasian Bittern | | 3 | 1988 |
| <i>Bubulcus ibis</i> | Cattle Egret | | 3 | 2007 |
| <i>Egretta garzetta</i> | Little Egret | | <1 | 2001 |
| <i>Egretta novaehollandiae</i> | White-faced Heron | | 1 | 2008 |
| <i>Egretta sacra</i> | Eastern Reef Egret | | <1 | 1975 |
| <i>Nycticorax caledonicus</i> | Nankeen Night Heron | | <1 | 1981 |
| Artamidae | Butcherbirds, Currawongs and Woodswallows | (7 out of 15 species in Australia) | | |
| <i>Artamus cyanopterus</i> | Dusky Woodswallow | | 2 | 2002 |
| <i>Cracticus quoyi</i> | Black Butcherbird | | <1 | 1975 |
| <i>Cracticus torquatus</i> | Grey Butcherbird | | 1 | 2004 |
| <i>Gymnorhina tibicen</i> | Australian Magpie | | <1 | 2003 |
| <i>Strepera fuliginosa</i> | Black Currawong | | 38 | 2007 |
| <i>Strepera graculina</i> | Pied Currawong | | <1 | 1992 |
| <i>Strepera versicolor</i> | Grey Currawong | | 3 | 2007 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------|--|--|---|--------------------|
| Burhinidae | Thicknees (Stone-Curlews) | (1 out of 2 species in Australia) | | |
| Burhinus grallarius | Bush Stone-curlew | | <1 | 1995 |
| Cacatuidae | Cockatoos | (7 out of 14 species in Australia) | | |
| Cacatua galerita | Sulphur-crested Cockatoo | | 1 | 2003 |
| Cacatua tenuirostris | Long-billed Corella | | <1 | 2001 |
| Callocephalon fimbriatum | Gang-gang Cockatoo | | <1 | 1995 |
| Calyptorhynchus funereus | Yellow-tailed Black-Cockatoo | | 6 | 2009 |
| Calyptorhynchus lathami | Glossy Black-Cockatoo | | <1 | 1988 |
| Eolophus roseicapillus | Galah | | <1 | 2002 |
| Nymphicus hollandicus | Cockatiel | | <1 | 1994 |
| Campephagidae | Cuckoo-shrikes, Trillers and Minivets | (2 out of 8 species in Australia) | | |
| Coracina novaehollandiae | Black-faced Cuckoo-shrike | | 1 | 2007 |
| Lalage tricolor | White-winged Triller | | <1 | 1977 |
| Casuariidae | Cassowary and Emus | (1 out of 2 species in Australia) | | |
| Dromaius novaehollandiae | Emu | | <1 | 1997 |
| Charadriidae | Plovers, Dotterels and Lapwings | (13 out of 15 species in Australia) | | |
| Charadrius bicinctus | Double-banded Plover | | 6 | 2002 |
| Charadrius dubius | Little Ringed Plover | | 6 | 1982 |
| Charadrius leschenaultii | Greater Sand Plover | | <1 | 1981 |
| Charadrius mongolus | Lesser Sand Plover | | <1 | 1981 |
| Charadrius ruficapillus | Red-capped Plover | | 2 | 2002 |
| Charadrius veredus | Oriental Plover | | <1 | 1978 |
| Elseyonis melanops | Black-fronted Dotterel | | <1 | 2002 |
| Erythrogonys cinctus | Red-kneed Dotterel | | <1 | 1987 |
| Pluvialis fulva | Pacific Golden Plover | | 2 | 2001 |
| Pluvialis squatarola | Grey Plover | | 1 | 2001 |
| Thinornis rubricollis | Hooded dotterel | | 9 | 2008 |
| Vanellus miles | Masked Lapwing | | 2 | 2007 |
| Vanellus tricolor | Banded Lapwing | | 1 | 2001 |
| Climacteridae | Treecreepers | (1 out of 6 species in Australia) | | |
| Cormobates leucophaeus | White-throated Treecreeper | | <1 | 1994 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|--|---|---|--------------------|
| Columbidae | Pigeons and Doves | (5 out of 23 species in Australia) | | |
| <i>Geopelia humeralis</i> | Bar-shouldered Dove | | <1 | 1993 |
| <i>Lopholaimus antarcticus</i> | Topknot Pigeon | | <1 | 1952 |
| <i>Phaps chalcoptera</i> | Common Bronzewing | | 1 | 2001 |
| <i>Phaps elegans</i> | Brush Bronzewing | | 5 | 2002 |
| <i>Phaps histrionica</i> | Flock Bronzewing | | <1 | 1973 |
| Corvidae | Crows and Jays | (3 out of 5 species in Australia) | | |
| <i>Corvus coronoides</i> | Australian Raven | | <1 | 1995 |
| <i>Corvus mellori</i> | Little Raven | | <1 | 1993 |
| <i>Corvus tasmanicus</i> | Forest Raven | | 23 | 2007 |
| Cuculidae | Cuckoos | (7 out of 12 species in Australia) | | |
| <i>Cacomantis flabelliformis</i> | Fan-tailed Cuckoo | | 3 | 2007 |
| <i>Cacomantis pallidus</i> | Pallid Cuckoo | | 2 | 2004 |
| <i>Cacomantis variolosus</i> | Brush Cuckoo | | <1 | 1991 |
| <i>Chrysococcyx basalis</i> | Horsfield's Bronze-Cuckoo | | <1 | 2002 |
| <i>Chrysococcyx lucidus</i> | Shining Bronze-Cuckoo | | 3 | 2006 |
| <i>Chrysococcyx minutillus</i> | Little Bronze-Cuckoo | | <1 | 1991 |
| <i>Scythrops novaehollandiae</i> | Channel-billed Cuckoo | | <1 | 1967 |
| Dicaeidae | Flower-peckers | (1 out of 2 species in Australia) | | |
| <i>Dicaeum hirundinaceum</i> | Mistletoebird | | <1 | 1990 |
| Dicruridae | Fan-tails, Drongos and Monarchs | (7 out of 23 species in Australia) | | |
| <i>Dicrurus bracteatus</i> | Spangled Drongo | | <1 | 1978 |
| <i>Grallina cyanoleuca</i> | Magpie-lark | | <1 | 1994 |
| <i>Myiagra cyanoleuca</i> | Satin Flycatcher | | 6 | 2002 |
| <i>Myiagra rubecula</i> | Leaden Flycatcher | | <1 | 1968 |
| <i>Rhipidura albiscapa</i> | Grey Fantail | | <1 | 2009 |
| <i>Rhipidura fuliginosa</i> | New Zealand Fantail | | 2 | 2003 |
| <i>Rhipidura leucophrys</i> | Willie Wagtail | | <1 | 1994 |
| Diomedeidae | Albatrosses | (4 out of 5 species in Australia) | | |
| <i>Diomedea exulans</i> | Wandering Albatross | Vulnerable | 3 | 2004 |
| <i>Phoebastria palpebrata</i> | Light-mantled Sooty Albatross | | 2 | 2005 |
| <i>Thalassarche cauta</i> | Shy Albatross | | 12 | 2005 |
| <i>Thalassarche melanophris</i> | Black-browed Albatross | | 6 | 2005 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------|---|-------------|---|--------------------|
| Estrildidae | Grass Finches | | (5 out of 18 species in Australia) | |
| Lonchura castaneothorax | Chestnut-breasted Mannikin, Chestnut-bre | | <1 | 1991 |
| Neochmia temporalis | Red-browed Finch | | <1 | 1991 |
| Poephila cincta | Black-throated Finch | | <1 | 1993 |
| Stagonopleura bella | Beautiful Firetail | | 17 | 2007 |
| Taeniopygia guttata | Zebra Finch | | <1 | 1992 |
| Eupetidae | Whipbirds, Quail-thrushes and Jewel-babblers | | (2 out of 9 species in Australia) | |
| Cinclosoma castanotus | Chestnut Quail-thrush | | <1 | 1993 |
| Cinclosoma punctatum | Spotted Quail-thrush | | 3 | 2001 |
| Falconidae | Falcons and Kestrels | | (5 out of 6 species in Australia) | |
| Falco berigora | Brown Falcon | | 1 | 2008 |
| Falco cenchroides | Nankeen Kestrel | | <1 | 2001 |
| Falco longipennis | Australian Hobby | | <1 | 2008 |
| Falco peregrinus | Peregrine Falcon | | 3 | 2008 |
| Falco subniger | Black Falcon | | <1 | 1966 |
| Haematopodidae | Oystercatchers | | (2 out of 2 species in Australia) | |
| Haematopus fuliginosus | Sooty Oystercatcher | | 5 | 2002 |
| Haematopus longirostris | Pied Oystercatcher | | 5 | 2006 |
| Hirundinidae | Swallows and Martins | | (4 out of 6 species in Australia) | |
| Cheramoeca leucosternus | White-backed Swallow | | <1 | 1991 |
| Hirundo neoxena | Welcome Swallow | | 2 | 2004 |
| Petrochelidon ariel | Fairy Martin | | <1 | 1997 |
| Petrochelidon nigricans | Tree Martin | | 2 | 2002 |
| Hydrobatidae | Storm-petrels | | (2 out of 3 species in Australia) | |
| Garrodia nereis | Grey-backed Storm-Petrel | | 18 | 1976 |
| Pelagodroma marina | White-faced Storm-Petrel | | 4 | 1988 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|---|-------------|---|--------------------|
| Laridae | Gulls, Terns, Skuas and Jaegers | | (11 out of 18 species in Australia) | |
| <i>Chlidonias hybridus</i> | Whiskered Tern | | <1 | 1985 |
| <i>Chroicocephalus novaehollandiae</i> | Silver Gull | | 3 | 2005 |
| <i>Gelochelidon nilotica</i> | Gull-billed Tern | | <1 | 1991 |
| <i>Larus dominicanus</i> | Kelp Gull | | 29 | 2005 |
| <i>Larus pacificus</i> | Pacific Gull | | 9 | 2005 |
| <i>Onychoprion fuscata</i> | Sooty Tern | | <1 | 1990 |
| <i>Sterna albifrons</i> | Little Tern | | <1 | 2001 |
| <i>Sterna caspia</i> | Caspian Tern | | 2 | 2002 |
| <i>Sterna striata</i> | White-fronted Tern | | 3 | 1999 |
| <i>Sterna vittata</i> | Antarctic Tern | | 20 | 1965 |
| <i>Thalasseus bergii</i> | Crested Tern | | 3 | 2004 |
| Maluridae | Fairy-wrens and Grasswrens | | (3 out of 22 species in Australia) | |
| <i>Malurus cyaneus</i> | Superb Fairy-wren | | 3 | 2007 |
| <i>Malurus melanocephalus</i> | Red-backed Fairy-wren | | <1 | 1994 |
| <i>Stipiturus malachurus</i> | Southern Emu-wren | | 5 | 2001 |
| Meliphagidae | Honeyeaters and Australian Chats | | (23 out of 72 species in Australia) | |
| <i>Acanthorhynchus tenuirostris</i> | Eastern Spinebill | | 5 | 2007 |
| <i>Anthochaera carunculata</i> | Red Wattlebird | | <1 | 1988 |
| <i>Anthochaera chrysoptera</i> | Little Wattlebird | | 5 | 2002 |
| <i>Anthochaera paradoxa</i> | Yellow Wattlebird | | 40 | 2007 |
| <i>Epthianura albifrons</i> | White-fronted Chat | | 2 | 2002 |
| <i>Gliciphila melanops</i> | Tawny-crowned Honeyeater | | <1 | 2001 |
| <i>Grantiella picta</i> | Painted Honeyeater | | <1 | 1994 |
| <i>Lichenostomus chrysops</i> | Yellow-faced Honeyeater | | <1 | 1996 |
| <i>Lichenostomus flavicollis</i> | Yellow-throated Honeyeater | | 38 | 2007 |
| <i>Lichenostomus fuscus</i> | Fuscous Honeyeater | | <1 | 1986 |
| <i>Lichenostomus leucotis</i> | White-eared Honeyeater | | <1 | 1994 |
| <i>Lichenostomus melanops</i> | Yellow-tufted Honeyeater | | <1 | 1991 |
| <i>Lichenostomus penicillatus</i> | White-plumed Honeyeater | | <1 | 1990 |
| <i>Manorina flavigula</i> | Yellow-throated Miner | | <1 | 1991 |
| <i>Manorina melanocephala</i> | Noisy Miner | | 2 | 2003 |
| <i>Meliphaga lewinii</i> | Lewin's Honeyeater | | <1 | 1990 |
| <i>Melithreptus affinis</i> | Black-headed Honeyeater | | 44 | 2007 |
| <i>Melithreptus albogularis</i> | White-throated Honeyeater | | <1 | 1988 |
| <i>Melithreptus brevirostris</i> | Brown-headed Honeyeater | | <1 | 2003 |
| <i>Melithreptus validirostris</i> | Strong-billed Honeyeater | | 42 | 2003 |
| <i>Phylidonyris novaehollandiae</i> | New Holland Honeyeater | | 5 | 2007 |
| <i>Phylidonyris pyrrhoptera</i> | Crescent Honeyeater | | 14 | 2007 |
| <i>Plectorhyncha lanceolata</i> | Striped Honeyeater | | <1 | 1991 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------|--|---|---|--------------------|
| Menuridae | Lyrebirds | (1 out of 2 species in Australia) | | |
| <i>Menura novaehollandiae</i> | Superb Lyrebird | | 2 | 2007 |
| Motacillidae | Pipits and Wagtails | (1 out of 5 species in Australia) | | |
| <i>Anthus australis</i> | Australian Pipit | | 1 | 2002 |
| Muscicapidae | Old World Flycatchers and Thrushes | (1 out of 6 species in Australia) | | |
| <i>Zoothera lunulata</i> | Bassian Thrush | | 6 | 2002 |
| Pachycephalidae | Whistlers, Shrike-thrushes and Allies | (4 out of 16 species in Australia) | | |
| <i>Colluricincla harmonica</i> | Grey Shrike-thrush | | 1 | 2007 |
| <i>Pachycephala olivacea</i> | Olive Whistler | | 13 | 2004 |
| <i>Pachycephala pectoralis</i> | Golden Whistler | | 2 | 2007 |
| <i>Pachycephala rufiventris</i> | Rufous Whistler | | <1 | 1993 |
| Paradisaeidae | Birds-of-Paradise | (1 out of 4 species in Australia) | | |
| <i>Ptiloris paradisus</i> | Paradise Riflebird | | <1 | 1994 |
| Pardalotidae | Pardalotes | (3 out of 4 species in Australia) | | |
| <i>Pardalotus punctatus</i> | Spotted Pardalote | | 3 | 2007 |
| <i>Pardalotus quadragintus</i> | Forty-spotted Pardalote | Endangered | 74 | 2009 |
| <i>Pardalotus striatus</i> | Striated Pardalote | | 1 | 2007 |
| Pelecanidae | Pelicans | (1 out of 1 species in Australia) | | |
| <i>Pelecanus conspicillatus</i> | Australian Pelican | | <1 | 2002 |
| Petroicidae | Australasian Robins | (8 out of 23 species in Australia) | | |
| <i>Eopsaltria australis</i> | Eastern Yellow Robin | | <1 | 1990 |
| <i>Melanodryas cucullata</i> | Hooded Robin | | <1 | 1994 |
| <i>Melanodryas vittata</i> | Dusky Robin | | 38 | 2002 |
| <i>Petroica boodang</i> | Scarlet Robin | | 6 | 2000 |
| <i>Petroica multicolor</i> | Pacific Robin | | 6 | 2007 |
| <i>Petroica phoenicea</i> | Flame Robin | | 7 | 2002 |
| <i>Petroica rodinogaster</i> | Pink Robin | | 14 | 2002 |
| <i>Tregellasia capito</i> | Pale-yellow Robin | | <1 | 1988 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------------------|---|--|---|--------------------|
| Phalacrocoracidae | Cormorants and Shags | (6 out of 6 species in Australia) | | |
| <i>Leucocarbo atriceps</i> | Imperial Shag | | 17 | 2000 |
| <i>Microcarbo melanoleucos</i> | Little Pied Cormorant | | 2 | 2003 |
| <i>Phalacrocorax carbo</i> | Great Cormorant | | 3 | 2004 |
| <i>Phalacrocorax fuscescens</i> | Black-faced Cormorant | | 13 | 2005 |
| <i>Phalacrocorax sulcirostris</i> | Little Black Cormorant | | 1 | 2002 |
| <i>Phalacrocorax varius</i> | Pied Cormorant | | <1 | 2001 |
| Phasianidae | Quail, Fowl and Allies | (2 out of 3 species in Australia) | | |
| <i>Coturnix pectoralis</i> | Stubble Quail | | <1 | 2003 |
| <i>Coturnix ypsilophora</i> | Brown Quail | | 2 | 2002 |
| Podargidae | Frogmouths | (1 out of 3 species in Australia) | | |
| <i>Podargus strigoides</i> | Tawny Frogmouth | | 1 | 2004 |
| Podicipedidae | Grebes | (3 out of 3 species in Australia) | | |
| <i>Podiceps cristatus</i> | Great Crested Grebe | | 2 | 2002 |
| <i>Poliiocephalus poliocephalus</i> | Hoary-headed Grebe | | 3 | 2003 |
| <i>Tachybaptus novaehollandiae</i> | Australasian Grebe | | <1 | 2002 |
| Procellariidae | Fulmars, Petrels, Prions and Shearwaters | (18 out of 22 species in Australia) | | |
| <i>Ardenna bulleri</i> | Buller's Shearwater | | 4 | 1977 |
| <i>Ardenna carneipes</i> | Flesh-footed Shearwater | | <1 | 1980 |
| <i>Ardenna grisea</i> | Sooty Shearwater | | 9 | 2000 |
| <i>Ardenna pacificus</i> | Wedge-tailed Shearwater | | <1 | 1955 |
| <i>Ardenna tenuirostris</i> | Short-tailed Shearwater | | 10 | 2002 |
| <i>Daption capense</i> | Cape Petrel | | 2 | 1998 |
| <i>Halobaena caerulea</i> | Blue Petrel | Vulnerable | 3 | 1996 |
| <i>Macronectes giganteus</i> | Southern Giant-Petrel | Endangered | 1 | 2005 |
| <i>Macronectes halli</i> | Northern Giant-Petrel | Vulnerable | 7 | 2005 |
| <i>Pachyptila crassirostris</i> | Fulmar Prion | | 12 | 1975 |
| <i>Pachyptila desolata</i> | Antarctic Prion | | 2 | 2000 |
| <i>Pachyptila turtur</i> | Fairy Prion | Vulnerable | 12 | 2002 |
| <i>Pachyptila vittata</i> | Broad-billed Prion | | 9 | 1976 |
| <i>Pelecanoides urinatrix</i> | Common Diving-Petrel | | 21 | 2002 |
| <i>Pterodroma lessonii</i> | White-headed Petrel | | 3 | 1991 |
| <i>Pterodroma leucoptera</i> | Gould's Petrel | Endangered | 4 | 1992 |
| <i>Pterodroma macroptera</i> | Great-winged Petrel | | <1 | 1999 |
| <i>Puffinus assimilis</i> | Little Shearwater | | <1 | 1973 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|---|--|---|--------------------|
| Psittacidae | Parrots | (13 out of 39 species in Australia) | | |
| Barnardius zonarius | Australian Ringneck | | <1 | 1994 |
| Glossopsitta concinna | Musk Lorikeet | | 3 | 2002 |
| Glossopsitta pusilla | Little Lorikeet | | <1 | 1991 |
| Lathamus discolor | Swift Parrot | Endangered | 14 | 2008 |
| Neophema chrysogaster | Orange-bellied Parrot | Critically endangered | 14 | 2001 |
| Neophema chrysostoma | Blue-winged Parrot | | 7 | 2002 |
| Neophema elegans | Elegant Parrot | | <1 | 1991 |
| Neophema petrophila | Rock Parrot | | <1 | 1984 |
| Pezoporus wallicus | Ground Parrot | | 19 | 2001 |
| Platycercus caledonicus | Green Rosella | | 39 | 2007 |
| Platycercus elegans | Crimson Rosella | | <1 | 1995 |
| Platycercus eximius | Eastern Rosella | | 2 | 2004 |
| Trichoglossus haematodus | Rainbow Lorikeet | | <1 | 1991 |
| Rallidae | Crakes, Moorhens, Rails and Allies | (10 out of 16 species in Australia) | | |
| Fulica atra | Eurasian Coot | | 2 | 2002 |
| Gallinula mortierii | Tasmanian Native-hen | | 44 | 2008 |
| Gallinula tenebrosa | Dusky Moorhen | | <1 | 2000 |
| Gallinula ventralis | Black-tailed Native-hen | | <1 | 2001 |
| Gallirallus philippensis | Buff-banded Rail | | <1 | 1989 |
| Lewinia pectoralis | Lewin's Rail | | 5 | 2001 |
| Porphyrio porphyrio | Purple Swamphen | | 1 | 2001 |
| Porzana fluminea | Australian Spotted Crake | | <1 | 2001 |
| Porzana pusilla | Baillon's Crake | | <1 | 2001 |
| Porzana tabuensis | Spotless Crake | | 1 | 2002 |
| Recurvirostridae | Stilts and Avocet | (2 out of 3 species in Australia) | | |
| Cladorhynchus leucocephalus | Banded Stilt | | 1 | 2000 |
| Himantopus himantopus | Black-winged Stilt | | <1 | 2001 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|---|--|---|--------------------|
| Scolopacidae | Waders (Snipe, Godwits, Curlew, Sandpipers and Allies) | (21 out of 31 species in Australia) | | |
| <i>Actitis hypoleucos</i> | Common Sandpiper | | <1 | 1981 |
| <i>Arenaria interpres</i> | Ruddy Turnstone | | 1 | 2000 |
| <i>Calidris acuminata</i> | Sharp-tailed Sandpiper | | <1 | 2000 |
| <i>Calidris alba</i> | Sanderling | | <1 | 1999 |
| <i>Calidris canutus</i> | Red Knot | | 2 | 2000 |
| <i>Calidris ferruginea</i> | Curlew Sandpiper | | 1 | 2002 |
| <i>Calidris melanotos</i> | Pectoral Sandpiper | | 2 | 1985 |
| <i>Calidris ruficollis</i> | Red-necked Stint | | 2 | 2002 |
| <i>Calidris subminuta</i> | Long-toed Stint | | 1 | 1998 |
| <i>Calidris tenuirostris</i> | Great Knot | | <1 | 1980 |
| <i>Gallinago hardwickii</i> | Latham's Snipe | | 5 | 2004 |
| <i>Heteroscelus brevipes</i> | Grey-tailed Tattler | | 1 | 1988 |
| <i>Limicola falcinellus</i> | Broad-billed Sandpiper | | <1 | 1999 |
| <i>Limosa lapponica</i> | Bar-tailed Godwit | | 1 | 2002 |
| <i>Limosa limosa</i> | Black-tailed Godwit | | <1 | 1979 |
| <i>Numenius madagascariensis</i> | Eastern Curlew | | 2 | 2002 |
| <i>Numenius minutus</i> | Little Curlew | | <1 | 1975 |
| <i>Numenius phaeopus</i> | Whimbrel | | <1 | 2002 |
| <i>Philomachus pugnax</i> | Ruff | | <1 | 1965 |
| <i>Tringa glareola</i> | Wood Sandpiper | | <1 | 1967 |
| <i>Tringa nebularia</i> | Common Greenshank | | <1 | 2002 |
| Spheniscidae | Penguins | (6 out of 6 species in Australia) | | |
| <i>Aptenodytes patagonicus</i> | King Penguin | | 55 | 2003 |
| <i>Eudyptes chrysocome</i> | Rockhopper Penguin | | 7 | 1999 |
| <i>Eudyptes chrysolophus</i> | Macaroni Penguin | | 41 | 2001 |
| <i>Eudyptes schlegeli</i> | Royal Penguin | | 100 | 1984 |
| <i>Eudyptula minor</i> | Little Penguin | | 9 | 2003 |
| <i>Pygoscelis antarctica</i> | Chinstrap Penguin | | 43 | 1994 |
| Strigidae | Hawk-type Owls | (1 out of 5 species in Australia) | | |
| <i>Ninox novaeseelandiae</i> | Southern Boobook | | 2 | 2008 |
| Sturnidae | Starlings and Mynas | (1 out of 3 species in Australia) | | |
| <i>Aplonis metallica</i> | Metallic Starling | | <1 | 1994 |
| Sulidae | Boobies and Gannets | (1 out of 5 species in Australia) | | |
| <i>Morus serrator</i> | Australasian Gannet | | 9 | 2002 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------------|--------------------------------|--|---|--------------------|
| Sylviidae | Old World Warblers | (3 out of 9 species in Australia) | | |
| <i>Acrocephalus australis</i> | Australian Reed-Warbler | | <1 | 2001 |
| <i>Cincloramphus cruralis</i> | Brown Songlark | | <1 | 1993 |
| <i>Megalurus gramineus</i> | Little Grassbird | | 2 | 2001 |
| Threskiornithidae | Ibis and Spoonbills | (4 out of 5 species in Australia) | | |
| <i>Platalea flavipes</i> | Yellow-billed Spoonbill | | <1 | 1977 |
| <i>Platalea regia</i> | Royal Spoonbill | | <1 | 2001 |
| <i>Plegadis falcinellus</i> | Glossy Ibis | | <1 | 1999 |
| <i>Threskiornis molucca</i> | Australian White Ibis | | <1 | 1980 |
| Turnicidae | Button-quail | (2 out of 7 species in Australia) | | |
| <i>Turnix pyrrhothorax</i> | Red-chested Button-quail | | <1 | 1987 |
| <i>Turnix varia</i> | Painted Button-quail | | <1 | 2008 |
| Tytonidae | Barn-type Owls | (2 out of 6 species in Australia) | | |
| <i>Tyto alba</i> | Barn Owl | | <1 | 1980 |
| <i>Tyto novaehollandiae</i> | Masked Owl | | 8 | 2008 |
| Zosteropidae | White-eyes (Silvereyes) | (1 out of 6 species in Australia) | | |
| <i>Zosterops lateralis</i> | Silvereye | | 3 | 2007 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|--|---|---|--------------------|
| Mammalia | | | | |
| Burramyidae | | | | |
| | Pygmy-possums | (2 out of 5 species in Australia) | | |
| Cercartetus lepidus | Little pygmy-possum | | 16 | 2005 |
| Cercartetus nanus | Eastern pygmy-possum | | 3 | 2003 |
| Dasyuridae | | | | |
| | Dasyurids (Quolls, Antechinus, Dunnarts and Allies) | (7 out of 55 species in Australia) | | |
| Antechinus minimus | Swamp antechinus | | 13 | 2003 |
| Antechinus stuartii | Brown antechinus | | <1 | |
| Antechinus swainsonii | Dusky antechinus | | 3 | 2001 |
| Dasyurus maculatus | Spotted-tailed quoll | Vulnerable | 5 | 2008 |
| Dasyurus viverrinus | Eastern quoll | | 41 | 2009 |
| Sarcophilus harrisii | Tasmanian devil | Endangered | 39 | 2009 |
| Sminthopsis leucopus | White-footed dunnart | | 6 | 2009 |
| Macropodidae | | | | |
| | Wallabies, Kangaroos and Tree-kangaroos | (4 out of 41 species in Australia) | | |
| Macropus agilis | Agile wallaby | | <1 | 2009 |
| Macropus giganteus | Eastern grey kangaroo | | <1 | 1996 |
| Macropus rufogriseus | Red-necked wallaby | | 6 | 2008 |
| Thylogale billardieri | Tasmanian pademelon | | 39 | 2008 |
| Muridae | | | | |
| | Rats and Mice | (5 out of 60 species in Australia) | | |
| Hydromys chrysogaster | Water-rat | | 3 | 2009 |
| Mastacomys fuscus | Broad-toothed Rat | | 3 | 1999 |
| Pseudomys higginsii | Long-tailed mouse | | 47 | 2009 |
| Pseudomys novaehollandiae | New Holland mouse | Vulnerable | 2 | 1987 |
| Rattus lutreolus | Swamp Rat | | 5 | 2009 |
| Ornithorhynchidae | | | | |
| | Platypus | (1 out of 1 species in Australia) | | |
| Ornithorhynchus anatinus | Platypus | | 4 | 2009 |
| Otariidae | | | | |
| | Eared Seals | (3 out of 5 species in Australia) | | |
| Arctocephalus gazella | Antarctic fur-seal | | 60 | 1997 |
| Arctocephalus pusillus | Australo-African Fur-seal | | <1 | 1979 |
| Neophoca cinerea | Australian sea-lion | Vulnerable | <1 | 1979 |
| Peramelidae | | | | |
| | Bandicoots and Spiny Bandicoot | (3 out of 7 species in Australia) | | |
| Isodon obesulus | Southern brown bandicoot | Endangered | 14 | 2003 |
| Perameles gunnii | Eastern barred bandicoot | Endangered | 42 | 2009 |
| Perameles nasuta | Long-nosed bandicoot | | <1 | |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|--|--|---|--------------------|
| Petauridae | Wrist-winged Gliders, Striped Possum and Leadbeaters Possum | (1 out of 6 species in Australia) | | |
| <i>Petaurus breviceps</i> | Sugar glider | | 1 | 2008 |
| Phalangeridae | Brushtail Possums, Cuscuses and Scaly-tailed Possum | (1 out of 6 species in Australia) | | |
| <i>Trichosurus vulpecula</i> | Common brushtail possum | | 3 | 2008 |
| Phocidae | True Seals | (2 out of 4 species in Australia) | | |
| <i>Hydrurga leptonyx</i> | Leopard seal | | 13 | 1997 |
| <i>Mirounga leonina</i> | Southern elephant seal | Vulnerable | 22 | 2001 |
| Potoroidae | Potoroos, Bettongs and the Musky Rat Kangaroo | (3 out of 11 species in Australia) | | |
| <i>Aepyprymnus rufescens</i> | Rufous bettong | | <1 | |
| <i>Bettongia gaimardi</i> | Tasmanian bettong | | 50 | 2009 |
| <i>Potorous tridactylus</i> | Long-nosed potoroo | | 3 | 2008 |
| Pseudocheiridae | Ring-tailed Possums and Greater Glider | (1 out of 7 species in Australia) | | |
| <i>Pseudocheirus peregrinus</i> | Common ringtail possum | | 4 | 2008 |
| Tachyglossidae | Echidna | (1 out of 1 species in Australia) | | |
| <i>Tachyglossus aculeatus</i> | Short-beaked Echidna | | 2 | 2009 |
| Vespertilionidae | Vespertilionid Bats | (10 out of 37 species in Australia) | | |
| <i>Chalinolobus gouldii</i> | Gould's wattled bat | | <1 | 1998 |
| <i>Chalinolobus morio</i> | Chocolate wattled bat | | <1 | 2000 |
| <i>Falsistrellus tasmaniensis</i> | Eastern false pipistrelle | | <1 | 2000 |
| <i>Nyctophilus geoffroyi</i> | Lesser long-eared bat | | <1 | 2008 |
| <i>Nyctophilus gouldi</i> | Gould's long-eared bat | | <1 | 1990 |
| <i>Nyctophilus timoriensis</i> | Greater Long-eared Bat | | <1 | 1986 |
| <i>Vespadelus darlingtoni</i> | Large Forest Bat | | <1 | 2008 |
| <i>Vespadelus pumilus</i> | Eastern Forest Bat | | <1 | |
| <i>Vespadelus regulus</i> | Southern Forest Bat | | <1 | 2008 |
| <i>Vespadelus vulturnus</i> | Little Forest Bat | | <1 | 2000 |
| Vombatidae | | (1 out of 3 species in Australia) | | |
| <i>Vombatus ursinus</i> | Common wombat | | 3 | 2004 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|------------------------------|---|---|--------------------|
| Petromyzontida | | | | |
| Petromyzontidae | | | | |
| | Lampreys | (2 out of 3 species in Australia) | | |
| Geotria australis | Pouch Lamprey | | 15 | 2009 |
| Mordacia mordax | Shorthead Lamprey | | 3 | 1983 |
| Reptilia | | | | |
| Agamidae | | | | |
| | Dragon lizards | (1 out of 70 species in Australia) | | |
| Rankinia diemensis | Mountain Heath Dragon | | 3 | 1994 |
| Chelidae | | | | |
| | Side-necked Tortoises | (1 out of 26 species in Australia) | | |
| Chelodina longicollis | Long-necked Turtle | | <1 | 1998 |
| Elapidae | | | | |
| | Front fang snakes | (3 out of 90 species in Australia) | | |
| Austrelaps superbus | Lowland Copperhead | | 3 | 2007 |
| Drysdalia coronoides | White-lipped Snake | | 8 | 2008 |
| Notechis scutatus | Tiger Snake | | 5 | 2008 |
| Scincidae | | | | |
| | Skinks | (18 out of 393 species in Australia) | | |
| Acritoscincus duperreyi | Eastern Three-lined Skink | | 3 | 1995 |
| Cyclodomorphus casuarinae | She-oak skink | | 45 | 2008 |
| Lampropholis delicata | Garden Sun-skink | | <1 | 2008 |
| Lampropholis guichenoti | Guichenots Sun-skink | | <1 | |
| Liopholis whitii | White's Skink | | 2 | 1996 |
| Nannoscincus maccoyi | McCoy's Skink | | <1 | |
| Niveoscincus coventryi | Snow-skink | | <1 | |
| Niveoscincus greeni | Northern Snow-skink | | 57 | 1994 |
| Niveoscincus metallicus | Metallic Snow-skink | | 25 | 2008 |
| Niveoscincus microlepidotus | Southern Snow-skink | | 72 | 1999 |
| Niveoscincus ocellatus | Ocellated Snow-skink | | 50 | 1996 |
| Niveoscincus orocryptus | Mountain Snow-skink | | 86 | 1999 |
| Niveoscincus palfreymani | Pedra Branca Snow-skink | Vulnerable | 100 | 2001 |
| Niveoscincus pretiosa | Tasmanian Tree Skink | | 42 | 1999 |
| Pseudemoia entrecasteauxii | Southern Grass Skink | | 4 | 2008 |
| Pseudemoia pagenstecheri | Tussock Skink | | 4 | 2000 |
| Pseudemoia rawlinsoni | Glossy Grass Skink | | <1 | 2008 |
| Tiliqua nigrolutea | Blotched-Blue-Tongued Lizard | | 5 | 2008 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Invertebrate fauna | | | |
| Amphipoda | | | |
| Hyalidae (1 out of 4 species in Australia) | | | |
| Austrochiltonia australis | | 50 | 1993 |
| Neoniphargidae (3 out of 23 species in Australia) | | | |
| Neoniphargus alpinus | | 100 | 1989 |
| Neoniphargus exiguus | | 100 | 1989 |
| Tasniphargus tyleri | | 100 | |
| Paramelitidae (1 out of 43 species in Australia) | | | |
| Antipodeus niger | | 100 | 1989 |
| Talitridae sandfleas, sandhoppers, landhoppers (14 out of 29 species in Australia) | | | |
| Austrotroides leptomerus | | 92 | 1990 |
| Austrotroides longicornis | | 86 | 1988 |
| Austrotroides maritimus | | 75 | 1990 |
| Keratroides albidus | | 29 | 1987 |
| Keratroides angulosus | | 42 | 1993 |
| Keratroides rex | | 50 | 1990 |
| Keratroides vulgaris | | 62 | 1995 |
| Mysticotalitrus cryptus | | 91 | 1995 |
| Mysticotalitrus tasmaniae | | 92 | 1993 |
| Neorchestia plicibrancha | | 62 | 1993 |
| Orchestiella neambulans | | 75 | 1995 |
| Orchestiella quasimodo | | 58 | 1988 |
| Protorchestia lakei | | 100 | 1993 |
| Tasmanorchestia annulata | | 20 | 1979 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|------------------------------|-------------|---|--------------------|
| Araneae | | | | |
| Hexathelidae | | | | |
| Funnel-web Spiders (3 out of 75 species in Australia) | | | | |
| | Funnel-web Spider | | 64 | 1986 |
| Hadronyche venenata | Funnel-web Spider | | 64 | 1986 |
| Plesiothele fentoni | Lake Fenton Funnel-web | | 100 | 1995 |
| Teranodes montana | Mountain Funnel-web | | 16 | 1987 |
| Idiopidae | | | | |
| Trapdoor Spiders (4 out of 153 species in Australia) | | | | |
| Arbanitis annulipes | Tasmanian Trapdoor | | 91 | 1982 |
| Arbanitis tasmanica | Tasmanian Trapdoor | | 100 | 1992 |
| Misgolas crispus | Trapdoor Spider | | 60 | 1992 |
| Misgolas mestoni | Mestons Trapdoor | | 82 | 1990 |
| Lycosidae | | | | |
| Wolf Spiders (17 out of 149 species in Australia) | | | | |
| Artoria albopilata | | | 9 | 2001 |
| Artoria berenice | | | 2 | 1996 |
| Artoria lineata | | | <1 | 1998 |
| Artoria triangularis | | | 2 | 1999 |
| Artoria victoriensis | | | 1 | 2004 |
| Artoriopsis expolita | | | 5 | 1998 |
| Artoriopsis melissae | | | 2 | 1994 |
| Diahogna martensii | | | 6 | 1957 |
| Hogna subrufa | | | 24 | 1989 |
| Lycosa godeffroyi | | | 6 | 2004 |
| Lycosa leuckartii | | | <1 | 1966 |
| Tasmanicosa tasmanica | | | 33 | |
| Tetrallycosa oraria | | | 6 | 1999 |
| Trochosa tristicula | | | 3 | 2000 |
| Venatrix esposica | | | 8 | 1987 |
| Venatrix funesta | | | 22 | 2006 |
| Venatrix pictiventris | | | 4 | 2003 |
| Migidae | | | | |
| Brushless-legged Trapdoor Spiders (1 out of 17 species in Australia) | | | | |
| Migas nitens | Silver Migas Trapdoor | | 100 | 1987 |
| Nemesiidae | | | | |
| Funnel-web Spiders (2 out of 227 species in Australia) | | | | |
| Chenistonia trevallynia | Trevalleyn Funnel-web Spider | | 40 | 1960 |
| Stanwellia pexa | Tasmanian Funnel-web | | 31 | 1980 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------|--|--|---|--------------------|
| Sparassidae | Huntsmen Spiders (formerly Heteropodidae) | (5 out of 119 species in Australia) | | |
| Delena cancerides | Crab-like Huntsman Spider | | 2 | 2008 |
| Eodelena spenceri | Spencers Huntsman Spider | | 25 | 1996 |
| Neosparassus diana | Dianas Huntsman Spider | | 4 | |
| Neosparassus n50 | Huntsman Spider | | 100 | |
| Neosparassus patellatus | Huntsman Spider | | 17 | 2000 |
| Basommatophora | | | | |
| Amphibolidae | Supra-littoral Globose "Land" Snails | (1 out of 2 species in Australia) | | |
| Salinator solida | Supra-littoral Globose "Land" Snail | | 14 | 2000 |
| Ancylidae | Freshwater Limpets | (2 out of 3 species in Australia) | | |
| Ferrissia petterdi | Freshwater Limpet | | 2 | |
| Ferrissia tasmanicus | Freshwater Limpet | | 7 | 1995 |
| Glacidorbidae | Freshwater Snails | (8 out of 23 species in Australia) | | |
| Benthodorbis fultoni | Fultons Freshwater Snail | | 100 | 1981 |
| Benthodorbis pawpela | Freshwater Snail | | 100 | 1977 |
| Glacidorbis pawpela | Great Lake Freshwater Snail | | 100 | 1981 |
| Glacidorbis pedderi | Lake Pedder Freshwater Snail | | 80 | 1995 |
| Glacidorbis tasmanicus | Tasmanian Freshwater Snail | | 11 | 1987 |
| Striadorbis janetae | Janets Freshwater Snail | | 50 | 1982 |
| Striadorbis pedderi | Lake Pedder Freshwater Snail | | 71 | 1989 |
| Tasmodorbis punctatus | Freshwater Snail | | 20 | 1988 |
| Lymnaeidae | Eutrophic Freshwater Snails | (1 out of 4 species in Australia) | | |
| Austropeplea tormentosa | South-east Eutrophic Freshwater Snail | | 6 | 2003 |
| Planorbidae | Freshwater Snails | (10 out of 71 species in Australia) | | |
| Ancylastrum cumingianus | Cumingians Freshwater Snail | | 100 | |
| Ancylastrum irvinae | Irvins Freshwater Snail | | 100 | 1966 |
| Glyptophysa gibbosa | Freshwater Snail | | 6 | 2003 |
| Glyptophysa huonensis | Tasmanian Freshwater Snail | | 50 | |
| Gyraulus atkinsoni | Atkinsons Freshwater Snail | | 33 | 1982 |
| Gyraulus meridionalis | Freshwater Snail | | 11 | 1995 |
| Gyraulus scottianus | Freshwater Snail | | 3 | 1995 |
| Gyraulus tasmanicus | Southern Freshwater Snail | | 25 | 1993 |
| Helicorbis australiensis | Freshwater Snail | | 2 | |
| Isidorella newcombi | Newcombs Freshwater Snail | | 2 | 1995 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------|-----------------------------|---|---|--------------------|
| Bivalvia | | | | |
| Hyriidae | | | | |
| | Freshwater Mussels | (1 out of 21 species in Australia) | | |
| Velesunio moretonicus | South Esk Freshwater Mussel | | 38 | 1953 |
| Sphaeriidae | | | | |
| | Pea-shells | (1 out of 17 species in Australia) | | |
| Pisidium casertanum | Pea-shell | | 67 | 1995 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Caenogastropoda | | | |
| Hydrobiidae | | | |
| Freshwater Operculate (62 out of 330 species in Australia) | | | |
| Snails | | | |
| Austropyrgus colludens | | 54 | 1995 |
| Austropyrgus diemensis | | 10 | 1988 |
| Austropyrgus dyerianus | | 100 | 1995 |
| Austropyrgus elongatus | | 100 | 1995 |
| Austropyrgus gordonensis | | 22 | 1987 |
| Austropyrgus gunnii | | 3 | 1987 |
| Austropyrgus nanus | | 100 | 1995 |
| Austropyrgus niger | | 100 | 1995 |
| Austropyrgus pagodoides | | 100 | 1995 |
| Austropyrgus salvus | | 11 | 1988 |
| Austropyrgus simsonianus | | 97 | 1995 |
| Beddomeia tumida | | 100 | 1992 |
| Nanocochlea 1 | | 100 | 1990 |
| Nanocochlea 2 | | 100 | 1988 |
| Nanocochlea 20 | | 100 | 1990 |
| Nanocochlea 22 | | 100 | 1989 |
| Nanocochlea 24 | | 100 | 1989 |
| Nanocochlea 3 | | 100 | 1989 |
| Nanocochlea 44 | | 100 | 1986 |
| Nanocochlea monticola | | 100 | 1982 |
| Nanocochlea parva | | 11 | 1996 |
| Nanocochlea pupoides | | 100 | 1982 |
| Phrantela 13 | | 100 | 1996 |
| Phrantela 15 | | 100 | 1987 |
| Phrantela 18 | | 100 | 1972 |
| Phrantela 5 | | 100 | 1989 |
| Phrantela 8 | | 50 | 1989 |
| Phrantela daveyensis | | 27 | 1989 |
| Phrantela pupiformis | | 89 | 2005 |
| Phrantela warwicki | | 94 | 1999 |
| Ps 10 | | 100 | 1994 |
| Ps 11 | | 100 | 1994 |
| Ps 12 | | 100 | 1994 |
| Ps 13 | | 100 | 1994 |
| Ps 14 | | 100 | 1994 |
| Ps 16 | | 100 | 1998 |
| Ps 17 | | 100 | 1994 |
| Ps 22 | | 100 | 1989 |
| Ps 23 | | 100 | 1989 |
| Ps 24 | | 100 | 1998 |
| Ps 27 | | 100 | 1989 |
| Ps 29 | | 100 | 1984 |
| Ps 3 | | 100 | 1990 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|------------------------------|-------------------------|---|--------------------|
| Ps 32 | Cave Freshwater Snail | | 100 | 1989 |
| Ps 33 | Cave Freshwater Snail | | 100 | 1985 |
| Ps 34 | Cave Freshwater Snail | | 100 | 1990 |
| Ps 38 | Cave Freshwater Snail | | 100 | 1990 |
| Ps 39 | Cave Freshwater Snail | | 100 | 1990 |
| Ps 45 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 46 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 47 | Cave Freshwater Snail | | 100 | 1991 |
| Ps 48 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 49 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 51 | Cave Freshwater Snail | | 100 | 1998 |
| Ps 6 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 7 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 8 | Cave Freshwater Snail | | 100 | 1994 |
| Ps 9 | Cave Freshwater Snail | | 100 | 1994 |
| Pseudotricula 10 | Persephone Pot Cave Snail | | 100 | 1991 |
| Pseudotricula 11 | Cueva Blanca Cave Snail | | 100 | 1988 |
| Pseudotricula 13 | Precipitous Bluff Cave Snail | | 100 | 1990 |
| Pseudotricula eberhardi | Eberhards Cave Snail | | 100 | 1996 |
| Pomatiopsidae | | Salt-lake Snails | (1 out of 14 species in Australia) | |
| Coxiella striata | Salt-lake Snail | | 5 | 1970 |
| Chordeumatida | | | | |
| Metopidiotrichidae | | | (2 out of 18 species in Australia) | |
| Australeuma simile | | | 100 | 2000 |
| Reginaterreuma tarkinensis | | | 50 | 1987 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------|----------------------|-------------|---|--------------------|
| Coleoptera | | | | |
| Buprestidae | Jewel Beetles | | (2 out of 1204 species in Australia) | |
| Castiarina erythromelas | | | 100 | 2008 |
| Castiarina insculpta | | | 100 | 1965 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|-------------|---|--------------------|
| Carabidae | | | |
| Ground Beetles | | (71 out of 2305 species in Australia) | |
| Adelotopus haemorrhoidalis | | 20 | 1965 |
| Agonocheila antipodum | | 14 | |
| Agonocheila bimaculata | | 100 | |
| Agonocheila curtula | | 2 | 1965 |
| Amblystomus nigrinus | | 25 | 1990 |
| Amblystomus ovalis | | 33 | |
| Amblytelus brevis | | 18 | 1994 |
| Amblytelus curtus | | 2 | |
| Amblytelus placidus | | 100 | 1998 |
| Anomotarus illawarrae | | 25 | |
| Bembidion jacksoniense | | 4 | |
| Chylinus ater | | 68 | 2004 |
| Clivina quadratifrons | | 4 | |
| Clivina suturalis | | 15 | |
| Clivina vagans | | 100 | |
| Demetrida angustula | | 25 | |
| Demetrida infuscata | | 67 | |
| Demetrida lineata | | 17 | |
| Demetrida setosa | | 100 | 1954 |
| Dicrochile brevicollis | | 50 | |
| Dicrochile goryi | | 12 | 1995 |
| Goedetrechus mendumae | | 100 | 1996 |
| Goedetrechus parallelus | | 100 | 1996 |
| Homethes guttifer | | 33 | 1995 |
| Hypharpax australis | | 7 | |
| Idacarabus cordicollis | | 100 | 1996 |
| Idacarabus longicollis | | 100 | 1996 |
| Idacarabus troglodytes | | 100 | 1996 |
| Lestignathus cursor | | 11 | 1965 |
| Mecyclothorax ambiguus | | 26 | 1998 |
| Mecyclothorax punctipennis | | 17 | 1991 |
| Notagonum marginellum | | 33 | 1994 |
| Notonomus politulus | | 17 | 1992 |
| Notonomus tubercauda | | 23 | 1995 |
| Pentagonica vittipennis | | 20 | |
| Percodermus niger | | 100 | 1953 |
| Percosoma carenoides | | 100 | 1995 |
| Pericompsus australis | | 7 | |
| Philophloeus distinguendus | | 6 | |
| Philophloeus eucalypti | | 25 | |
| Philophloeus moestus | | 33 | |
| Plagiotelum opalescens | | 50 | |
| Promecoderus brunnicornis | | 46 | 1994 |
| Promecoderus cordicollis | | 50 | 1985 |
| Promecoderus curvipes | | 100 | 1995 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|---------------------------------------|-------------|---|--------------------|
| <i>Promecoderus gibbosus</i> | Ground Beetle | | 40 | |
| <i>Promecoderus ovicollis</i> | Ground Beetle | | 25 | |
| <i>Promecoderus tasmanicus</i> | Tasmanian Ground Beetle | | 100 | 1989 |
| <i>Prosopogmus monochrous</i> | Pterostichid Ground Beetle | | 12 | |
| <i>Prosopogmus oodiformis</i> | Pterostichid Ground Beetle | | 25 | |
| <i>Prosopogmus punctifer</i> | Pterostichid Ground Beetle | | 50 | 1994 |
| <i>Pseudoceneus sollicitus</i> | Pterostichid Ground Beetle | | 12 | 1995 |
| <i>Pterocyrtus rubescens</i> | Trechitid Ground Beetle | | 100 | |
| <i>Rhabdotus floridus</i> | Pterostichid Ground Beetle | | 100 | 1995 |
| <i>Rhabdotus reflexus</i> | Pterostichid Ground Beetle | | 38 | 2009 |
| <i>Rhytisternus cyathoderus</i> | Pterostichid Ground Beetle | | 5 | |
| <i>Sarthrocrepis benefica</i> | Masoreitid Ground Beetle | | 10 | |
| <i>Sarthrocrepis corticalis</i> | Masoreitid Ground Beetle | | 21 | |
| <i>Scaraphites rotundipennis</i> | Scaritid Ground Beetle | | 4 | |
| <i>Scopodes aterrimus</i> | Odacanthid Ground Beetle | | 50 | 1995 |
| <i>Scopodes boops</i> | Odacanthid Ground Beetle | | 40 | |
| <i>Scopodes sigillatus</i> | Odacanthid Ground Beetle | | 12 | 1995 |
| <i>Sloaneana tasmaniae</i> | Tasmanian Trechitid Ground Beetle | | 23 | 1995 |
| <i>Stichonotus piceus</i> | Ground Beetle | | 40 | 1956 |
| <i>Tasmanitachoides hobarti</i> | Hobart Trechitid Ground Beetle | | 100 | 1957 |
| <i>Tasmanorites flavipes</i> | Ida Bay Caves Trechitid Ground Beetle | | 100 | 1996 |
| <i>Tasmanorites intermedius</i> | Trechitid Ground Beetle | | 100 | 1998 |
| <i>Tasmanorites nitens</i> | Silver Trechitid Ground Beetle | | 100 | 1954 |
| <i>Tasmanorites tasmaniae</i> | Tasmanian Trechitid Ground Beetle | | 100 | 1998 |
| <i>Trechimorphus diemenensis</i> | Trechitid Ground Beetle | | 13 | 1994 |
| <i>Trigonothops pacifica</i> | Lepitid Ground Beetle | | 6 | |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------------|----------------------------------|-------------|---|--------------------|
| Dytiscidae | Dytiscidine Water Beetles | | (33 out of 302 species in Australia) | |
| <i>Allodessus bistrigatus</i> | Water Beetle | | 1 | 2000 |
| <i>Antiporus blakei</i> | Blakes Water Beetle | | 19 | 2000 |
| <i>Antiporus femoralis</i> | Water Beetle | | 8 | 2000 |
| <i>Chostonectes gigas</i> | Giant Water Beetle | | 8 | 2000 |
| <i>Copelatus australiae</i> | Australian Water Beetle | | 3 | 2000 |
| <i>Copelatus boulevardi</i> | Boulevards Water Beetle | | 33 | 2001 |
| <i>Hyderodes schuckardi</i> | Schuckards Water Beetle | | 7 | 1961 |
| <i>Hyphodrus elegans</i> | Elegant Water Beetle | | 1 | 1985 |
| <i>Lancetes lanceolatus</i> | Lanceolate Water Beetle | | 2 | 2000 |
| <i>Limbodessus amabilis</i> | Water Beetle | | 21 | 2003 |
| <i>Limbodessus gemellus</i> | Water Beetle | | 17 | 2002 |
| <i>Limbodessus praelargus</i> | Water Beetle | | 13 | 2000 |
| <i>Megaporus fischeri</i> | Fischers Water Beetle | | 20 | 2000 |
| <i>Megaporus gardnerii</i> | Gardners Water Beetle | | 11 | 2000 |
| <i>Megaporus hamatus</i> | Water Beetle | | 5 | 2000 |
| <i>Megaporus howittii</i> | Howitts Water Beetle | | 1 | 2000 |
| <i>Necterosoma penicillatum</i> | Water Beetle | | 5 | 2003 |
| <i>Necterosoma undecimlineatum</i> | Water Beetle | | 2 | 2000 |
| <i>Onychodrus scutellaris</i> | Water Beetle | | 2 | 2000 |
| <i>Platynectes bakewellii</i> | Bakewells Water Beetle | | 14 | 2000 |
| <i>Platynectes decempunctatus</i> | Water Beetle | | 1 | 2000 |
| <i>Platynectes octodecimmaculatus</i> | Eighteen-spotted Water Beetle | | 3 | 1982 |
| <i>Rhantus suturalis</i> | Water Beetle | | 2 | 2000 |
| <i>Sternopriscus alpinus</i> | Alpine Water Beetle | | 75 | 2000 |
| <i>Sternopriscus barbarae</i> | Barbaras Water Beetle | | 14 | 2000 |
| <i>Sternopriscus clavatus</i> | | | 15 | 2000 |
| <i>Sternopriscus hansardii</i> | Hansards Water Beetle | | 9 | 2000 |
| <i>Sternopriscus meadfootii</i> | Meadfoots Water Beetle | | 10 | 2000 |
| <i>Sternopriscus montanus</i> | Montane Water Beetle | | 100 | 2000 |
| <i>Sternopriscus multimaculatus</i> | Multi-spotted Water Beetle | | 2 | 2000 |
| <i>Sternopriscus tasmanicus</i> | Tasmans Water Beetle | | 12 | 2000 |
| <i>Sternopriscus weckwerthi</i> | Weckwerths Water Beetle | | 44 | 2000 |
| <i>Sternopriscus williamsi</i> | Williams Water Beetle | | 100 | 1963 |
| Collothecaceae | Sessile Wheel Animals | | (7 out of 15 species in Australia) | |
| <i>Collothea campanulata</i> | Sessile Wheel Animal | | 47 | 1989 |
| <i>Collothea libera</i> | Sessile Wheel Animal | | 100 | |
| <i>Collothea mutabilis</i> | Sessile Wheel Animal | | 17 | 1989 |
| <i>Collothea ornata</i> | Ornate Sessile Wheel Animal | | 8 | 1989 |
| <i>Collothea pelagica</i> | Deep Water Sessile Wheel Animal | | 32 | 1989 |
| <i>Collothea trilobata</i> | Sessile Wheel Animal | | 17 | |
| <i>Stephanoceros fimbriatus</i> | Sessile Wheel Animal | | 33 | |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Diptera | | | |
| Asilidae Robber Flies (2 out of 360 species in Australia) | | | |
| Colepia rufiventris | | 8 | 1979 |
| Daptolestes limbipennis | | 33 | |
| Therevidae Stiletto Flies (22 out of 252 species in Australia) | | | |
| Agapophytus quatiens | | 100 | 1998 |
| Anabarhynchus danielsi | | 12 | 1963 |
| Anabarhynchus dimidiatus | | 11 | |
| Anabarhynchus helvenacus | | 24 | 1992 |
| Anabarhynchus hyalipennis | | 4 | |
| Anabarhynchus lanatus | | 100 | 1992 |
| Anabarhynchus maritimus | | 10 | 1994 |
| Anabarhynchus montanus | | 13 | 1992 |
| Anabarhynchus nudifemoratus | | 57 | 1992 |
| Anabarhynchus passus | | 33 | 1962 |
| Anabarhynchus planifrons | | 10 | 1992 |
| Anabarhynchus plumbeoides | | 6 | 1988 |
| Anabarhynchus plumbeus | | 15 | 1992 |
| Anabarhynchus whitei | | 43 | 1992 |
| Bonjeania actuosa | | 25 | 1998 |
| Bonjeania nitidifrons | | 17 | |
| Bonjeania segnis | | 20 | 1951 |
| Johnmannia tasmanica | | 100 | 1963 |
| Laxotela whitei | | 50 | 1998 |
| Neodialineura nitens | | 100 | |
| Neodialineura saxatilis | | 20 | 1983 |
| Parapsilocephala bifasciata | | 100 | 1998 |
| Ephemeroptera | | | |
| Nesameletidae Mayflies (1 out of 1 species in Australia) | | | |
| Ameletoides lacusalbinae | | 67 | 1995 |
| Oniscigastridae Mayflies (1 out of 3 species in Australia) | | | |
| Tasmanophlebia lacustris | | 64 | 1995 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|---|--|---|--------------------|
| Eupulmonata | | | | |
| Bulimulidae | | | | |
| | Elongate High-spired Land Snails | (1 out of 115 species in Australia) | | |
| Bothriembryon tasmanicus | Tasman Land Snail | | 82 | 2005 |
| Caryodidae | | | | |
| | Giant Land Snails | (3 out of 18 species in Australia) | | |
| Anoglypta launcestonensis | Launceston Land Snail | | 6 | 1996 |
| Caryodes dufresnii | Dufreys Land Snail | | 49 | 2009 |
| Hedleyella falconeri | Falconers Land Snail | | <1 | |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------------------------|---|--------------------|
| Charopidae | | | |
| Flattened and Turbinate (43 out of 642 species in Australia) | | | |
| Land Snails | | | |
| Allocharopa barossa hill | | 100 | 2004 |
| Allocharopa kershawi | | 17 | 1992 |
| Allocharopa legrandi | | 78 | 2009 |
| Allocharopa macgregor | | 50 | 1999 |
| Allocharopa sandspit | | 50 | 2004 |
| Allocharopa wellington | | 100 | 2009 |
| B swallet | | 100 | 2000 |
| Bischoffena bischoffensis | | 17 | 1970 |
| Dentherona aff tasmaniae | | 100 | 1966 |
| Dentherona dispar | | 33 | 2009 |
| Dentherona subrugosa | | 20 | 2000 |
| Discocharopa mimosa | Land Snail | 32 | 2009 |
| Discocharopa vigens | Land Snail | 46 | 2003 |
| Elsothera a | | 50 | 1999 |
| Elsothera limula | | 3 | 1966 |
| Elsothera ricei | Rices Charopid Land Snail | 34 | 2004 |
| Elsothera sp. | | 7 | 2000 |
| Geminoropa a | | 33 | 1998 |
| Mulathena fordei | Fords Charopid Land Snail | 35 | 2009 |
| Nautiliropa omicron | Charopid Land Snail | 1 | |
| Oreomava johnstoni | Johnstons Charopid Land Snail | 8 | 2000 |
| Pernagera a | | 60 | 2001 |
| Pernagera albolabris | Land Snail | 100 | 1966 |
| Pernagera kingstonensis | Kingston Land Snail | 52 | 2009 |
| Pernagera officeri | Officers Land Snail | 18 | 2005 |
| Pernagera tamarensis | Tamar Land Snail | 8 | 1995 |
| Pernagera tasmaniae | Tasmanian Land Snail | 70 | 2009 |
| Pernagera waterfall | Waterfall Land Snail | 67 | 2004 |
| Planilaoma luckmanii | Luckmans Land Snail | 60 | 2005 |
| Roblinella aff. mathinnae | | 100 | 1966 |
| Roblinella agnewi | Agnews Land Snail | 100 | 2001 |
| Roblinella b | | 100 | 2001 |
| Roblinella curacoae | | 85 | 2004 |
| Roblinella gadensis | Gadens Land Snail | 43 | 2003 |
| Roblinella mathinnae | Mathins Land Snail | 25 | 1998 |
| Roblinella sugarloaf | Sugarloaf Land Snail | 100 | 2004 |
| Stenacapha a | | 4 | 1994 |
| Stenacapha c | | 91 | 2002 |
| Stenacapha hamiltoni | Hamiltons Charopid Land Snail | 25 | 2004 |
| Stenacapha vitriniformis | Land Snail | 75 | 2000 |
| Thryasona a | | 75 | 2002 |
| Thryasona aff. diemenensis | | 100 | 1966 |
| Thryasona diemenensis | Van Diemens Land Snail | 31 | 2004 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|--|-------------|---|--------------------|
| Cystopeltidae | Australian "Half" Slugs (2 out of 13 species in Australia) | | | |
| <i>Cystopelta bicolor</i> | Two-coloured Half-slug | | 39 | 2005 |
| <i>Cystopelta petterdi</i> | Petterds Half-slug | | 5 | 2000 |
| Ellobiidae | Primitive Supralittoral Pulmonate Snails (2 out of 42 species in Australia) | | | |
| <i>Marinula xanthostoma</i> | Supralittoral Snail | | 15 | 2004 |
| <i>Ophicardelus ornata</i> | Supralittoral Snail | | 5 | 1999 |
| Helicarionidae | Helicarionid Land Snails (2 out of 300 species in Australia) | | | |
| <i>Helicarion cuvieri</i> | Cuviers Helicarionid Land Snail | | 33 | 2004 |
| <i>Helicarion rubicundus</i> | Helicarionid Land Snail | | 100 | 2004 |
| Punctidae | Translucent Frosted Land Snails (11 out of 107 species in Australia) | | | |
| <i>Laomavix collisi</i> | Collis's Punctid Land Snail | | 28 | 2005 |
| <i>Magilaoma a</i> | Punctid Land Snail | | 100 | 2002 |
| <i>Magilaoma penolensis</i> | Punctid Land Snail | | 14 | 2005 |
| <i>Paralaoma a</i> | Punctid Land Snail | | 30 | 1975 |
| <i>Paralaoma b</i> | Punctid Land Snail | | 33 | 2000 |
| <i>Paralaoma caputspinulae</i> | Punctid Land Snail | | 9 | 2005 |
| <i>Pedicamista a</i> | Punctid Land Snail | | 33 | 2001 |
| <i>Pedicamista bull hill</i> | Land Snail | | 67 | 1992 |
| <i>Pedicamista coesus</i> | Punctid Land Snail | | 71 | 2004 |
| <i>Trocholaoma a</i> | Punctid Land Snail | | 100 | 2001 |
| <i>Trocholaoma parvissimia</i> | Punctid Land Snail | | 25 | 2005 |
| Pupillidae | Pupa-shaped Shelled Land Snails (1 out of 62 species in Australia) | | | |
| <i>Pupilla australis</i> | Land Snail | | 2 | 1994 |
| Rhytididae | Carnivorous Land Snails (9 out of 107 species in Australia) | | | |
| <i>Prolesophanta a</i> | Carnivorous Land Snail | | 100 | 2002 |
| <i>Prolesophanta dyeri</i> | Dyers Carnivorous Land Snail | | 5 | 1998 |
| <i>Prolesophanta nelsonensis</i> | Nelsons Carnivorous Land Snail | | 55 | 2005 |
| <i>Tasmaphena a</i> | Carnivorous Land Snail | | 20 | 2000 |
| <i>Tasmaphena ruga</i> | Carnivorous Land Snail | | 41 | 2004 |
| <i>Tasmaphena sinclairi</i> | Sinclairs Carnivorous Land Snail | | 58 | 2005 |
| <i>Victaphanta a</i> | Carnivorous Land Snail | | 100 | 2002 |
| <i>Victaphanta lampra</i> | Carnivorous Land Snail | | 7 | 2001 |
| <i>Victaphanta milligani</i> | Milligans Carnivorous Land Snail | | 13 | 2002 |
| Succineidae | Amber Land Snails (1 out of 25 species in Australia) | | | |
| <i>Succinea australis</i> | Australian Amber Land Snail | | 2 | 1970 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|--|-------------|---|--------------------|
| Flosculariaceae | | | | |
| Conochilidae | | | | |
| | Six-"armed" Wheel Animals | | (4 out of 5 species in Australia) | |
| Conochilus coenobasis | Six-"armed" Wheel Animal | | 67 | 1989 |
| Conochilus dossuarius | Six-"armed" Wheel Animal | | 16 | 1989 |
| Conochilus natans | Six-"armed" Wheel Animal | | 22 | 1989 |
| Conochilus unicornis | Unicorn Six-"armed" Wheel Animal | | 20 | 1989 |
| Flosculariidae | | | | |
| | Sessile-footed Wheel Animals | | (12 out of 43 species in Australia) | |
| Beauchampia crucigera | Beauchamps Sessile-footed Wheel Anima | | 10 | |
| Floscularia janus | Sessile-footed Wheel Animal | | 20 | |
| Floscularia melicerta | Sessile-footed Wheel Animal | | 33 | |
| Lacinularia flosculosa | Sessile-footed Wheel Animal | | 25 | 1989 |
| Limnias ceratophylli | Sessile-footed Wheel Animal | | 17 | |
| Limnias melicerta | Sessile-footed Wheel Animal | | 42 | |
| Ptygura barbata | Sessile-footed Wheel Animal | | 100 | |
| Ptygura beauchampi | Sessile-footed Wheel Animal | | 100 | 1989 |
| Ptygura crystallina | Crystal Sessile-footed Wheel Animal | | 25 | |
| Ptygura melicerta | Sessile-footed Wheel Animal | | 40 | 1989 |
| Ptygura tacita | Sessile-footed Wheel Animal | | 25 | |
| Sinantherina socialis | Social Sessile-footed Wheel Animal | | 12 | |
| Hexarthridae | | | | |
| | Gelatinous-colonial Wheel Animals | | (2 out of 8 species in Australia) | |
| Hexarthra fennica | Gelatinous-colonial Wheel Animal | | 10 | |
| Hexarthra mira | Miras Gelatinous-colonial Wheel Animal | | 24 | 1989 |
| Testudinellidae | | | | |
| | Free-swimming Round-shelled Wheel Animals | | (8 out of 24 species in Australia) | |
| Pompholyx complanata | Free-swimming Round-shelled Wheel Ani | | 9 | |
| Pompholyx sulcata | Free-swimming Round-shelled Wheel Ani | | 6 | |
| Testudinella ahlstromi | Free-swimming Round-shelled Wheel Ani | | 50 | |
| Testudinella amphora | Vase-shaped Free-swimming Round-shell | | 57 | 1989 |
| Testudinella husseyi | Husseys Free-swimming Round-shelled | | 50 | |
| Testudinella parva | Free-swimming Round-shelled Wheel Ani | | 33 | 1989 |
| Testudinella tasmaniensis | Tasmanian Free-swimming Round-shelled | | 75 | 1989 |
| Testudinella unicornuta | Unicorn Free-swimming Round-shelled W | | 86 | 1989 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------------|--|-------------|---|--------------------|
| Trochosphaeridae | Globular Wheel Animals (7 out of 17 species in Australia) | | | |
| <i>Filinia australiensis</i> | Australian Globular Wheel Animal | | 16 | |
| <i>Filinia grandis</i> | Grand Globular Wheel Animal | | 27 | |
| <i>Filinia longiseta</i> | Globular Wheel Animal | | 17 | |
| <i>Filinia opoliensis</i> | Globular Wheel Animal | | 5 | |
| <i>Filinia pejleri</i> | Globular Wheel Animal | | 15 | |
| <i>Filinia terminalis</i> | Globular Wheel Animal | | 4 | |
| <i>Horaella brehmi</i> | Brehms Globular Wheel Animal | | 43 | |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|-----------------------------|--|---|--------------------|
| Hemiptera | | | | |
| Acanthosomatidae | | | | |
| | Shield Bugs | (3 out of 44 species in Australia) | | |
| Eupolemus insularis | | | 12 | |
| Eupolemus tasmanicus | | | 15 | |
| Eupolemus virescens | | | 100 | |
| Aradidae | | | | |
| | Flat bugs, Bark bugs | (5 out of 166 species in Australia) | | |
| Aradus australis | | | 8 | |
| Brachyrhynchus wilsoni | | | 12 | |
| Calisius tasmanicus | | | 40 | |
| Neuroctenus woodwardi | | | 5 | |
| Prosypniestus nasutus | | | 17 | 1956 |
| Artheneidae | | | | |
| | | (1 out of 2 species in Australia) | | |
| Dilompus robustus | | | 7 | |
| Berytidae | | | | |
| | Stilt bugs | (1 out of 7 species in Australia) | | |
| Bezu maiponga | maiponga | | 50 | |
| Ceratocombidae | | | | |
| | | (1 out of 1 species in Australia) | | |
| Ceratocombus australiensis | | | 38 | 1990 |
| Cicadidae | | | | |
| | | (7 out of 233 species in Australia) | | |
| Cicadetta abdominalis | | | 6 | 1957 |
| Cicadetta torrida | | | 25 | 1977 |
| Diemeniana euronotiana | | | 4 | 1966 |
| Diemeniana hirsuta | | | 100 | 1971 |
| Diemeniana tillyardi | | | 89 | 1998 |
| Pauropsalta encaustica | Black Squeaker | | 100 | 1954 |
| Psaltoda moerens | Redeye | | 12 | 1956 |
| Coreidae | | | | |
| | | (4 out of 81 species in Australia) | | |
| Amorbus angustior | Red Gum-bush Bug | | 14 | |
| Amorbus obscuricornis | | | 100 | 1993 |
| Amorbus rubiginosus | | | 4 | |
| Gelonus tasmanicus | | | 8 | |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------------------|-----------------------|---|--------------------|
| Corixidae | Waterboatmen | (6 out of 31 species in Australia) | |
| <i>Agraptocorixa parvipunctata</i> | | <1 | |
| <i>Diaprepocoris barycephalus</i> | | 18 | 1966 |
| <i>Micronecta annae</i> | | 23 | |
| <i>Micronecta robusta</i> | | 50 | 1995 |
| <i>Sigara neboissi</i> | | 100 | 2000 |
| <i>Sigara tasmaniae</i> | | 100 | 1995 |
| Cryptorhamphidae | | (1 out of 4 species in Australia) | |
| <i>Cryptorhamphus orbus</i> | | 2 | |
| Cydnidae | Burrower Bugs | (1 out of 82 species in Australia) | |
| <i>Adrisa atra</i> | | 25 | |
| Cymidae | | (1 out of 10 species in Australia) | |
| <i>Ontiscus barberi</i> | | 40 | |
| Gerridae | Water-striders | (1 out of 35 species in Australia) | |
| <i>Rheumatometra philarete</i> | | 10 | 1995 |
| Idiostolidae | | (1 out of 3 species in Australia) | |
| <i>Trisecus pictus</i> | | 33 | |
| Lygaeidae | | (1 out of 80 species in Australia) | |
| <i>Nysius turneri</i> | Invermay Bug | 67 | |
| Mesoveliidae | | (1 out of 7 species in Australia) | |
| <i>Mesovelia hungerfordi</i> | | 2 | 1973 |
| Miridae | Miridae | (6 out of 189 species in Australia) | |
| <i>Calocoris norvegicus</i> | Potato Bug | 100 | 1973 |
| <i>Chaetodus longiceps</i> | | 33 | |
| <i>Porophoroptera excellens</i> | | 50 | |
| <i>Pseudopantilius australis</i> | | 4 | |
| <i>Sejanus vividus</i> | | 22 | |
| <i>Sidnia kinbergi</i> | Crop Mirid | 6 | 1973 |
| Naucoridae | | (1 out of 9 species in Australia) | |
| <i>Naucoris congrex</i> | | 8 | 2000 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|---------------------------|---|---|--------------------|
| Notonectidae | Backswimmers | (4 out of 39 species in Australia) | | |
| Anisops deanei | | | 1 | 1995 |
| Anisops gratus | | | 1 | 1995 |
| Anisops hackeri | | | 7 | |
| Enithares woodwardi | | | 6 | 2000 |
| Ochteridae | Velvety Shore Bugs | (1 out of 11 species in Australia) | | |
| Ochterus eurythorax | | | 6 | |
| Peloridiidae | Moss Bugs | (2 out of 8 species in Australia) | | |
| Hemiodoecellus fidelis | | | 100 | |
| Hemiodoecus leai | | | 50 | 1997 |
| Pentatomidae | Stinkbugs | (14 out of 361 species in Australia) | | |
| Agonoscelis rutila | Horehound Bug | | 2 | 1983 |
| Cuspicona simplex | Green Potato Bug | | 3 | |
| Diemenia immarginata | | | 10 | |
| Diemenia rubromarginata | | | 3 | 1975 |
| Mycoolona atricornis | | | 14 | |
| Notius depressus | | | 6 | |
| Ocirrhoe lutescens | | | 4 | |
| Omyta centrolineata | | | 3 | |
| Oncocoris geniculatus | | | 5 | |
| Oncocoris similis | | | 14 | |
| Platycoris rugosus | | | 21 | |
| Poecilotoma grandicornis | | | 6 | |
| Tholosanus proximus | | | 2 | |
| Tinganina dimorpha | | | 20 | |
| Psyllidae | | (3 out of 319 species in Australia) | | |
| Anoeconeossa assimilis | | | 100 | 1966 |
| Anoeconeossa copidiformis | | | 100 | 1966 |
| Platybrija maddeni | | | 100 | 1985 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|------------------------|---|--------------------|
| Reduviidae | Assassin Bugs | (15 out of 226 species in Australia) | |
| <i>Coranus trabeatus</i> | | 3 | |
| <i>Dicrotelus prolixus</i> | | 20 | |
| <i>Ectomocoris decoratus</i> | | 9 | |
| <i>Empicoris rubromaculatus</i> | | 7 | |
| <i>Gminatus australis</i> | | 2 | |
| <i>Leistarches serripes</i> | | 7 | |
| <i>Nyllius asperatus</i> | | 11 | |
| <i>Nyllius australicus</i> | | 8 | |
| <i>Peirates fuliginosus</i> | | 10 | |
| <i>Pristhesancus plagipennis</i> | Bee-Killer | 3 | |
| <i>Pseudobargyia addititia</i> | | 100 | |
| <i>Pseudobargyia iuncea</i> | | 33 | 2009 |
| <i>Ptilocnemus sidnicus</i> | | 20 | |
| <i>Pygolampis frenchi</i> | | 25 | |
| <i>Stenolemus edwardsii</i> | | 6 | |
| Rhyparochromidae | Seed Bugs | (11 out of 191 species in Australia) | |
| <i>Austroxestus australiensis</i> | | 12 | 1967 |
| <i>Brentiscerus australis</i> | | 3 | |
| <i>Euander lacertosus</i> | Strawberry Bug | 5 | |
| <i>Fontejus collaris</i> | | 25 | |
| <i>Geratarma tasmaniensis</i> | | 75 | 1957 |
| <i>Lamproceps australis</i> | | 10 | |
| <i>Myocara acuminatum</i> | | 67 | 1957 |
| <i>Paradrymus exilirostris</i> | | 7 | 1956 |
| <i>Plinthisus tasmaniensis</i> | | 100 | |
| <i>Plinthisus woodwardi</i> | | 5 | |
| <i>Udeocoris nigroaeneus</i> | | 14 | 1957 |
| Schizopteridae | | (2 out of 61 species in Australia) | |
| <i>Pateena elimata</i> | | 33 | |
| <i>Pateena polymitarior</i> | | 40 | |
| Scutelleridae | | (3 out of 22 species in Australia) | |
| <i>Choerocoris paganus</i> | | <1 | |
| <i>Morbora hirtula</i> | | 11 | |
| <i>Scutiphora pedicellata</i> | Metallic Shield Bug | 9 | |
| Tettigarctidae | | (1 out of 2 species in Australia) | |
| <i>Tettigarcta tomentosa</i> | Tasmanian Hairy Cicada | 40 | 1972 |
| Thaumastocoridae | | (1 out of 11 species in Australia) | |
| <i>Baclozygum depressum</i> | | 14 | |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------|--|--|---|--------------------|
| Tingidae | Lacebugs | (3 out of 145 species in Australia) | | |
| Epimixia vittata | | | 12 | |
| Froggattia olivinia | Olive Lace Bug | | 20 | |
| Oncophysa vesiculata | | | 14 | |
| Triozidae | | (1 out of 24 species in Australia) | | |
| Aacanthocnema dobsoni | | | 100 | 1997 |
| Veliidae | Small Water Striders, Riffle Bugs | (3 out of 53 species in Australia) | | |
| Drepanovelia dubia | | | 8 | 1976 |
| Microvelia oceanica | Water Bug | | 2 | 1995 |
| Microvelia peramoena | | | 3 | 1995 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------------|-------------|---|--------------------|
| Hymenoptera | | | |
| Apidae | | | |
| Social Bees | | (3 out of 195 species in Australia) | |
| Exoneura angophorae | | 14 | 1967 |
| Exoneura bicolor | | 7 | 1967 |
| Exoneura turneri | | 100 | 1972 |
| Colletidae | | | |
| Short-tongued Bees | | (13 out of 878 species in Australia) | |
| Callomelitta picta | | 20 | 2009 |
| Euryglossa nigrocaerulea | | 20 | 1971 |
| Euryglossina healesvillensis | | 25 | 1967 |
| Hylaeus nubilosus | | 50 | |
| Leioproctus amabilis | | 3 | |
| Leioproctus carinatus | | 25 | |
| Leioproctus clarki | | 1 | 1989 |
| Leioproctus leai | | 20 | 1989 |
| Leioproctus obscurus | | 29 | 1972 |
| Leioproctus plumosus | | 3 | 1971 |
| Leioproctus providellus | | 33 | 1965 |
| Leioproctus providus | | 71 | 1971 |
| Leioproctus versicolor | | 4 | 1965 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|-------------|---|--------------------|
| Formicidae | | | |
| Ants | | (35 out of 1944 species in Australia) | |
| <i>Amblyopone australis</i> | | 2 | 2005 |
| <i>Anonychomyrma nitidiceps</i> | | 33 | 1995 |
| <i>Camponotus novaehollandiae</i> | | 25 | |
| <i>Camponotus terebrans</i> | | 2 | 1962 |
| <i>Colobostruma alinodis</i> | | 10 | 1992 |
| <i>Colobostruma elliotti</i> | | 5 | 1991 |
| <i>Discothyrea turtoni</i> | | 7 | 1992 |
| <i>Dolichoderus australis</i> | | 8 | 1994 |
| <i>Epopostruma alata</i> | | 15 | 1992 |
| <i>Iridomyrmex alpinus</i> | | 22 | 1993 |
| <i>Iridomyrmex bicknelli</i> | | 2 | 1993 |
| <i>Iridomyrmex coeruleus</i> | | 5 | 1983 |
| <i>Iridomyrmex dromus</i> | | <1 | 1972 |
| <i>Iridomyrmex mattirolói</i> | | 33 | 1993 |
| <i>Iridomyrmex meridianus</i> | | 22 | 1992 |
| <i>Iridomyrmex mjobergi</i> | | <1 | 1962 |
| <i>Iridomyrmex splendens</i> | | <1 | 1962 |
| <i>Linepithema humile</i> | | 7 | 1994 |
| <i>Monomorium leae</i> | | 6 | |
| <i>Myrmecia esuriens</i> | | 50 | 1980 |
| <i>Myrmecia forficata</i> | | 3 | 2005 |
| <i>Myrmecia fulvipes</i> | | 7 | 1967 |
| <i>Myrmecia pilosula</i> | | 5 | 1980 |
| <i>Myrmecia pilosula</i> gp | | 33 | 1992 |
| <i>Myrmecorhynchus carteri</i> | | 10 | 1993 |
| <i>Notoncus spinisquamis</i> | | 10 | 1995 |
| <i>Orectognathus clarki</i> | | 6 | 1977 |
| <i>Paratrechina anic_sos_sp.3</i> | | 1 | 1962 |
| <i>Pheidole tasmaniensis</i> | | 100 | |
| <i>Ponera leae</i> | | 4 | 1962 |
| <i>Rhytidoponera tasmaniensis</i> | | 15 | 1999 |
| <i>Rhytidoponera victoriae</i> | | 1 | 1993 |
| <i>Stigmacros barretti</i> | | 17 | 1994 |
| <i>Strumigenys perplexa</i> | | 10 | 1994 |
| <i>Technomyrmex jocosus</i> | | 10 | 1993 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------------------|-------------|---|--------------------|
| Halictidae | | | |
| (37 out of 438 species in Australia) | | | |
| Lasioglossum altichum | | 10 | 1992 |
| Lasioglossum asperithorax | | 5 | 1983 |
| Lasioglossum aus-sp1 | | 17 | 1988 |
| Lasioglossum bicingulatum | | 1 | 1988 |
| Lasioglossum blighi | | 24 | 1997 |
| Lasioglossum brazieri | | 4 | 1992 |
| Lasioglossum brunnesetum | | 6 | 1983 |
| Lasioglossum calophyllae | | 2 | 1983 |
| Lasioglossum clelandi | | 8 | 1983 |
| Lasioglossum cognatum | | 1 | 1975 |
| Lasioglossum cte-sp1 | | 8 | |
| Lasioglossum cte-sp13 | | 12 | 1997 |
| Lasioglossum disclusum | | 42 | 1997 |
| Lasioglossum dissimulator | | 8 | 1997 |
| Lasioglossum erythrurum | | <1 | 1988 |
| Lasioglossum excultum | | 32 | 1997 |
| Lasioglossum gilesi | | 2 | 1986 |
| Lasioglossum globosum | | 3 | |
| Lasioglossum hemichalceum | | <1 | 1972 |
| Lasioglossum hilactum | | 2 | 1988 |
| Lasioglossum imitans | | 6 | 1988 |
| Lasioglossum inteny | | 28 | 2001 |
| Lasioglossum lanarium | | 2 | 1992 |
| Lasioglossum lithuscum | | 27 | 1997 |
| Lasioglossum littleri | | 12 | 1983 |
| Lasioglossum macrops | | 50 | 1991 |
| Lasioglossum melbournense | | 18 | 1991 |
| Lasioglossum mirriji | | 80 | 1988 |
| Lasioglossum mundulum | | 6 | 1989 |
| Lasioglossum orbatum | | 2 | 1989 |
| Lasioglossum plorator | | 16 | 1997 |
| Lasioglossum repraesentans | | 13 | 1983 |
| Lasioglossum seductum | | 9 | 1975 |
| Lasioglossum sulthicum | | 8 | 1997 |
| Lasioglossum taluche | | 32 | 1997 |
| Lasioglossum turneri | | 12 | 1993 |
| Lasioglossum willsi | | 2 | 1997 |
| Megachilidae | | | |
| (1 out of 165 species in Australia) | | | |
| Megachile erythropyga | | 6 | |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|-----------------------|-------------|---|--------------------|
| Isopoda | | | | |
| Hypsimetopodidae | | | | |
| Phreatoicidean Isopods (1 out of 9 species in Australia) | | | | |
| Phreatoicoides longicollis | Phreatoicidean Isopod | | 100 | 2006 |
| Phreatoicidae | | | | |
| Phreatoicidean Isopods (7 out of 44 species in Australia) | | | | |
| Colubotelson thomsoni | Phreatoicidean Isopod | | 100 | 1956 |
| Mesacanthotelson setosus | Phreatoicidean Isopod | | 100 | 1981 |
| Mesacanthotelson tasmaniae | Phreatoicidean Isopod | | 100 | 1981 |
| Onchotelson brevicaudatus | Phreatoicidean Isopod | | 100 | 1983 |
| Onchotelson spatulatus | Phreatoicidean Isopod | | 100 | |
| Paraphreatoicus relictus | Phreatoicidean Isopod | | 100 | 1995 |
| Uramphisopus pearsoni | Phreatoicidean Isopod | | 100 | 1981 |
| Isoptera | | | | |
| Kalotermitidae | | | | |
| Termite (2 out of 33 species in Australia) | | | | |
| Bifiditermes improbus | Termite | | 3 | 1983 |
| Kalotermes convexus | | | 28 | 1983 |
| Termopsidae | | | | |
| Termite (2 out of 5 species in Australia) | | | | |
| Porotermes adamsoni | Termite | | 12 | 1983 |
| Stolotermes brunneicornis | Termite | | 60 | 1991 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-----------------------------|--|---|--------------------|
| Lepidoptera | | | | |
| Hesperiidae | | | | |
| Skippers, flats, awls & darts | | (9 out of 123 species in Australia) | | |
| Anisynta dominula | Two-brand Grass-skipper | | 24 | 1981 |
| Antipodia chaostola | Heath Sand-skipper | | 27 | 2008 |
| Hesperilla chrysotricha | Golden-haired Sedge-skipper | | 15 | 2009 |
| Hesperilla donnysa | Varied Sedge-skipper | | 15 | 1988 |
| Hesperilla idothea | Flame Sedge-skipper | | 7 | 1990 |
| Ocybadistes walkeri | Green Grass-dart | | 1 | 1960 |
| Pasma tasmanica | Two-spotted Grass-skipper | | 20 | 1960 |
| Taractrocera papyria | White-banded Grass-dart | | 5 | 2000 |
| Trapezites lutea | Yellow Ochre | | 14 | 1960 |
| Lycaenidae | | | | |
| Blues, Coppers, Hairstreaks and Metalmarks | | (8 out of 145 species in Australia) | | |
| Candalides acasta | Blotched Dusky-blue | | 2 | 1956 |
| Lampides boeticus | Long-tailed Pea-blue | | 1 | 1992 |
| Neolucia agricola | Fringed Heath-blue | | 3 | 1978 |
| Neolucia hobartensis | Montane Heath-blue | | 16 | 1990 |
| Paralucia aurifera | Bright Copper | | 4 | 1969 |
| Pseudalmenus chlorinda | Silky Hairstreak | | 25 | 2008 |
| Theclinessthes serpentata | Saltbush Blue | | 1 | 2009 |
| Zizina labradus | Common Grass-blue | | <1 | 1953 |
| Nymphalidae | | | | |
| Browns, Nymphs, Danaines | | (15 out of 83 species in Australia) | | |
| Argynnina hobartia | Tasmanian Brown | | 68 | 1988 |
| Danaus affinis | Swamp Tiger | | <1 | 1982 |
| Danaus chrysippus | Lesser Wanderer | | 1 | 1975 |
| Danaus plexippus | Monarch Butterfly | | 1 | 1995 |
| Geitoneura klugii | Marbled Xenica | | 6 | 1981 |
| Heteronympha cordace | Bright-eyed Brown | | 19 | 2006 |
| Heteronympha merope | Common Brown | | 4 | 2009 |
| Heteronympha penelope | Shouldered Brown | | 14 | 2009 |
| Junonia villida | Meadow Argus | | 2 | 2009 |
| Nesoxenica leprea | Tasmanian Xenica | | 64 | 1996 |
| Oreixenica lathoniella | Silver Xenica | | 16 | 1986 |
| Oreixenica orichora | Spotted Alpine Xenica | | 16 | 1983 |
| Oreixenica ptunarra | Ptunarra Xenica | | 65 | 2007 |
| Vanessa itea | Red Admiral | | <1 | 1957 |
| Vanessa kershawii | Australian Painted Lady | | <1 | 1975 |
| Papilionidae | | | | |
| Swallowtails | | (1 out of 19 species in Australia) | | |
| Graphium macleayanus | Macleay's Swordtail | | 7 | 2009 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------|---------------------------|---|---|--------------------|
| Pieridae | Whites and Yellows | (1 out of 35 species in Australia) | | |
| Belenois java | Caper White | | <1 | 1973 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|--|---|---|--------------------|
| Odonata | | | | |
| Aeshnidae | | | | |
| | Hawker, Duskhawker and Emperor Damselflies | (2 out of 13 species in Australia) | | |
| <i>Adversaeschna brevistyla</i> | Blue-spotted Hawker | | 3 | 1973 |
| <i>Hemianax papuensis</i> | Australian Emperor | | <1 | |
| Archipetalidae | | | | |
| | Tasmanian Redspot Dragonfly | (1 out of 1 species in Australia) | | |
| <i>Archipetalia auriculata</i> | Tasmanian Redspot | | 62 | 1997 |
| Coenagrionidae | | | | |
| | Bluetail and Thin Damselflies | (4 out of 32 species in Australia) | | |
| <i>Austroagrion cyane</i> | South-western Billabongfly | | 5 | 1995 |
| <i>Coenagrion lyelli</i> | Swamp Bluetail | | 22 | 1995 |
| <i>Ischnura aurora</i> | Aurora Bluetail | | 5 | 1997 |
| <i>Ischnura heterosticta</i> | Common Bluetail | | 4 | 1995 |
| Gomphidae | | | | |
| | Dragon, Hunter and Vicetail Dragonflies | (1 out of 35 species in Australia) | | |
| <i>Austrogomphus guerini</i> | Yellow-striped Hunter | | 7 | 1995 |
| Hemicorduliidae | | | | |
| | Emerald and Swamp Emerald Dragonflies | (3 out of 11 species in Australia) | | |
| <i>Hemicordulia australiae</i> | Australian Emerald | | 2 | 1995 |
| <i>Hemicordulia tau</i> | Tau Emerald | | 3 | 1993 |
| <i>Procordulia jacksoniensis</i> | Eastern Swamp Emerald | | 21 | 1995 |
| Lestidae | | | | |
| | Ringtail, Reedling and Dusky Spreadwing Damselflies | (5 out of 14 species in Australia) | | |
| <i>Austrolestes analis</i> | Slender Ringtail | | 4 | 1995 |
| <i>Austrolestes annulosus</i> | Blue Ringtail | | 11 | 1995 |
| <i>Austrolestes cingulatus</i> | Metallic Ringtail | | 13 | 1999 |
| <i>Austrolestes io</i> | Iota Ringtail | | 13 | 1995 |
| <i>Austrolestes psyche</i> | Cup Ringtail | | 14 | 1995 |
| Libellulidae | | | | |
| | Perchers, Archtails and other Dragonflies | (1 out of 52 species in Australia) | | |
| <i>Nannophya dalei</i> | Eastern Pygmyfly | | 5 | 1961 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|---|---|---|--------------------|
| Synthemistidae | Spottail and Tigertail Dragonflies | (3 out of 26 species in Australia) | | |
| <i>Archaeosynthemis orientalis</i> | Eastern Brown Tigertail | | 12 | |
| <i>Synthemopsis gomphomacromioides</i> | Tasmanian Spotwing | | 61 | 1994 |
| <i>Synthemis tasmanica</i> | Tasmanian Swamp Tigertail | | 54 | 1997 |
| Telephlebiidae | Darner Dragonflies | (4 out of 36 species in Australia) | | |
| <i>Austroaeschna hardyi</i> | Lesser Tasmanian Darner | | 49 | 1994 |
| <i>Austroaeschna parvistigma</i> | Swamp Darner | | 22 | 1995 |
| <i>Austroaeschna tasmanica</i> | Tasmanian Darner | | 48 | 1995 |
| <i>Austroaeschna unicornis</i> | Unicorn Darner | | 13 | 1995 |
| Philodinida | | | | |
| Habrotrochidae | Rotifers with Paired Trochi | (1 out of 23 species in Australia) | | |
| <i>Habrotrocha angusticollis</i> | Rotifer with Paired Trochi | | 17 | 1989 |
| Philodinidae | Rotifers with Paired Trochi | (4 out of 69 species in Australia) | | |
| <i>Dissotrocha macrostyla</i> | Rotifer with Paired Trochi | | 47 | 1989 |
| <i>Mniobia scarlatina</i> | Rotifer with Paired Trochi | | 33 | |
| <i>Rotaria rotatoria</i> | Rotifer with Paired Trochi | | 38 | |
| <i>Rotaria tardigrada</i> | Rotifer with Paired Trochi | | 20 | |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Ploimida | | | |
| Asplanchnidae | | | |
| Sac-like Wheel Animals (1 out of 13 species in Australia) | | | |
| Asplanchna sieboldi | | Siebolds Sac-like Wheel Animal | 12 |
| Brachionidae | | | |
| Sac-like Wheel Animal (15 out of 62 species in Australia) | | | |
| Anuraeopsis navicula | | Sac-like Wheel Animal | 7 |
| Brachionus angularis | | Angular Sac-like Wheel Animal | 15 |
| Brachionus calyciflorus | | Sac-like Wheel Animal | 2 |
| Brachionus lyratus | | Sac-like Wheel Animal | 14 |
| Brachionus quadridentatus | | Sac-like Wheel Animal | 7 |
| Brachionus tasmaniensis | | Tamanian Sac-like Wheel Animal | 100 |
| Brachionus urceolaris | | Sac-like Wheel Animal | 3 |
| Keratella australis | | Australian Sac-like Wheel Animal | 24 |
| Keratella cochlearis | | Ear-shaped Sac-like Wheel Animal | 22 1989 |
| Keratella hispida | | Sac-like Wheel Animal | 25 |
| Keratella javana | | Sac-like Wheel Animal | 38 1989 |
| Keratella procurva | | Sac-like Wheel Animal | 17 1989 |
| Keratella quadrata | | Sac-like Wheel Animal | 16 |
| Keratella slacki | | Slacks Sac-like Wheel Animal | 33 1989 |
| Keratella tropica | | Tropical Sac-like Wheel Animal | 2 |
| Dicranophoridae | | | |
| Tong-grabbing Wheel Animals (7 out of 41 species in Australia) | | | |
| Aspelta aper | | Tong-grabbing Wheel Animal | 50 |
| Dicranophorus forcipatus | | Tong-grabbing Wheel Animal | 33 |
| Dicranophorus hauerianus | | Tong-grabbing Wheel Animal | 12 |
| Dicranophorus hercules | | Tong-grabbing Wheel Animal | 100 |
| Dicranophorus robustus | | Tong-grabbing Wheel Animal | 53 1994 |
| Encentrum cruentum | | Tong-grabbing Wheel Animal | 100 |
| Encentrum diglandula | | Tong-grabbing Wheel Animal | 100 |
| Epiphanidae | | | |
| Shell-less Wheel Animals (4 out of 11 species in Australia) | | | |
| Epiphanes brachionus | | Shell-less Wheel Animal | 33 |
| Epiphanes clavulata | | Shell-less Wheel Animal | 18 |
| Epiphanes macroura | | Shell-less Wheel Animal | 12 |
| Proalides subtilis | | Shell-less Wheel Animal | 33 |
| Euchlanidae | | | |
| Wheel Animals (4 out of 19 species in Australia) | | | |
| Beauchampiella eudactylota | | Wheel Animal | 33 |
| Euchlanis dilatata | | Wheel Animal | 16 1989 |
| Euchlanis incisa | | Wheel Animal | 14 |
| Euchlanis meneta | | Wheel Animal | 50 1989 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|---------------------------------|-------------|---|--------------------|
| Gastropodidae | | | | |
| Shelled Wheel Animals (5 out of 8 species in Australia) | | | | |
| Ascomorpha ecaudis | Shelled Wheel Animal | | 31 | 1989 |
| Ascomorpha ovalis | Oval Shelled Wheel Animal | | 40 | 1989 |
| Ascomorpha saltans | Shelled Wheel Animal | | 20 | 1989 |
| Gastropus hyptopus | Shelled Wheel Animal | | 12 | |
| Gastropus stylifer | Shelled Wheel Animal | | 29 | |
| Ituridae | | | | |
| Shelled Wheel Animals (1 out of 4 species in Australia) | | | | |
| Itura aurita | Shelled Wheel Animal | | 14 | |
| Lecanidae | | | | |
| Shelled Wheel Animal (18 out of 78 species in Australia) | | | | |
| Lecane bulla | Shelled Wheel Animal | | 32 | 1989 |
| Lecane closterocerca | Shelled Wheel Animal | | 31 | 1989 |
| Lecane eylesi | Shelled Wheel Animal | | 100 | |
| Lecane flexilis | Shelled Wheel Animal | | 50 | 1989 |
| Lecane furcata | Shelled Wheel Animal | | 33 | |
| Lecane halicysta | Shelled Wheel Animal | | 100 | |
| Lecane hamata | Shelled Wheel Animal | | 43 | 1989 |
| Lecane hornemanni | Hornemanns Shelled Wheel Animal | | 22 | |
| Lecane inermis | Shelled Wheel Animal | | 67 | 1989 |
| Lecane inopinata | Shelled Wheel Animal | | 50 | |
| Lecane luna | Shelled Wheel Animal | | 10 | |
| Lecane lunaris | Shelled Wheel Animal | | 39 | 1989 |
| Lecane nana | Shelled Wheel Animal | | 75 | 1989 |
| Lecane pumila | Shelled Wheel Animal | | 100 | |
| Lecane quadridentata | Shelled Wheel Animal | | 50 | |
| Lecane stichaea | Shelled Wheel Animal | | 33 | |
| Lecane tenuiseta | Shelled Wheel Animal | | 50 | 1989 |
| Lecane thalera | Shelled Wheel Animal | | 100 | |
| Lepadellidae | | | | |
| Wheel Animals (13 out of 52 species in Australia) | | | | |
| Colurella adriatica | Adriatic Wheel Animal | | 11 | |
| Colurella colurus | Coloured Wheel Animal | | 100 | 1989 |
| Colurella obtusa | Obtuse Wheel Animal | | 30 | 1989 |
| Colurella tessellata | Tesselated Wheel Animal | | 100 | 1989 |
| Colurella uncinata | Wheel Animal | | 45 | 1989 |
| Lepadella acuminata | Wheel Animal | | 29 | |
| Lepadella amphitropis | Wheel Animal | | 100 | |
| Lepadella ovalis | Oval Wheel Animal | | 20 | |
| Lepadella patella | Wheel Animal | | 56 | 1989 |
| Lepadella rhomboides | Rhomboid Wheel Animal | | 9 | |
| Lepadella tanae | Wheel Animal | | 100 | |
| Lepadella triptera | Wheel Animal | | 50 | 1989 |
| Squatinella mutica | Wheel Animal | | 20 | |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------------|---------------------------------------|--|---|--------------------|
| Lindiidae | Wheel Animals | (1 out of 6 species in Australia) | | |
| <i>Lindia torulosa</i> | Wheel Animal | | 17 | |
| Microcodonidae | Wheel Animals | (1 out of 1 species in Australia) | | |
| <i>Microcodon clavus</i> | Wheel Animal | | 50 | |
| Mytilinidae | Wheel Animals | (2 out of 11 species in Australia) | | |
| <i>Lophocharis salpina</i> | Wheel Animal | | 29 | |
| <i>Mytilina ventralis</i> | Wheel Animal | | 11 | |
| Notommatidae | Pseudo-segmented Wheel Animals | (24 out of 93 species in Australia) | | |
| <i>Cephalodella catellina</i> | Pseudo-segmented Wheel Animal | | 33 | |
| <i>Cephalodella exigua</i> | Pseudo-segmented Wheel Animal | | 100 | |
| <i>Cephalodella f</i> | Pseudo-segmented Wheel Animal | | 100 | |
| <i>Cephalodella forficula</i> | Pseudo-segmented Wheel Animal | | 33 | |
| <i>Cephalodella gibba</i> | Pseudo-segmented Wheel Animal | | 29 | 1989 |
| <i>Cephalodella hoodi</i> | Hoods Pseudo-segmented Wheel Animal | | 100 | 1989 |
| <i>Cephalodella intuta</i> | Pseudo-segmented Wheel Animal | | 50 | |
| <i>Cephalodella mucronata</i> | Pseudo-segmented Wheel Animal | | 27 | 1989 |
| <i>Cephalodella nana</i> | Pseudo-segmented Wheel Animal | | 100 | |
| <i>Cephalodella rotunda</i> | Rotund Pseudo-segmented Wheel Animal | | 67 | 1989 |
| <i>Cephalodella ventripes</i> | Pseudo-segmented Wheel Animal | | 25 | |
| <i>Eosphora najas</i> | Pseudo-segmented Wheel Animal | | 50 | 1989 |
| <i>Eothinia elongata</i> | Pseudo-segmented Wheel Animal | | 50 | 1989 |
| <i>Monommata aeshyna</i> | Pseudo-segmented Wheel Animal | | 50 | 1989 |
| <i>Monommata dentata</i> | Pseudo-segmented Wheel Animal | | 33 | |
| <i>Monommata grandis</i> | Grand Pseudo-segmented Wheel Animal | | 33 | 1989 |
| <i>Monommata maculata</i> | Pseudo-segmented Wheel Animal | | 33 | 1989 |
| <i>Notommata aurita</i> | Pseudo-segmented Wheel Animal | | 100 | |
| <i>Notommata tripus</i> | Pseudo-segmented Wheel Animal | | 33 | |
| <i>Pleurata tyleri</i> | Tylers Pseudo-segmented Wheel Animal | | 50 | |
| <i>Pleurotrocha petromyzon</i> | Pseudo-segmented Wheel Animal | | 50 | |
| <i>Resticula cf. plicata</i> | Pseudo-segmented Wheel Animal | | 100 | |
| <i>Resticula melandocus</i> | Pseudo-segmented Wheel Animal | | 25 | |
| <i>Taphrocampa selenura</i> | Pseudo-segmented Wheel Animal | | 33 | |
| Proalidae | Worm-like Wheel Animals | (3 out of 15 species in Australia) | | |
| <i>Proales decipiens</i> | Worm-like Wheel Animal | | 50 | 1989 |
| <i>Proales fallaciosa</i> | Worm-like Wheel Animal | | 33 | |
| <i>Proalinopsis staurus</i> | Worm-like Wheel Animal | | 50 | |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------|---|-------------|---|--------------------|
| Scaridiidae | Wheel Animals | | (1 out of 4 species in Australia) | |
| Scaridium longicaudum | Wheel Animal | | 27 | 1989 |
| Synchaetidae | Sac-, Bell- or Cone-shaped Wheel Animals | | (6 out of 22 species in Australia) | |
| Polyarthra dolichoptera | Rotifer | | 28 | 1989 |
| Polyarthra remata | Rotifer | | 50 | |
| Polyarthra vulgaris | Rotifer | | 16 | |
| Synchaeta longipes | Sac-, Bell-, or Cone-shaped Wheel Animal | | 5 | |
| Synchaeta oblonga | Sac-, Bell- or Cone-shaped Wheel Animal | | 23 | |
| Synchaeta pectinata | Sac-, Bell- or Cone-shaped Wheel Animal | | 16 | 1989 |
| Trichocercidae | Comma-like Wheel Animals | | (17 out of 50 species in Australia) | |
| Trichocerca bidens | Comma-like Wheel Animal | | 20 | 1989 |
| Trichocerca brachyura | Comma-like Wheel Animal | | 36 | |
| Trichocerca capucina | Comma-like Wheel Animal | | 9 | |
| Trichocerca cavia | Comma-like Wheel Animal | | 100 | |
| Trichocerca collaris | Collared Comma-like Wheel Animal | | 67 | 1989 |
| Trichocerca gracilis | Gracile Comma-like Wheel Animal | | 100 | |
| Trichocerca insignis | Comma-like Wheel Animal | | 25 | |
| Trichocerca insulana | Comma-like Wheel Animal | | 100 | |
| Trichocerca intermedia | Intermediate Comma-like Wheel Animal | | 100 | |
| Trichocerca longiseta | Comma-like Wheel Animal | | 31 | |
| Trichocerca myersi | Myers Comma-like Wheel Animal | | 25 | |
| Trichocerca porcellus | Comma-like Wheel Animal | | 47 | 1989 |
| Trichocerca pusilla | Comma-like Wheel Animal | | 17 | 1989 |
| Trichocerca rattus | Rat-like Comma-like Wheel Animal | | 18 | |
| Trichocerca scipio | Scipios Comma-like Wheel Animal | | 50 | 1989 |
| Trichocerca similis | Comma-like Wheel Animal | | 32 | 1989 |
| Trichocerca tenuior | Comma-like Wheel Animal | | 17 | |
| Trichotriidae | Wheel Animals | | (3 out of 10 species in Australia) | |
| Macrochaetus collinsi | Collins Wheel Animal | | 67 | 1989 |
| Trichotria tetractis | Wheel Animal | | 29 | 1989 |
| Wolga spinifera | Wheel Animal | | 50 | |
| Polydesmida | | | | |
| Dalodesmidae | | | (2 out of 97 species in Australia) | |
| Dasystigma margaretae | | | 25 | 1995 |
| Lissodesmus perporosus | | | 33 | 2000 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------------------|-------------|---|--------------------|
| Pseudoscorpiones | | | |
| Chthoniidae | | | |
| (1 out of 32 species in Australia) | | | |
| Pseudotyrannochthonius tasmanicus | | 100 | 1996 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------------|--|---|--------------------|
| Trichoptera | | | |
| Atriplectididae | | | |
| Caddisflies | (1 out of 2 species in Australia) | | |
| Atriplectides dubius | | 85 | 1995 |
| Calocidae | | | |
| Caddisflies | (4 out of 18 species in Australia) | | |
| Caenota plicata | | 35 | 1989 |
| Caloca ascita | | 100 | 1977 |
| Caloca tertia | | 50 | 1990 |
| Tamasia variegata | | 44 | 1989 |
| Conoesucidae | | | |
| Caddisflies | (17 out of 22 species in Australia) | | |
| Conoesucus adiaxolus | | 38 | 1989 |
| Conoesucus brontensis | | 50 | 1988 |
| Conoesucus digitiferus | | 46 | 1989 |
| Conoesucus fromus | | 43 | 1989 |
| Conoesucus nepotulus | | 35 | 1988 |
| Conoesucus norelus | | 32 | 1989 |
| Conoesucus notialis | | 100 | 1988 |
| Costora delora | | 26 | 1989 |
| Costora ebenina | | 41 | 1988 |
| Costora krene | | 100 | 1988 |
| Costora luxata | | 33 | 1988 |
| Costora ramosa | | 55 | 1989 |
| Costora rotosca | | 56 | 1989 |
| Costora seposita | | 50 | 1988 |
| Hampa patona | | 27 | 1989 |
| Lingora aurata | | 44 | 1989 |
| Matasia satana | | 47 | 1988 |
| Ecnomidae | | | |
| Caseless Caddisflies | (5 out of 93 species in Australia) | | |
| Daternomina irrorata | | 80 | 1988 |
| Ecnomina batyle | | 100 | 1965 |
| Ecnomina legula | | 67 | 1988 |
| Ecnomus russellius | | 100 | 1988 |
| Ecnomus tillyardi | | 75 | 1988 |
| Glossosomatidae | | | |
| Stone-case Caddisflies | (2 out of 10 species in Australia) | | |
| Agapetus cralus | | 100 | 1965 |
| Agapetus tasmanicus | | 71 | 1988 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------|---------------------------------|---|--------------------|
| Helicophidae | Caddisflies | (4 out of 7 species in Australia) | |
| Alloecella grisea | | 32 | 1988 |
| Alloecella longispina | | 64 | 1989 |
| Alloecella pilosa | | 52 | 1988 |
| Helicopha astia | | 67 | 1987 |
| Helicopsychidae | Snail Case Caddisflies | (1 out of 14 species in Australia) | |
| Helicopsyche murrumba | | 100 | 1988 |
| Hydrobiosidae | Caddisflies | (21 out of 67 species in Australia) | |
| Apsilochorema gisbum | | 83 | 1988 |
| Apsilochorema obliquum | | 55 | 1988 |
| Austrochorema pegidion | | 25 | 1966 |
| Ethochorema kelion | | 33 | 1988 |
| Ethochorema nesydrion | | 56 | 1988 |
| Ipsobiosis spicula | | 100 | 1965 |
| Koetonga clivicola | | 75 | 1988 |
| Moruya charadra | | 83 | 1988 |
| Moruya opora | | 75 | 1988 |
| Poecilochorema complexum | | 100 | 1965 |
| Poecilochorema evansi | | 100 | 1953 |
| Ptychobiosis nigrita | | 57 | 1988 |
| Taschorema apobamum | | 71 | 1988 |
| Taschorema asmanum | | 67 | 1988 |
| Taschorema evansi | | 86 | 1988 |
| Taschorema ithyphallicum | | 50 | 1977 |
| Taschorema viridarium | | 75 | 1988 |
| Ulmerochorema breve | | 67 | 1966 |
| Ulmerochorema rubiconum | | 80 | 1988 |
| Ulmerochorema seonum | | 67 | 1988 |
| Ulmerochorema tasmanicum | | 75 | 1977 |
| Hydropsychidae | Net Spinning Caddisflies | (5 out of 35 species in Australia) | |
| Asmicridea edwardsii | whitemoth | 64 | 1989 |
| Cheumatopsyche modica | | 67 | 1993 |
| Diemeniluma tasmanica | | 50 | 1977 |
| Diplectrona castanea | Mount Field caddis fly | 100 | |
| Diplectrona lyella | King River caddis fly | 25 | 1999 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|--------------------------------|---|--------------------|
| Hydroptilidae | Micro Caddisflies | (7 out of 139 species in Australia) | |
| Hellyethira simplex | | 67 | 1988 |
| Maydenoptila cuneola | | 67 | 1988 |
| Orphninostrichia maculata | | 100 | 1999 |
| Orthotrichia adornata | | 100 | 1979 |
| Orthotrichia zonata | | 100 | 1966 |
| Oxyethira columba | | 100 | 1988 |
| Oxyethira mienica | Ouse River caddis fly | 100 | 1988 |
| Kokiriidae | Caddisflies | (3 out of 5 species in Australia) | |
| Taskiria austera | | 100 | 1978 |
| Taskiria mccubbini | | 100 | 1999 |
| Taskiropsyche lacustris | Lake Pedder caddis fly | 100 | 1999 |
| Leptoceridae | Long-horned Caddisflies | (26 out of 179 species in Australia) | |
| Condocerus paludosus | | 100 | 1995 |
| Lectrides varians | | 50 | 1965 |
| Leptorussa darlingtoni | | 100 | 1988 |
| Notalina bifaria | | 71 | 1988 |
| Notalina fulva | | 67 | 1988 |
| Notalina nigra | | 100 | 1977 |
| Notalina parkeri | | 83 | 1988 |
| Notoperata maculata | | 100 | 1988 |
| Notoperata sparsa | | 88 | 1988 |
| Oecetis arcada | | 75 | 1988 |
| Oecetis asmanista | | 100 | 1988 |
| Oecetis australis | | 100 | 1988 |
| Oecetis inscripta | | 100 | 1988 |
| Oecetis laustra | | 100 | 1988 |
| Oecetis minasata | | 67 | 1988 |
| Oecetis pechana | | 100 | 1988 |
| Oecetis scirpicula | | 100 | 1965 |
| Symphitoneuria opposita | | 50 | 1995 |
| Triaenodes intricata | | 100 | 1995 |
| Triplectides bilobus | | 58 | 1988 |
| Triplectides magnus | | 100 | 1965 |
| Triplectides proximus | | 100 | 1993 |
| Triplectides similis | | 100 | 1988 |
| Triplectides truncatus | | 50 | 1988 |
| Triplectidina nigricornis | | 67 | 1988 |
| Westriplectes pedderensis | | 100 | 1988 |
| Oeconesidae | Caddisflies | (1 out of 1 species in Australia) | |
| Tascuna ignota | | 100 | 1977 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|---|---|---|--------------------|
| Philopotamidae | Finger Net caddisflies, Silken Tube Spinners | (6 out of 48 species in Australia) | | |
| <i>Hydrobiosella armata</i> | Mount Wellington caddis fly | | 100 | 1967 |
| <i>Hydrobiosella cerula</i> | | | 100 | 1977 |
| <i>Hydrobiosella cognata</i> | | | 50 | 1977 |
| <i>Hydrobiosella orba</i> | | | 50 | 1965 |
| <i>Hydrobiosella tasmanica</i> | | | 100 | 1989 |
| <i>Hydrobiosella waddama</i> | | | 25 | 1966 |
| Philorheithridae | Caddisflies | (4 out of 14 species in Australia) | | |
| <i>Aphilorheithrus decoratus</i> | | | 83 | 1989 |
| <i>Aphilorheithrus luteolus</i> | | | 100 | 1966 |
| <i>Aphilorheithrus stepheni</i> | | | 67 | 1988 |
| <i>Tasmanthrus angustipennis</i> | | | 67 | 1989 |
| Plectrotarsidae | Caddisflies | (3 out of 5 species in Australia) | | |
| <i>Liapota lavara</i> | | | 67 | 1988 |
| <i>Nanoplectrus truchanasi</i> | | | 100 | 1965 |
| <i>Plectrotarsus tasmanicus</i> | | | 50 | 1977 |
| Polycentropodidae | Caseless Caddisflies | (5 out of 18 species in Australia) | | |
| <i>Paranyctiophylax repandus</i> | | | 75 | 1988 |
| <i>Plectrocnemia altera</i> | | | 60 | 1988 |
| <i>Plectrocnemia caudata</i> | | | 100 | 1988 |
| <i>Plectrocnemia manicata</i> | | | 75 | 1988 |
| <i>Tasmanoplegas spilota</i> | | | 50 | 1988 |
| Tasimiidae | Tasimiidae | (1 out of 7 species in Australia) | | |
| <i>Tasimia drepana</i> | Huon River caddis fly | | 100 | 1966 |
| Valvatida | | | | |
| Oreasteridae | | (1 out of 1 species in Australia) | | |
| <i>Nectria ocellata</i> | | | 43 | 1994 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|--------------------------|---|--------------------|
| Vascular flora | | | |
| Alismatales | | | |
| Alismataceae | | | |
| Water Plantains | | (2 out of 7 species in Australia) | |
| Alisma plantago-aquatica | Water Plantain | 3 | 2003 |
| Damasonium minus | | 2 | 2006 |
| Araceae | | | |
| Settlers Flax, Cunjevoi, Native Lilies | | (3 out of 46 species in Australia) | |
| Lemna disperma | | 5 | 2006 |
| Lemna trisulca | | 6 | 2004 |
| Wolffia australiana | | 5 | 2006 |
| Juncaginaceae | | | |
| Water Ribbons | | (7 out of 25 species in Australia) | |
| Triglochin alcockiae | | 13 | 2007 |
| Triglochin centrocarpa | | <1 | 1995 |
| Triglochin minutissima | | 1 | 1998 |
| Triglochin nana | | <1 | 1994 |
| Triglochin procera | | 12 | 2006 |
| Triglochin rheophila | | 7 | 2003 |
| Triglochin striata | Three-ribbed Arrow Grass | 5 | 2007 |
| Potamogetonaceae | | | |
| Sea Grasses and Horned Pondweeds, Pondweed | | (11 out of 16 species in Australia) | |
| Lepilaena bilocularis | | 3 | 1966 |
| Lepilaena cylindrocarpa | | 14 | 2006 |
| Lepilaena marina | | 5 | 1984 |
| Lepilaena patentifolia | | 20 | 1989 |
| Lepilaena preissii | | 9 | 1990 |
| Potamogeton australiensis | | 22 | 2004 |
| Potamogeton crispus | Curly-leaved Pondweed | 4 | 2007 |
| Potamogeton ochreatus | | 8 | 2007 |
| Potamogeton pectinatus | | 3 | 1992 |
| Potamogeton perfoliatus | | 5 | 2004 |
| Potamogeton tricarinatus | | 5 | 2007 |
| Ruppiaceae | | | |
| | | (4 out of 4 species in Australia) | |
| Ruppia maritima | | 2 | 1988 |
| Ruppia megacarpa | | 9 | 2006 |
| Ruppia polycarpa | | 13 | 2002 |
| Ruppia tuberosa | | 3 | 2006 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|---------------------|-------------|---|--------------------|
| Asparagales | | | | |
| Asparagaceae (5 out of 177 species in Australia) | | | | |
| Arthropodium milleflorum | Vanilla Lily | | 10 | 2005 |
| Arthropodium minus | Small Vanilla Lily | | 2 | 2005 |
| Arthropodium strictum | Chocolate Lily | | 1 | 2007 |
| Laxmannia orientalis | Dwarf Paper-Lily | | 6 | 2000 |
| Sowerbaea juncea | Vanilla Lily | | <1 | 1979 |
| Asphodelaceae Native Leek (3 out of 7 species in Australia) | | | | |
| Bulbine bulbosa | Bulbine Lily, Pike | | 2 | 2008 |
| Bulbine glauca | Rock Lily | | 33 | 2007 |
| Bulbine semibarbata | Leek Lily | | 2 | 2008 |
| Asteliaceae Palm-Lilies and Allies (6 out of 9 species in Australia) | | | | |
| Astelia alpina | Pineapple Grass | | 29 | 2007 |
| Milligania densiflora | Milligania Lily | | 62 | 2006 |
| Milligania johnstonii | Milligania Lily | | 25 | 2005 |
| Milligania lindoniana | Milligania Lily | | 47 | 2006 |
| Milligania longifolia | Milligania Lily | | 42 | 1989 |
| Milligania stylosa | Milligania Lily | | 100 | 2006 |
| Blandfordiaceae Christmas Bells (1 out of 4 species in Australia) | | | | |
| Blandfordia punicea | | | 38 | 2004 |
| Hemerocallidaceae (12 out of 79 species in Australia) | | | | |
| Caesia alpina | Alpine Grass-Lily | | 17 | 2006 |
| Caesia calliantha | Blue Grass-Lily | | 1 | 1990 |
| Caesia parviflora | Pale Grass-Lily | | 2 | 2003 |
| Dianella amoena | Matted Flax-lily | Endangered | 22 | 2009 |
| Dianella brevicaulis | Flax-lily | | 5 | 2007 |
| Dianella longifolia | Flax-lily | | <1 | 1994 |
| Dianella revoluta | Spreading Flax-lily | | 3 | 2009 |
| Dianella tasmanica | Tasman Flax-lily | | 14 | 2009 |
| Herpolirion novae-zelandiae | Sky-lily | | 28 | 2007 |
| Thelionema caespitosum | Native Lily | | 11 | 2007 |
| Thelionema umbellatum | Blue Lily | | 6 | 1996 |
| Tricoryne elatior | Native Lily | | <1 | 1984 |
| Hypoxidaceae Native Lilies (3 out of 15 species in Australia) | | | | |
| Hypoxis glabella | Tiny Stars | | 3 | 2008 |
| Hypoxis hygrometrica | Golden Weathergrass | | 9 | 2007 |
| Hypoxis vaginata | Yellow Star | | 4 | 2008 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------------|--------------------------|---|---|--------------------|
| Iridaceae | Irises and Allies | (7 out of 28 species in Australia) | | |
| <i>Diplarrena latifolia</i> | Native Iris | | 43 | 2005 |
| <i>Diplarrena moraea</i> | Native Iris | | 36 | 2008 |
| <i>Isophysis tasmanica</i> | Native Iris | | 65 | 2000 |
| <i>Libertia pulchella</i> | Pretty Grass-Flag | | 20 | 1998 |
| <i>Patersonia fragilis</i> | Short Purple Flag | | 12 | 2099 |
| <i>Patersonia occidentalis</i> | Purple Flag | | 1 | 2006 |
| <i>Patersonia sericea</i> | Native Flag | | <1 | |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|-------------|---|--------------------|
| Orchidaceae | | | |
| Orchids | | (207 out of 1248 species in Australia) | |
| Acianthus exsertus | | 2 | 1997 |
| Acianthus pusillus | | 12 | 2009 |
| Anzybas unguiculatus | | 13 | 2007 |
| Arachnorchis anthracina | Endangered | 14 | 1992 |
| Arachnorchis caudata | Vulnerable | 37 | 2009 |
| Arachnorchis clavigera | | 12 | 2001 |
| Arachnorchis dilatata | | 5 | 2001 |
| Arachnorchis echidnachila | | 81 | 2009 |
| Arachnorchis helvina | | 68 | 2002 |
| Arachnorchis lindleyana | Endangered | 15 | 1992 |
| Arachnorchis pallida | Endangered | 17 | 1991 |
| Arachnorchis patersonii | | 10 | 1996 |
| Arachnorchis saggicola | Endangered | 75 | 2009 |
| Arachnorchis venusta | | 2 | 1994 |
| Bunochilus longifolius | | 5 | 1996 |
| Bunochilus melagrammus | | 21 | 2008 |
| Bunochilus stenochilus | | 84 | 2009 |
| Bunochilus williamsonii | | 82 | 2008 |
| Burnettia cuneata | | 21 | 2007 |
| Caladenia latifolia | | 1 | 1990 |
| Caladenia similis | | 50 | 1991 |
| Caleana major | | 9 | 2008 |
| Calochilus campestris | | <1 | 1964 |
| Calochilus herbaceus | | 34 | 2008 |
| Calochilus imberbis | | 7 | 1992 |
| Calochilus paludosus | | 6 | 2007 |
| Calochilus robertsonii | | 5 | 2008 |
| Chiloglottis reflexa | | 14 | 2008 |
| Chiloglottis trilabra | | 2 | 2003 |
| Corunastylis archeri | | 14 | 2009 |
| Corunastylis brachystachya | Endangered | 18 | 1989 |
| Corunastylis despectans | | 11 | 2006 |
| Corunastylis firthii | Endangered | 100 | 1999 |
| Corunastylis morrisii | | 5 | 2008 |
| Corunastylis nigricans | | <1 | 1984 |
| Corunastylis nuda | | 10 | 2008 |
| Corunastylis nudiscapa | | 3 | 2008 |
| Corunastylis pumila | | 11 | 2006 |
| Corunastylis rufa | | 2 | 1996 |
| Corunastylis tasmanica | | 58 | 2002 |
| Corybas aconitiflorus | | 9 | 2007 |
| Corysanthes diemenica | | 28 | 2008 |
| Corysanthes dilatata | | 3 | 1994 |
| Corysanthes fimbriata | | 5 | 2004 |
| Corysanthes incurva | | 11 | 2007 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------------------|---|-------------|---|--------------------|
| <i>Crangonorchis pedoglossa</i> | Prawn Greenhood | | 32 | 2007 |
| <i>Cryptostylis subulata</i> | Large Tongue Orchid | | 6 | 2001 |
| <i>Cyrtostylis reniformis</i> | Common Gnat Orchid | | 9 | 2009 |
| <i>Cyrtostylis robusta</i> | Large Gnat Orchid | | 2 | 2009 |
| <i>Diplodium alatum</i> | Striped Greenhood | | 22 | 2009 |
| <i>Diplodium atrans</i> | | | 8 | 2006 |
| <i>Diplodium coccinum</i> | | | 1 | 1992 |
| <i>Diplodium decurvum</i> | Summer Greenhood | | 21 | 2006 |
| <i>Diplodium grandiflorum</i> | Cobra Greenhood | | 3 | 2008 |
| <i>Diplodium obtusum</i> | Blunt-tongue Greenhood | | <1 | 1991 |
| <i>Dipodium roseum</i> | Rosy Hyacinth Orchid | | 8 | 2007 |
| <i>Diuris chryseopsis</i> | Common Golden Moths | | 15 | 2007 |
| <i>Diuris corymbosa</i> | Wallflower Donkey Orchid | | <1 | 1985 |
| <i>Diuris lanceolata</i> | Snake Orchid, Large Golden Moths | Endangered | 2 | 1994 |
| <i>Diuris maculata</i> | Leopard Orchid, Spotted Doubletail | | 3 | 1996 |
| <i>Diuris monticola</i> | Highland Golden Moths | | 26 | 2004 |
| <i>Diuris orientis</i> | Eastern Wallflower Diuris | | 2 | 1990 |
| <i>Diuris palustris</i> | Swamp Diuris | | 5 | 2002 |
| <i>Diuris pardina</i> | Leopard Orchid, Spotted Doubletail | | 8 | 2008 |
| <i>Diuris pedunculata</i> | Small Snake Orchid | Endangered | 4 | 1993 |
| <i>Diuris sulphurea</i> | Tiger Orchid | | 8 | 2008 |
| <i>Dockrillia striolata</i> | Streaked Rock Orchid | | 6 | 2008 |
| <i>Eriochilus aff. cucullatus</i> 3 | | | 33 | 1992 |
| <i>Eriochilus autumnalis</i> | Parson's Bands | | 1 | 1984 |
| <i>Eriochilus cucullatus</i> | Parson's Bands | | 11 | 2009 |
| <i>Gastrodia procera</i> | Tall Potato Orchid | | 18 | 2008 |
| <i>Gastrodia sesamoides</i> | Cinnamon Bells, Common Potato Orchid | | 11 | 2008 |
| <i>Glossodia major</i> | Wax-lip Orchid | | 7 | 2009 |
| <i>Glossodia minor</i> | Small Wax Lip | | 4 | 1998 |
| <i>Hymenochilus cycnocephalus</i> | Swan Greenhood | | 1 | 1993 |
| <i>Hymenochilus muticus</i> | Midget Greenhood | | <1 | 1952 |
| <i>Hymenochilus wapstrarum</i> | Fleshy Greenhood | Endangered | 71 | 2007 |
| <i>Jonesiopsis filamentosa</i> | | | 12 | 2007 |
| <i>Linguella nana</i> | Dwarf Greenhood | | 3 | 2009 |
| <i>Lyperanthus suaveolens</i> | Brown Beaks | | 9 | 2006 |
| <i>Microtidium atratum</i> | Yellow Onion Orchid | | 1 | 2005 |
| <i>Microtis arenaria</i> | | | 6 | 2007 |
| <i>Microtis parviflora</i> | Slender Onion Orchid | | 3 | 2008 |
| <i>Microtis rara</i> | Scented Onion Orchid, Sweet Onion Orchi | | 5 | 2001 |
| <i>Microtis unifolia</i> | Common Onion Orchid | | 6 | 2008 |
| <i>Nemacianthus caudatus</i> | Dead Horse, Mayfly Orchid | | 7 | 1998 |
| <i>Nematoceras dienema</i> | Windswept Helmet Orchid | | 50 | 1998 |
| <i>Oligochaetochilus commutatus</i> | Midland Greenhood | Endangered | 20 | 2001 |
| <i>Orthoceras strictum</i> | Horned Orchid | | 3 | 2006 |
| <i>Paracaleana minor</i> | Small Duck Orchid | | 10 | 2006 |
| <i>Petalochilus alatus</i> | | | 23 | 2001 |
| <i>Petalochilus carneus</i> | Pink Fingers | | 5 | 2008 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|---|-----------------------|---|--------------------|
| Petalochilus catenatus | White Caladenia | | 4 | 1995 |
| Petalochilus fuscatus | | | 8 | 2009 |
| Petalochilus mentiens | Lesser Caladenia | | 9 | 2000 |
| Petalochilus prolatus | Long-leaved caladenia | | 2 | 2001 |
| Petalochilus pusillus | Tiny Caladenia | | 3 | 2003 |
| Petalochilus quadrifarius | | | 3 | 1992 |
| Petalochilus sylvicola | Forest Fingers | Endangered | 100 | 2009 |
| Petalochilus tonellii | Robust Fingers | Endangered | 12 | 1974 |
| Petalochilus vulgaris | Summer Fingers | | 26 | 2000 |
| Pheladenia deformis | Blue Fairies | | 4 | 2007 |
| Plumatchilos plumosum | Bearded Greenhood | | 5 | 1999 |
| Plumatchilos tasmanicum | | | 20 | 2005 |
| Prasophyllum alpestre | Highland Leek Orchid | | 13 | 2001 |
| Prasophyllum alpinum | Alpine Leek Orchid | | 31 | 2004 |
| Prasophyllum amoenum | Dainty Leek Orchid | Endangered | 100 | 2009 |
| Prasophyllum apoxychilum | Tapered Leek Orchid | Endangered | 63 | 2002 |
| Prasophyllum australe | Austral Leek Orchid | | 14 | 2006 |
| Prasophyllum brevilabre | Short-lip Leek Orchid | | 11 | 2006 |
| Prasophyllum castaneum | Chestnut Leek Orchid | Critically endangered | 88 | 2003 |
| Prasophyllum concinnum | Trim Leek Orchid | | 61 | 2008 |
| Prasophyllum elatum | Tall Leek Orchid | | 7 | 2002 |
| Prasophyllum flavum | Yellow Leek Orchid | | 19 | 1998 |
| Prasophyllum fuscum | Tailed Leek Orchid | Vulnerable | 2 | 1985 |
| Prasophyllum incurvum | Horseshoe Leek Orchid | | 78 | 2009 |
| Prasophyllum lindleyanum | Green Leek Orchid | | 26 | 2006 |
| Prasophyllum milfordense | Milford Leek Orchid | Critically endangered | 100 | 2007 |
| Prasophyllum montanum | Leek Orchid | | 7 | 1993 |
| Prasophyllum odoratum | Leek Orchid | | 3 | 1999 |
| Prasophyllum patens | Broad-lip Leek Orchid | | 3 | 1969 |
| Prasophyllum perangustum | Knocklofty Leek Orchid | Critically endangered | 100 | 2001 |
| Prasophyllum pulchellum | Pretty Leek Orchid | Critically endangered | 56 | 2003 |
| Prasophyllum rogersii | Marsh Leek Orchid | | 6 | 1988 |
| Prasophyllum rostratum | Slaty Leek Orchid | | 8 | 1990 |
| Prasophyllum secutum | Northern Leek Orchid | Endangered | 9 | 1990 |
| Prasophyllum similis | Leek Orchid | | 100 | 1989 |
| Prasophyllum sphacelatum | Leek Orchid | | 14 | 2005 |
| Prasophyllum suttonii | Mauve Leek Orchid, Sutton's Leek Orchid | | 14 | 1992 |
| Prasophyllum tadgellianum | Tadgell's Leek Orchid | | 5 | 2004 |
| Prasophyllum truncatum | Truncate Leek Orchid | | 36 | 2005 |
| Prasophyllum tunbridgense | Tunbridge Leek Orchid | Endangered | 17 | 2009 |
| Pterostylis alpina | Mountain Greenhood | | 2 | 1992 |
| Pterostylis curta | Blunt Greenhood | | 4 | 2004 |
| Pterostylis dubia | Blue-tongued Greenhood | | 52 | 2004 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------------------|---|-------------|---|--------------------|
| <i>Pterostylis erecta</i> | Upright Maroonhood | | 2 | 1992 |
| <i>Pterostylis falcata</i> | Sickle Greenhood | | 2 | 1991 |
| <i>Pterostylis foliata</i> | Slender Greenhood | | 18 | 2005 |
| <i>Pterostylis furcata</i> | Sickle Greenhood | | 5 | 1996 |
| <i>Pterostylis nutans</i> | Nodding Greenhood | | 8 | 2008 |
| <i>Pterostylis pedunculata</i> | Maroonhood | | 10 | 2009 |
| <i>Pterostylis pratensis</i> | Liawenee Greenhood | Vulnerable | 92 | 2009 |
| <i>Pterostylis scabrada</i> | | | 30 | 2005 |
| <i>Pterostylis ziegeleri</i> | Cape Portland Greenhood | Vulnerable | 25 | 2009 |
| <i>Pyrorchis nigricans</i> | Red Beaks, Undertaker Orchid | | <1 | 2001 |
| <i>Sarcochilus australis</i> | Butterfly Orchid, Gunn's Tree Orchid | | 6 | 1999 |
| <i>Simpliglottis cornuta</i> | Green Bird Orchid | | 22 | 2005 |
| <i>Simpliglottis grammata</i> | Small Bird-orchid | | 51 | 2005 |
| <i>Simpliglottis gunnii</i> | Common Bird Orchid | | 9 | 2008 |
| <i>Simpliglottis triceratops</i> | Three-horned Bird-orchid | | 56 | 2008 |
| <i>Simpliglottis valida</i> | Large Bird-orchid | | 2 | 1993 |
| <i>Speculantha aphylla</i> | | | 27 | 2002 |
| <i>Speculantha atriola</i> | | Endangered | 21 | 2009 |
| <i>Speculantha parviflora</i> | | | 8 | 2007 |
| <i>Speculantha uliginosa</i> | | | 11 | 1996 |
| <i>Spiranthes australis</i> | Pink Spiral Orchid | | 4 | 2009 |
| <i>Stegostyla alpina</i> | | | 25 | 2006 |
| <i>Stegostyla angustata</i> | | | 15 | 1996 |
| <i>Stegostyla atrata</i> | Dark Caladenia | | 75 | 2008 |
| <i>Stegostyla congesta</i> | Black-tongue Caladenia | | <1 | 1980 |
| <i>Stegostyla cracens</i> | | | 74 | 2008 |
| <i>Stegostyla cucullata</i> | Hooded Caladenia | | 4 | 1992 |
| <i>Stegostyla gracilis</i> | Musky Caladenia | | 11 | 2008 |
| <i>Stegostyla iridescens</i> | Bronze Caladenia | | 4 | 1992 |
| <i>Stegostyla lyallii</i> | Mountain Caladenia | | 38 | 1999 |
| <i>Stegostyla praecox</i> | | | 3 | 1990 |
| <i>Stegostyla transitoria</i> | | | 6 | 2004 |
| <i>Taurantha concinna</i> | | | 11 | 2009 |
| <i>Thelymitra aff. pauciflora</i> 1 | | | 15 | 2004 |
| <i>Thelymitra aggericola</i> | Bleak Sun Orchid | | 38 | 2008 |
| <i>Thelymitra antennifera</i> | Vanilla Orchid, Lemon-scented Sun Orchi | | <1 | 1999 |
| <i>Thelymitra arenaria</i> | Lazy Sun Orchid | | 15 | 2004 |
| <i>Thelymitra aristata</i> | Great Sun Orchid | | 16 | 2008 |
| <i>Thelymitra atronitida</i> | Sun Orchid | | 23 | 2008 |
| <i>Thelymitra brevifolia</i> | Sun Orchid | | 16 | 2008 |
| <i>Thelymitra carnea</i> | Tiny Sun Orchid, Pink Sun Orchid | | 5 | 1999 |
| <i>Thelymitra circumsepta</i> | Sun Orchid | | 24 | 2004 |
| <i>Thelymitra cyanea</i> | Veined Sun Orchid | | 22 | 2009 |
| <i>Thelymitra decora</i> | Sun Orchid | | 20 | 1986 |
| <i>Thelymitra erosa</i> | Striped Sun Orchid | | 29 | 2008 |
| <i>Thelymitra exigua</i> | Sun Orchid | | 10 | 2006 |
| <i>Thelymitra flexuosa</i> | Twisted Sun Orchid | | 3 | 1994 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|--|-----------------------|---|--------------------|
| <i>Thelymitra holmesii</i> | Holme's Sun Orchid | | 10 | 2006 |
| <i>Thelymitra imbricata</i> | Broad Sun Orchid | | 38 | 2004 |
| <i>Thelymitra inflata</i> | Sun Orchid | | 7 | 1992 |
| <i>Thelymitra ixioides</i> | Dotted Sun Orchid | | 5 | 2008 |
| <i>Thelymitra jonesii</i> | Sun Orchid | Critically endangered | 60 | 2006 |
| <i>Thelymitra juncifolia</i> | Large Spotted Sun Orchid | | 11 | 2005 |
| <i>Thelymitra longiloba</i> | Lobed Sun Orchid | | 6 | 1970 |
| <i>Thelymitra lucida</i> | Sun Orchid | | 7 | 2001 |
| <i>Thelymitra luteocilium</i> | Fringed Sun Orchid | | <1 | 1971 |
| <i>Thelymitra malvina</i> | Mauve-tufted Sun Orchid | | 7 | 2008 |
| <i>Thelymitra media</i> | Tall Sun Orchid | | 7 | 1996 |
| <i>Thelymitra mucida</i> | Hoary Sun Orchid, Plum Orchid | | 3 | 1980 |
| <i>Thelymitra nuda</i> | Plain Sun Orchid, Scented Sun Orchid | | 3 | 2003 |
| <i>Thelymitra pauciflora</i> | Slender Sun Orchid | | 7 | 2008 |
| <i>Thelymitra peniculata</i> | Sun Orchid | | 5 | 2008 |
| <i>Thelymitra polychroma</i> | Rainbow Sun Orchid | | 8 | 2008 |
| <i>Thelymitra pulchella</i> | Beautiful Sun Orchid, Rare Veined Sun Or | | 25 | 1996 |
| <i>Thelymitra rubra</i> | Pink Sun Orchid, Salmon Sun Orchid | | 12 | 2005 |
| <i>Thelymitra silena</i> | Madonna Sun Orchid | | 40 | 2002 |
| <i>Thelymitra spadicea</i> | Browntop Sun Orchid | | 17 | 2001 |
| <i>Thelymitra sparsa</i> | Wispy Sun Orchid | | 75 | 2003 |
| <i>Thelymitra venosa</i> | Large Veined Sun Orchid | | 9 | 1983 |
| <i>Thelymitra viridis</i> | Sun Orchid | | 29 | 2008 |
| <i>Thynniorchis nothofagicola</i> | | | 100 | 1995 |
| <i>Townsonia viridis</i> | Beech Orchid | | 52 | 2006 |
| Xanthorrhoeaceae | Grass Trees, Lomandras and Allies | | (2 out of 30 species in Australia) | |
| <i>Xanthorrhoea arenaria</i> | | Vulnerable | 12 | 2008 |
| <i>Xanthorrhoea australis</i> | | | 11 | 2008 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|-----------------------|---|--------------------|
| Asterales | | | |
| Asteraceae | | | |
| Daisies and Allies (197 out of 1019 species in Australia) | | | |
| Abrotanella forsteroides | | 62 | 2007 |
| Abrotanella scapigera | | 56 | 2000 |
| Actites megalocarpus | | 8 | 2006 |
| Allittia cardiocarpa | | 8 | 2005 |
| Ammobium calyceroides | | 2 | 1992 |
| Angianthus preissianus | | 2 | 2006 |
| Apalochlamys spectabilis | | 6 | 2007 |
| Argentipallium dealbatum | | 38 | 2009 |
| Argentipallium obtusifolium | | <1 | |
| Argentipallium spiceri | Critically endangered | 25 | 2000 |
| Argyrotegium fordianum | | 10 | 1980 |
| Argyrotegium mackayi | | 24 | 2002 |
| Argyrotegium nitidulum | Vulnerable | 12 | 1994 |
| Argyrotegium poliochlorum | | 8 | 1984 |
| Bedfordia linearis | | 51 | 2008 |
| Bedfordia salicina | | 52 | 2009 |
| Brachyglottis brunonis | | 100 | 2008 |
| Brachyscome aculeata | | 23 | 2008 |
| Brachyscome angustifolia | | 2 | 1995 |
| Brachyscome ciliaris | | <1 | 1990 |
| Brachyscome decipiens | | 11 | 2006 |
| Brachyscome diversifolia | | 4 | 2009 |
| Brachyscome graminea | | 12 | 2006 |
| Brachyscome nivalis | | 2 | 1984 |
| Brachyscome perpusilla | | <1 | |
| Brachyscome radicans | | 21 | 2005 |
| Brachyscome radicata | | 60 | 2009 |
| Brachyscome rigidula | | 7 | 2007 |
| Brachyscome sieberi | | 100 | 1997 |
| Brachyscome spathulata | | 14 | 2009 |
| Brachyscome tenuiscapa | | 14 | 1998 |
| Calocephalus brownii | | 1 | 2002 |
| Calocephalus citreus | | 1 | 2009 |
| Calocephalus lacteus | | 8 | 2009 |
| Cassinia aculeata | | 12 | 2009 |
| Cassinia trinerva | | 2 | 1996 |
| Celmisia asteliifolia | | 50 | 2007 |
| Celmisia longifolia | | 12 | 1993 |
| Celmisia saxifraga | | 59 | 2005 |
| Centipeda elatinoides | | 7 | 2007 |
| Centipeda minima | | 1 | 1995 |
| Chamaemelum nobile | | 9 | 1986 |
| Chrysocephalum apiculatum | | 2 | 2009 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------------------|----------------------|-------------|---|--------------------|
| <i>Chrysocephalum semipapposum</i> | Native Daisy | | 2 | 2007 |
| <i>Cotula alpina</i> | Alpine Cotula | | 30 | 2004 |
| <i>Cotula australis</i> | Common Brass-buttons | | 5 | 2007 |
| <i>Cotula longipes</i> | Creeping Cotula | | 50 | 1990 |
| <i>Cotula reptans</i> | Cotula | | 21 | 2006 |
| <i>Cotula vulgaris</i> | Cotula | | 3 | 2004 |
| <i>Craspedia coolaminica</i> | Native Daisy | | 8 | 2004 |
| <i>Craspedia glabrata</i> | Native Daisy | | 91 | 2002 |
| <i>Craspedia glauca</i> | Native Daisy | | 26 | 2007 |
| <i>Craspedia paludicola</i> | Native Daisy | | 2 | 1998 |
| <i>Craspedia richea</i> | Native Daisy | | 54 | 2005 |
| <i>Craspedia uniflora</i> | Native Daisy | | 3 | |
| <i>Craspedia variabilis</i> | Native Daisy | | <1 | 1984 |
| <i>Cymbonotus lawsonianus</i> | Austral Bears-ears | | 3 | 1989 |
| <i>Cymbonotus preissianus</i> | Bears-ears | | 2 | 2006 |
| <i>Euchiton argentifolius</i> | Native Daisy | | 28 | 1999 |
| <i>Euchiton gymnocephalus</i> | Native Daisy | | 18 | 2009 |
| <i>Euchiton involucratus</i> | Native Daisy | | 2 | 2006 |
| <i>Euchiton litticola</i> | Native Daisy | | 46 | 2007 |
| <i>Euchiton sphaericus</i> | Native Daisy | | <1 | 2004 |
| <i>Euchiton traversii</i> | Native Daisy | | 40 | 2004 |
| <i>Euchiton umbricolus</i> | Native Daisy | | 25 | 1997 |
| <i>Ewartia catipes</i> | Native Daisy | | 52 | 2007 |
| <i>Ewartia meredithae</i> | Native Daisy | | 58 | 2007 |
| <i>Ewartia planchonii</i> | Native Daisy | | 48 | 2005 |
| <i>Gnaphalium indutum</i> | Native Daisy | | <1 | 2001 |
| <i>Gnaphalium limosum</i> | Native Daisy | | 8 | 1981 |
| <i>Helichrysum ericifolius</i> | Native Daisy | | 88 | 1984 |
| <i>Helichrysum gunnii</i> | Native Daisy | | 33 | 2003 |
| <i>Helichrysum hookeri</i> | Native Daisy | | 50 | 1990 |
| <i>Helichrysum ledifolium</i> | Native Daisy | | 50 | 2007 |
| <i>Helichrysum leucopsidium</i> | Native Daisy | | <1 | 1989 |
| <i>Helichrysum milliganii</i> | Native Daisy | | 54 | 1998 |
| <i>Helichrysum paraliu</i> | Native Daisy | | 20 | 1990 |
| <i>Helichrysum pumilum</i> | Native Daisy | | 44 | 2008 |
| <i>Helichrysum purpurascens</i> | Native Daisy | | 67 | 1988 |
| <i>Helichrysum reticulatum</i> | Native Daisy | | 100 | 1991 |
| <i>Helichrysum rosmarinifolium</i> | Native Daisy | | 48 | 1992 |
| <i>Helichrysum rutidolepis</i> | Native Daisy | | <1 | 2000 |
| <i>Helichrysum scorpioides</i> | Native Daisy | | 10 | 2008 |
| <i>Helichrysum scutellifolium</i> | Native Daisy | | 87 | 1990 |
| <i>Hyalosperma demissum</i> | Everlasting | | 1 | 2007 |
| <i>Isoetopsis graminifolia</i> | Grass-cushions | | <1 | 2009 |
| <i>Ixodia angusta</i> | Hills Daisy | | 100 | 1980 |
| <i>Lagenophora gracilis</i> | Native Daisy | | 2 | 2004 |
| <i>Lagenophora huegelii</i> | Native Daisy | | 6 | 2007 |
| <i>Lagenophora stipitata</i> | Native Daisy | | 11 | 2009 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------|-----------------------------|-------------|---|--------------------|
| Leptinella filicula | Native Daisy | | 6 | 1998 |
| Leptorhynchos elongatus | Native Daisy | | 4 | 2006 |
| Leptorhynchos nitidulus | Native Daisy | | 18 | 2006 |
| Leptorhynchos squamatus | Native Daisy | | 12 | 2009 |
| Leucochrysum albicans | Everlasting | | 2 | 2008 |
| Microseris lanceolata | Native Daisy | | 7 | 2006 |
| Millotia tenuifolia | Native Daisy | | 3 | 2007 |
| Odidia achlaena | Native Daisy | | 100 | 2009 |
| Odidia angusta | Native Daisy | | 87 | 2006 |
| Olearia algida | Alpine Daisy Bush | | 39 | 2005 |
| Olearia archeri | Native Daisy | | 86 | 2008 |
| Olearia argophylla | Musk Daisy Bush | | 16 | 2009 |
| Olearia axillaris | Daisy Bush | | <1 | 2009 |
| Olearia ciliata | Daisy Bush | | 5 | 2008 |
| Olearia ericoides | Daisy Bush | | 65 | 2008 |
| Olearia erubescens | Daisy Bush | | 12 | 2008 |
| Olearia floribunda | Daisy Bush | | 11 | 2006 |
| Olearia glandulosa | Daisy Bush | | 17 | 2007 |
| Olearia hookeri | Hookers Daisy Bush | | 58 | 2009 |
| Olearia lanceolata | Daisy Bush | | 71 | 1987 |
| Olearia lanuginosa | Daisy Bush | | <1 | |
| Olearia ledifolia | Daisy Bush | | 52 | 2005 |
| Olearia lepidophylla | Daisy Bush | | 5 | 2009 |
| Olearia lirata | Snowy Daisy Bush | | 11 | 2009 |
| Olearia myrsinoides | Daisy Bush | | 11 | 2006 |
| Olearia obcordata | Daisy Bush | | 55 | 2006 |
| Olearia persoonioides | Daisy Bush | | 57 | 2005 |
| Olearia phlogopappa | Daisy Bush | | 20 | 2009 |
| Olearia pinifolia | Daisy Bush | | 45 | 2005 |
| Olearia ramulosa | Daisy Bush | | 11 | 2009 |
| Olearia stellulata | Daisy Bush | | 28 | 2008 |
| Olearia tasmanica | Tasmans Daisy Bush | | 51 | 2008 |
| Olearia viscosa | Daisy Bush | | 60 | 2009 |
| Ozothamnus antennaria | Everlasting | | 63 | 2007 |
| Ozothamnus costatifructus | Everlasting | | 70 | 2007 |
| Ozothamnus dendroideus | Everlasting | | 22 | 1984 |
| Ozothamnus ericifolius | Everlasting | | 79 | 2007 |
| Ozothamnus expansifolius | Everlasting | | 65 | 2005 |
| Ozothamnus ferrugineus | Everlasting | | 13 | 2009 |
| Ozothamnus hookeri | Hookers Everlasting | | 32 | 2005 |
| Ozothamnus lycopodioides | Everlasting | | 95 | 2007 |
| Ozothamnus obcordatus | Everlasting | | 9 | 2007 |
| Ozothamnus reflexifolius | Reflexed Everlasting | Vulnerable | 100 | 2007 |
| Ozothamnus reticulatus | Everlasting | | 97 | 2007 |
| Ozothamnus rodwayi | Rodways Everlasting | | 59 | 2007 |
| Ozothamnus rosmarinifolius | Rosemary-leaved Everlasting | | 18 | 2003 |
| Ozothamnus scutellifolius | Everlasting | | 87 | 2008 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|------------------------|-------------|---|--------------------|
| Ozothamnus thyrsoides | Everlasting | | 17 | 2006 |
| Ozothamnus turbinatus | Everlasting | | 10 | 2004 |
| Pappochroma bellidioides | Native Daisy | | 14 | 2004 |
| Pappochroma gunnii | Native Daisy | | 54 | 2005 |
| Pappochroma setosum | Native Daisy | | 14 | 1983 |
| Pappochroma stellatum | Native Daisy | | 64 | 2000 |
| Pappochroma tasmanicum | Native Daisy | | 37 | 2001 |
| Pappochroma trigonum | Native Daisy | | 50 | 2000 |
| Picris angustifolia | | | 4 | 1992 |
| Podolepis jaceoides | Native Daisy | | 6 | 2006 |
| Pseudognaphalium luteoalbum | Native Daisy | | 3 | 2009 |
| Pterygopappus lawrencei | Native Daisy | | 58 | 2007 |
| Rhodanthe anthemoides | Paper-daisy | | 2 | 2005 |
| Rutidosis multiflora | Native Daisy | | <1 | 1992 |
| Senecio albogilvus | Senecio | | 71 | 2005 |
| Senecio biserratus | Senecio | | 14 | 2006 |
| Senecio elegans | Senecio | | 3 | 2004 |
| Senecio extensus | Senecio | | 4 | 1984 |
| Senecio glomeratus | Senecio | | 7 | 2007 |
| Senecio gunnii | Senecio | | 20 | 2007 |
| Senecio hispidissimus | Senecio | | 7 | 1984 |
| Senecio hispidulus | Senecio | | 10 | 2007 |
| Senecio lautus | Senecio | | 7 | 1999 |
| Senecio leptocarpus | Senecio | | 56 | 2000 |
| Senecio linearifolius | Senecio | | 9 | 2009 |
| Senecio macrocarpus | Large-fruit Groundsel | Vulnerable | 2 | 1991 |
| Senecio microbasis | Senecio | | 12 | 2005 |
| Senecio minimus | Senecio | | 11 | 2009 |
| Senecio odoratus | Senecio | | 3 | 1988 |
| Senecio papillosus | Senecio | | 100 | 1993 |
| Senecio pectinatus | Senecio | | 44 | 2005 |
| Senecio phelleus | Senecio | | 2 | 2006 |
| Senecio pinnatifolius | Senecio | | 3 | 2006 |
| Senecio prenanthoides | Senecio | | 2 | 2006 |
| Senecio primulaefolius | Senecio | | 100 | 1993 |
| Senecio quadridentatus | Senecio | | 4 | 2007 |
| Senecio squarrosus | Senecio | | 10 | 2009 |
| Senecio velleioides | Senecio | | 7 | 2009 |
| Senecio vulgaris | Senecio | | 8 | 2008 |
| Sigesbeckia orientalis | Native Daisy | | <1 | 2005 |
| Solenogyne dominii | Native Daisy | | 9 | 2007 |
| Solenogyne gunnii | Native Daisy | | 22 | 2008 |
| Sonchus hydrophilus | Native Sowthistle | | <1 | 1982 |
| Taraxacum aristum | Native Dandelion | | 24 | 1994 |
| Taraxacum cygnorum | Coast Dandelion | Vulnerable | 20 | 1990 |
| Tragopogon porrifolius | Introduced Goats-beard | | 4 | 2004 |
| Triptilodiscus pygmaeus | Native Daisy | | <1 | 2008 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------|--------------------------|---------------------------------|---|--------------------|
| <i>Vellereophyton dealbatum</i> | Native Daisy | | 8 | 2007 |
| <i>Vittadinia burbridgeae</i> | Smooth New Holland Daisy | | 75 | 2009 |
| <i>Vittadinia cuneata</i> | Native Daisy | | 1 | 2008 |
| <i>Vittadinia dissecta</i> | Native Daisy | | <1 | 1983 |
| <i>Vittadinia gracilis</i> | Native Daisy | | 2 | 2009 |
| <i>Vittadinia muelleri</i> | Native Daisy | | 6 | 2009 |
| <i>Xerochrysum bicolor</i> | Native Daisy | | 5 | 1996 |
| <i>Xerochrysum bracteatum</i> | Native Daisy | | <1 | 1997 |
| <i>Xerochrysum collierianum</i> | Native Daisy | | 21 | 1995 |
| <i>Xerochrysum palustre</i> | Native Daisy | Vulnerable | 10 | 2007 |
| <i>Xerochrysum papillosa</i> | Native Daisy | | 27 | 2007 |
| <i>Xerochrysum subundulata</i> | Native Daisy | | 10 | 2004 |
| <i>Xerochrysum viscosum</i> | Native Daisy | | <1 | 2002 |
| Menyanthaceae | | Marshwort | (5 out of 29 species in Australia) | |
| <i>Liparophyllum gunnii</i> | Liparophyllum | | 56 | 1994 |
| <i>Nymphoides exigua</i> | Marshwort | | 39 | 2007 |
| <i>Nymphoides montana</i> | Entire Marshwort | | 2 | 2008 |
| <i>Villarsia exaltata</i> | Marshwort | | <1 | 1983 |
| <i>Villarsia reniformis</i> | Running Marshflower | | 11 | 2007 |
| Stylidiaceae | | Triggerplants and Allies | (8 out of 293 species in Australia) | |
| <i>Donatia novae-zelandiae</i> | | | 60 | 2007 |
| <i>Forstera bellidifolia</i> | | | 71 | 2006 |
| <i>Levenhookia dubia</i> | Hairy Stylewort | | <1 | |
| <i>Phyllachne colensoi</i> | | | 56 | 1996 |
| <i>Stylidium armeria</i> | | | 15 | 2005 |
| <i>Stylidium despectum</i> | Dwarf Triggerplant | | 1 | 1986 |
| <i>Stylidium dilatatum</i> | | | 25 | 2004 |
| <i>Stylidium graminifolium</i> | Grass Trigger-plant | | 15 | 2009 |
| Canellales | | Pepper Trees | (1 out of 11 species in Australia) | |
| <i>Tasmania lanceolata</i> | Mountain Pepper | | 23 | 2008 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------------|------------------------------|---|---|--------------------|
| Caryophyllales | | | | |
| Aizoaceae | | | | |
| | Pigfaces | (3 out of 44 species in Australia) | | |
| Carpobrotus rossii | Native Pig Face | | 9 | 2007 |
| Disphyma crassifolium | Round-leaved Pigface | | 1 | 2007 |
| Tetragonia implexicoma | Bower Spinach | | 6 | 2007 |
| Amaranthaceae | | | | |
| | Pigweed | (3 out of 157 species in Australia) | | |
| Alternanthera nodiflora | Common Joyweed | | <1 | 1998 |
| Hemichroa pentandra | Trailing Jointweed | | 7 | 2006 |
| Ptilotus spathulatus | Pussy Tails | | 3 | 2008 |
| Caryophyllaceae | | | | |
| | Pin Cushion, Wort | (16 out of 42 species in Australia) | | |
| Colobanthus affinis | Alpine Colobanth | | 35 | 2004 |
| Colobanthus apetalus var. apetalus | Tufted Cushion Plant | | 40 | 1986 |
| Colobanthus curtisiae | Grassland Hopflower | Vulnerable | 67 | 2009 |
| Colobanthus muscoides | Coastal cushion plants | | 100 | 2002 |
| Colobanthus pulvinatus | Cushion plant eyebright | | 46 | 1993 |
| Gypsophila tubulosa | Chalkwort | | <1 | 1996 |
| Sagina diemensis | Pearlwort | Critically endangered | 100 | 2005 |
| Sagina namadgi | Namadgi | | 5 | 2006 |
| Scleranthus biflorus | Two flowered Knawel | | 22 | 2009 |
| Scleranthus brockiei | Brock knawel | | 43 | 2009 |
| Scleranthus diander | Knawel | | 8 | 2009 |
| Scleranthus fasciculatus | Kanwel | | 30 | 2009 |
| Stellaria angustifolia | Starwort | | 1 | 2006 |
| Stellaria flaccida | Forest Starwort | | 4 | 2005 |
| Stellaria multiflora | Rayless Starwort | | 16 | 2009 |
| Stellaria pungens | Prickly Starwort | | 1 | 1996 |
| Chenopodiaceae | | | | |
| | Saltbushes and Allies | (13 out of 291 species in Australia) | | |
| Atriplex australasica | Native Orache | | 8 | 2004 |
| Atriplex billarderei | Glistening Saltbush | | 22 | 1998 |
| Atriplex cinerea | Grey Saltbush | | 2 | 2006 |
| Atriplex paludosa | Marsh Saltbush | | 1 | 1997 |
| Atriplex suberecta | Lagoon Saltbush | | <1 | 1999 |
| Chenopodium glaucum | Glaucous Goosefoot | | 8 | 2007 |
| Dysphania pumilio | Clammy Goosefoot | | 1 | 2005 |
| Einadia nutans | Climbing Saltbush | | 1 | 2008 |
| Rhagodia candolleana | Seaberry saltbush | | 5 | 2007 |
| Sarcocornia blackiana | Thick-head samphire | | 4 | 2007 |
| Sarcocornia quinqueflora | Beaded Glasswort | | 6 | 2007 |
| Sclerostegia arbuscula | Shrubby Glasswort | | 6 | 2006 |
| Suaeda australis | Austral Seablite | | 5 | 2007 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|---|-------------|---|--------------------|
| Droseraceae | Sundews | | (7 out of 117 species in Australia) | |
| <i>Drosera arcturi</i> | Alpine Sundew | | 42 | 2008 |
| <i>Drosera auriculata</i> | Sundew | | 8 | 2005 |
| <i>Drosera binata</i> | Sundew | | 14 | 2008 |
| <i>Drosera macrantha</i> | Bridal Rainbow, Climbing Sundew | | 1 | 2006 |
| <i>Drosera peltata</i> | Pale Sundew | | 4 | 2006 |
| <i>Drosera pygmaea</i> | Tiny Sundew | | 12 | 2008 |
| <i>Drosera spatulata</i> | Sundew | | 4 | 2007 |
| Portulacaceae | Purslane | | (5 out of 73 species in Australia) | |
| <i>Calandrinia eremaea</i> | Twining Purslane | | <1 | 1994 |
| <i>Calandrinia granulifera</i> | Pygmy Purslane | | <1 | 1992 |
| <i>Montia australasica</i> | White Purslane | | 33 | 1990 |
| <i>Montia fontana</i> | Water blinks | | 9 | 2002 |
| <i>Neopaxia australasica</i> | White Purslane | | 11 | 2004 |
| Commelinales | Bloodroots, Conostyles, Kangaroo Paws and their Allies | | | |
| Haemodoraceae | | | | |
| <i>Haemodorum distichophyllum</i> | Native Bloodroot | | 82 | 2008 |
| Dilleniales | Guinea Flowers | | (13 out of 253 species in Australia) | |
| Dilleniaceae | | | | |
| <i>Hibbertia acicularis</i> | | | 6 | 2007 |
| <i>Hibbertia appressa</i> | | | 23 | 2005 |
| <i>Hibbertia aspera</i> | | | <1 | 1992 |
| <i>Hibbertia basaltica</i> | Basalt Guinea-flower | Endangered | 100 | 2009 |
| <i>Hibbertia empetrifolia</i> | | | 9 | 2009 |
| <i>Hibbertia fasciculata</i> | | | 8 | 2009 |
| <i>Hibbertia hirsuta</i> | | | 50 | 2007 |
| <i>Hibbertia hirticalyx</i> | | | 8 | 2004 |
| <i>Hibbertia procumbens</i> | Spreading Guinea Flower | | 26 | 2009 |
| <i>Hibbertia riparia</i> | Erect Guinea-flower | | 7 | 2009 |
| <i>Hibbertia sericea</i> | | | 1 | 2007 |
| <i>Hibbertia serpyllifolia</i> | | | 14 | 2006 |
| <i>Hibbertia stricta</i> | | | 2 | 1979 |
| Dioscoreales | Fairy Lanterns | | (1 out of 3 species in Australia) | |
| Thismiaceae | | | | |
| <i>Thismia rodwayi</i> | Fairy Lanterns | | 50 | 2009 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-----------------------|---|--------------------|
| Ericales | | | |
| Epacridaceae | | | |
| Southern Heaths (86 out of 458 species in Australia) | | | |
| Acrotriche serrulata | | 8 | 2008 |
| Archeria comberi | | 65 | 2005 |
| Archeria eriocarpa | | 36 | 1994 |
| Archeria hirtella | | 42 | 1997 |
| Archeria serpyllifolia | | 70 | 2005 |
| Astroloma humifusum | | 8 | 2009 |
| Astroloma pinifolium | | 14 | 2006 |
| Brachyloma ciliatum | | 16 | 2006 |
| Brachyloma depressum | | 25 | 2009 |
| Cyathodes platystoma | | 100 | 2009 |
| Dracophyllum milliganii | | 74 | 2005 |
| Dracophyllum minimum | | 69 | 2005 |
| Epacris acuminata | Vulnerable | 83 | 2009 |
| Epacris apsleyensis | Endangered | 60 | 2009 |
| Epacris barbata | Critically endangered | 100 | 2004 |
| Epacris corymbiflora | | 60 | 2007 |
| Epacris exserta | Endangered | 15 | 1986 |
| Epacris franklinii | | 6 | 2002 |
| Epacris grandis | Endangered | 50 | 2000 |
| Epacris gunnii | | 30 | 2009 |
| Epacris heteronema | | 54 | 2004 |
| Epacris impressa | | 14 | 2009 |
| Epacris lanuginosa | | 31 | 2007 |
| Epacris limbata | Critically endangered | 67 | 2004 |
| Epacris marginata | | 100 | 2006 |
| Epacris microphylla | | 1 | 1965 |
| Epacris moscaliana | | 8 | 1980 |
| Epacris mucronulata | | 46 | 2004 |
| Epacris myrtifolia | | 88 | 2008 |
| Epacris navicularis | | 60 | 1998 |
| Epacris obtusifolia | | 9 | 2008 |
| Epacris petrophila | | 28 | 2000 |
| Epacris serpyllifolia | | 54 | 2007 |
| Epacris stuartii | Critically endangered | 100 | 2001 |
| Epacris tasmanica | | 78 | 2008 |
| Epacris virgata | Endangered | 69 | 2009 |
| Leptecophylla abietina | | 36 | 2007 |
| Leptecophylla dealbata | | 64 | 2007 |
| Leptecophylla divaricata | | 80 | 2008 |
| Leptecophylla glauca | | 60 | 2007 |
| Leptecophylla juniperina | | 42 | 2007 |
| Leptecophylla pendulosa | | 39 | 2004 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|--------------------------|-------------|---|--------------------|
| <i>Leptecophylla pogonocalyx</i> | Heath | | 66 | 2006 |
| <i>Leptecophylla straminea</i> | Heath | | 48 | 2006 |
| <i>Leucopogon australis</i> | Spiked Beard-heath | | 4 | 2004 |
| <i>Leucopogon collinus</i> | Fringed Beard-heath | | 39 | 2009 |
| <i>Leucopogon ericoides</i> | Pink Beard-heath | | 13 | 2009 |
| <i>Leucopogon fraseri</i> | Sharp Beard-heath | | 19 | 2007 |
| <i>Leucopogon lanceolatus</i> | Lance-leaved Beard-heath | | <1 | 1979 |
| <i>Leucopogon milliganii</i> | Milligans Beard-heath | | 50 | 1998 |
| <i>Leucopogon oreophilus</i> | Beard-heath | | 50 | 1999 |
| <i>Leucopogon parviflorus</i> | Coastal Beard-heath | | 9 | 2008 |
| <i>Leucopogon pilifer</i> | Thready Beard-heath | | 32 | 2005 |
| <i>Leucopogon virgatus</i> | Beard-heath | | 8 | 2009 |
| <i>Lissanthe</i> sp. <i>diggers camp</i> | Beard-heath | | 17 | |
| <i>Lissanthe strigosa</i> | Beard-heath | | 11 | 2009 |
| <i>Monotoca elliptica</i> | Broom-heath | | 7 | 2008 |
| <i>Monotoca empetrifolia</i> | Broom-heath | | 54 | 2004 |
| <i>Monotoca glauca</i> | Goldy-wood | | 31 | 2009 |
| <i>Monotoca linifolia</i> | Broom-heath | | 63 | 2005 |
| <i>Monotoca scoparia</i> | Prickly Broom-heath | | <1 | 2003 |
| <i>Monotoca submutica</i> | Broom-heath | | 49 | 2006 |
| <i>Pentachondra ericifolia</i> | Heath | | 69 | 2005 |
| <i>Pentachondra involucrata</i> | Heath | | 73 | 2005 |
| <i>Pentachondra pumila</i> | Heath | | 47 | 2007 |
| <i>Planocarpa nitida</i> | Heath | | 62 | 1992 |
| <i>Planocarpa petiolaris</i> | Heath | | 62 | 2005 |
| <i>Planocarpa sulcata</i> | Heath | | 46 | 1997 |
| <i>Prionotes cerinthoides</i> | Heath | | 59 | 2008 |
| <i>Richea acerosa</i> | Grass-tree | | 59 | 2007 |
| <i>Richea alpina</i> | Grass-tree | | 83 | 1990 |
| <i>Richea continentis</i> | Grass-tree | | 1 | 1974 |
| <i>Richea dracophylla</i> | Grass-tree | | 94 | 2006 |
| <i>Richea gunnii</i> | Grass-tree | | 49 | 2007 |
| <i>Richea milliganii</i> | Grass-tree | | 76 | 2008 |
| <i>Richea pandanifolia</i> | Giant Grass-tree | | 56 | 2008 |
| <i>Richea procera</i> | Grass-tree | | 80 | 2009 |
| <i>Richea scoparia</i> | Grass-tree | | 50 | 2007 |
| <i>Richea sprengelioides</i> | Grass-tree | | 60 | 2007 |
| <i>Sprengelia distichophylla</i> | Heath | | 60 | 2005 |
| <i>Sprengelia incarnata</i> | Pink Swamp Heath | | 21 | 2099 |
| <i>Styphelia adscendens</i> | Golden Heath | | 23 | 2007 |
| <i>Trochocarpa cunninghamii</i> | Heath | | 39 | 2008 |
| <i>Trochocarpa disticha</i> | Heath | | 75 | 2005 |
| <i>Trochocarpa gunnii</i> | Heath | | 47 | 2008 |
| <i>Trochocarpa thymifolia</i> | Heath | | 42 | 2005 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|-----------------------------------|-------------|---|--------------------|
| Ericaceae | Heathers and Rhododendrons | | (8 out of 17 species in Australia) | |
| <i>Acrothamnus hookeri</i> | Mountain Beard-Heath | | 16 | 2007 |
| <i>Acrothamnus montanus</i> | Snow Beard-Heath | | 40 | 2009 |
| <i>Gaultheria appressa</i> | Waxberry | | <1 | 1988 |
| <i>Gaultheria depressa</i> | Waxberry | | 54 | 2002 |
| <i>Gaultheria hispida</i> | Waxberry | | 47 | 2006 |
| <i>Gaultheria tasmanica</i> | Waxberry | | 55 | 2006 |
| <i>Pernettya lanceolata</i> | Lanceolate Waxberry | | 67 | 1994 |
| <i>Pernettya tasmanica</i> | Tasmanian Waxberry | | 42 | 1992 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Fabales | | | |
| Fabaceae | | | |
| Peas (94 out of 2583 species in Australia) | | | |
| Acacia aspera | | 1 | |
| Acacia axillaris | Vulnerable | 53 | 2003 |
| Acacia baileyana | | 3 | 2007 |
| Acacia betchei | | 3 | 1981 |
| Acacia boormanii | | 2 | 2004 |
| Acacia calamifolia | | <1 | |
| Acacia dealbata | | 12 | 2009 |
| Acacia decurrens | | <1 | 2007 |
| Acacia floribunda | | <1 | 2002 |
| Acacia genistifolia | | 13 | 2009 |
| Acacia gunnii | | 8 | 2006 |
| Acacia howittii | | 11 | 2006 |
| Acacia leucolobia | | 14 | 1989 |
| Acacia longifolia | | <1 | 2006 |
| Acacia mearnsii | | 7 | 2009 |
| Acacia melanoxylon | | 8 | 2009 |
| Acacia mucronata | | 15 | 2009 |
| Acacia myrtifolia | | 7 | 2008 |
| Acacia paradoxa | | <1 | 2001 |
| Acacia pataczekii | | 21 | 2002 |
| Acacia pravissima | | 6 | 2007 |
| Acacia pycnantha | | <1 | 2006 |
| Acacia retinodes | | 2 | 2004 |
| Acacia riceana | | 52 | 2007 |
| Acacia siculiformis | | 10 | 2009 |
| Acacia sophorae | | 7 | 2008 |
| Acacia stricta | | 11 | 2009 |
| Acacia suaveolens | | 7 | 2008 |
| Acacia terminalis | | 6 | 2008 |
| Acacia ulicifolia | | 3 | 2009 |
| Acacia verniciflua | | 9 | 2009 |
| Acacia verticillata | | 16 | 2009 |
| Almaleea subumbellata | | 30 | 2005 |
| Aotus ericoides | | 18 | 2009 |
| Bossiaea cinerea | | 22 | 2009 |
| Bossiaea cordigera | | 22 | 2009 |
| Bossiaea ensata | | <1 | |
| Bossiaea obcordata | | <1 | 2002 |
| Bossiaea prostrata | | 13 | 2009 |
| Bossiaea riparia | | 41 | 2009 |
| Cassia artemisioides | | 5 | 1993 |
| Chamaecytisus palmensis | | 92 | 1999 |
| Daviesia latifolia | | 8 | 2009 |
| Daviesia sejugata | | 62 | 2007 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------|-------------------------------|-------------|---|--------------------|
| <i>Daviesia ulicifolia</i> | Bitter-pea | | 7 | 2009 |
| <i>Desmodium gunnii</i> | Gunns Tick-trefoil | | <1 | 2006 |
| <i>Desmodium intortum</i> | Tick-trefoil | | 2 | 1968 |
| <i>Desmodium varians</i> | Tick-trefoil | | <1 | 2004 |
| <i>Dillwynia cinerascens</i> | Parrot-pea | | 16 | 2009 |
| <i>Dillwynia glaberrima</i> | Parrot-pea | | 9 | 2008 |
| <i>Dillwynia hispida</i> | Parrot-pea | | 2 | 2008 |
| <i>Dillwynia retorta</i> | Parrot-pea | | <1 | 0002 |
| <i>Dillwynia rudis</i> | Parrot-pea | | 8 | 1998 |
| <i>Eutaxia microphylla</i> | Small-leaved Bacon and Eggs | | <1 | 1988 |
| <i>Glycine clandestina</i> | Widespread Glycine | | <1 | 2007 |
| <i>Glycine latrobeana</i> | Clover Glycine | Vulnerable | 8 | 2008 |
| <i>Glycine microphylla</i> | Small-leaved Glycine | | <1 | 2006 |
| <i>Gompholobium huegelii</i> | Pale Wedge-pea | | 9 | 2008 |
| <i>Goodia lotifolia</i> | Native Clover | | 4 | 2006 |
| <i>Hardenbergia violacea</i> | Happy Wanderer | | <1 | 2009 |
| <i>Hovea heterophylla</i> | Hovea | | 4 | 2009 |
| <i>Hovea linearis</i> | Thin-leaved Hovea | | 13 | 1999 |
| <i>Hovea longifolia</i> | Long-leaved Hovea | | 2 | 1990 |
| <i>Hovea magnibractea</i> | Hovea | | 20 | 1978 |
| <i>Hovea montana</i> | Montane Hovea | | 13 | 2007 |
| <i>Hovea tasmanica</i> | Hovea | | 57 | 2009 |
| <i>Indigofera australis</i> | Native Indigo | | 3 | 2009 |
| <i>Kennedia prostrata</i> | Running Postman | | 4 | 2009 |
| <i>Kennedia rubicunda</i> | Red Kennedy Pea | | <1 | 2001 |
| <i>Lotus australis</i> | Australian Trefoil | | <1 | 2007 |
| <i>Mirbelia oxylobioides</i> | Mirbelia | | <1 | 1999 |
| <i>Neonotonia wightii</i> | Wights Glycine | | 1 | 1968 |
| <i>Oxylobium arborescens</i> | Golden Shaggy-pea | | 4 | 2005 |
| <i>Oxylobium ellipticum</i> | Golden Shaggy-pea | | 35 | 2009 |
| <i>Phyllota diffusa</i> | Bush-pea | | 24 | 2000 |
| <i>Platylobium formosum</i> | Native Pea | | <1 | 2009 |
| <i>Platylobium obtusangulum</i> | Native Pea | | 3 | 2000 |
| <i>Platylobium triangulare</i> | Native Pea | | 26 | 2009 |
| <i>Pultenaea daphnoides</i> | Bacon and Eggs | | 12 | 2009 |
| <i>Pultenaea dentata</i> | Bacon and Eggs | | 17 | 2007 |
| <i>Pultenaea fasciculata</i> | Bacon and Eggs | | 23 | 2000 |
| <i>Pultenaea gunnii</i> | Bacon and Eggs | | 19 | 2007 |
| <i>Pultenaea humilis</i> | Bacon and Eggs | | 2 | 2009 |
| <i>Pultenaea juniperina</i> | Juniper-leaved Bacon and Eggs | | 25 | 2009 |
| <i>Pultenaea pedunculata</i> | Matted Bush Pea | | 14 | 2008 |
| <i>Pultenaea prostrata</i> | Prostrate Bacon and Eggs | | 2 | 2005 |
| <i>Pultenaea stricta</i> | Bacon and Eggs | | 11 | 2006 |
| <i>Pultenaea tenuifolia</i> | Bacon and Eggs | | <1 | 1992 |
| <i>Sphaerolobium minus</i> | Reed Pea | | 14 | 2006 |
| <i>Sphaerolobium vimineum</i> | Reed Pea | | 1 | 2001 |
| <i>Stonesiella selaginoides</i> | Bush-pea | Endangered | 62 | 1998 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|-------------------|-------------|---|--------------------|
| Swainsona lessertiifolia | Darling Pea | | 1 | 2006 |
| Trifolium glomeratum | Introduced Clover | | 4 | 2004 |
| Viminaria juncea | Golden Spray | | <1 | 2007 |
| Fagales | | | | |
| Casuarinaceae | | | | |
| She-oaks (7 out of 69 species in Australia) | | | | |
| Allocasuarina crassa | She-oak | | 100 | 1998 |
| Allocasuarina duncanii | Duncan's Sheoak | Vulnerable | 100 | 2009 |
| Allocasuarina littoralis | Black Sheoak | | 6 | 2009 |
| Allocasuarina monilifera | She-oak | | 35 | 2008 |
| Allocasuarina paludosa | Scrub She-oak | | <1 | |
| Allocasuarina verticillata | Drooping Sheoak | | 8 | 2009 |
| Allocasuarina zephyrea | She-oak | | 23 | 2005 |
| Nothofagaceae (3 out of 3 species in Australia) | | | | |
| Nothofagus cunninghamii | Myrtle Beech | | 34 | 2005 |
| Nothofagus gunnii | Tanglefoot Beech | | 38 | 2006 |
| Nothofagus moorei | Antarctic Beech | | 2 | 1992 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|--|---|--------------------|
| Lamiales | | | |
| Lentibulariaceae | | | |
| Bladderworts (5 out of 58 species in Australia) | | | |
| Utricularia australis | | <1 | 1990 |
| Utricularia dichotoma | | 9 | 2008 |
| Utricularia lateriflora | | 14 | 2008 |
| Utricularia monanthos | | 30 | 2007 |
| Utricularia uniflora | | 11 | 2006 |
| Phrymaceae (3 out of 21 species in Australia) | | | |
| Glossostigma elatinoides | | 2 | 1999 |
| Mazus pumilio | | 11 | 2005 |
| Mimulus repens | | 7 | 2004 |
| Scrophulariaceae | | | |
| Figworts (21 out of 99 species in Australia) | | | |
| Euphrasia amphisysepala | Vulnerable | 100 | 2008 |
| Euphrasia collina | | 11 | 2008 |
| Euphrasia fragosa | Shy Eyebright Critically endangered | 100 | 2007 |
| Euphrasia gibbsiae | | 48 | 2007 |
| Euphrasia hookeri | | 47 | 1998 |
| Euphrasia phragmostoma | Vulnerable | 100 | 2008 |
| Euphrasia scabra | | 4 | 2008 |
| Euphrasia semipicta | Peninsula Eyebright | 100 | 2008 |
| Euphrasia sp. bivouac bay | Endangered | 100 | 2008 |
| Gratiola nana | | 25 | 2007 |
| Gratiola peruviana | | 1 | 2002 |
| Limosella australis | | 6 | 2004 |
| Ourisia integrifolia | | 56 | 2006 |
| Veronica calycina | Hairy Speedwell | 10 | 2009 |
| Veronica continua | | 100 | 2007 |
| Veronica densifolia | | 20 | |
| Veronica formosa | | 60 | 2007 |
| Veronica gracilis | | 18 | 2009 |
| Veronica novae-hollandiae | | 23 | 1997 |
| Veronica plebeia | | <1 | 2004 |
| Veronica serpyllifolia | | 11 | 2004 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|-----------------------|-------------|---|--------------------|
| Laurales | | | | |
| Atherospermataceae (1 out of 10 species in Australia) | | | | |
| Atherosperma moschatum | Tasmanian Sassafras | | 27 | 2009 |
| Lauraceae Laurels, Camphorwood and Australian Walnuts (3 out of 140 species in Australia) | | | | |
| Cassytha glabella | Tangled Dodder Laurel | | 4 | 2008 |
| Cassytha melantha | Dodder Laurel | | <1 | 2006 |
| Cassytha pubescens | Dodder Laurel | | 5 | 2008 |
| Liliales | | | | |
| Campynemataceae (1 out of 1 species in Australia) | | | | |
| Campynema lineare | Native Lily | | 61 | 2007 |
| Colchicaceae Nancies and Allies (3 out of 48 species in Australia) | | | | |
| Burchardia umbellata | Milk-maids | | 1 | 2004 |
| Wurmbea dioica | Early Nancy | | 1 | 2005 |
| Wurmbea uniflora | Nancy | | 30 | 2005 |
| Luzuriagaceae Orange and Turquoise Berries (1 out of 2 species in Australia) | | | | |
| Drymophila cyanocarpa | Turquoise Berry | | 26 | 2008 |
| Philesiaceae (1 out of 1 species in Australia) | | | | |
| Eustrephus latifolius | Wombat Berry | | <1 | |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--|-------------|---|--------------------|
| Malpighiales | | | |
| Euphorbiaceae | | | |
| Spurges (7 out of 315 species in Australia) | | | |
| Amperea xiphioclada | | 14 | 2009 |
| Bertya rosmarinifolia | | 6 | 1990 |
| Beyeria lechenaultii | | <1 | |
| Beyeria viscosa | | 18 | 2008 |
| Euphorbia drummondii | | <1 | |
| Euphorbia helioscopia | | 5 | 2004 |
| Ricinocarpos pinifolius | | 2 | 2004 |
| Phyllanthaceae | | | |
| (4 out of 113 species in Australia) | | | |
| Phyllanthus australis | | 29 | 2007 |
| Phyllanthus gunnii | | 3 | 2005 |
| Poranthera microphylla | | 7 | 2009 |
| Poranthera petalifera | Vulnerable | 100 | 2005 |
| Picrodendraceae | | | |
| (3 out of 29 species in Australia) | | | |
| Micrantheum hexandrum | | 12 | 2008 |
| Pseudanthus divaricatissimus | | 1 | 1996 |
| Pseudanthus ovalifolius | | 17 | 2003 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Malvales | | | |
| Sterculiaceae | | | |
| Kurrajongs and Bottle-trees | | | |
| (2 out of 273 species in Australia) | | | |
| Lasiopetalum macrophyllum | | 3 | 2007 |
| Lasiopetalum micranthum | Vulnerable | 88 | 2008 |
| Thymelaeaceae | | | |
| Pimeleas | | | |
| (17 out of 110 species in Australia) | | | |
| Kelleria dieffenbachii | | 45 | 1992 |
| Pimelea cinerea | | 38 | 2005 |
| Pimelea curviflora | | 1 | 2009 |
| Pimelea drupacea | | 35 | 2009 |
| Pimelea filiformis | | 6 | 2005 |
| Pimelea flava | | 6 | 2006 |
| Pimelea glauca | | 9 | 2009 |
| Pimelea ligustrina | | 6 | 1999 |
| Pimelea linifolia | | 5 | 2007 |
| Pimelea micrantha | | <1 | 1984 |
| Pimelea milliganii | | 43 | 1989 |
| Pimelea neo-anglica | | 2 | 2006 |
| Pimelea nivea | | 58 | 2008 |
| Pimelea pygmaea | | 82 | 2000 |
| Pimelea sericea | | 66 | 2005 |
| Pimelea serpyllifolia | | <1 | 1974 |
| Pimelea stricta | | <1 | 1978 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------|---------------------|---|---|--------------------|
| Myrtales | | | | |
| Lythraceae | | | | |
| | Loosestrifes | (2 out of 27 species in Australia) | | |
| Lythrum hyssopifolia | Loosestrife | | <1 | 2007 |
| Lythrum salicaria | Loosestrife | | 1 | 2008 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|----------------------------|--|---|--------------------|
| Myrtaceae | Native Myrtles | (78 out of 2211 species in Australia) | | |
| Angophora hispida | Dwarf Apple | | 2 | 2005 |
| Baeckea gunniana | Alpine Baeckea | | 28 | 2007 |
| Baeckea leptocaulis | Baeckea | | 42 | 2008 |
| Baeckea sp. walkaway | Baeckea | | 3 | |
| Callistemon citrinus | Crimson Bottlebrush | | <1 | 2002 |
| Callistemon pallidus | Lemon Bottlebrush | | 29 | 2009 |
| Callistemon rigidus | Bottlebrush | | 1 | |
| Callistemon rugulosus | Scarlet Bottlebrush | | <1 | |
| Callistemon viridiflorus | Green Bottlebrush | | 58 | 2006 |
| Calytrix tetragona | Star flower | | 1 | 2008 |
| Corymbia citriodora | Lemon-scented gum | | <1 | 1961 |
| Eucalyptus amygdalina | Black Peppermint | | 40 | 2099 |
| Eucalyptus archeri | Alpine Cider Gum | | 37 | 2004 |
| Eucalyptus barberi | Barbers Gum | | 80 | 2009 |
| Eucalyptus bicostata | Southern Blue Gum | | 24 | 2009 |
| Eucalyptus botryoides | Bangalay | | <1 | 2006 |
| Eucalyptus bridgesiana | Apple Gum | | <1 | 2006 |
| Eucalyptus brookeriana | Brookers Gum | | 25 | 2008 |
| Eucalyptus camaldulensis | River Red Gum | | <1 | 1989 |
| Eucalyptus coccifera | Tasmanian Snow Gum | | 66 | 2007 |
| Eucalyptus cordata | Heart-leaved silver gum | | 94 | 2009 |
| Eucalyptus crenulata | Buxton Gum | Endangered | 3 | 1979 |
| Eucalyptus dalrympleana | Mountain Gum | | 16 | 2009 |
| Eucalyptus delegatensis | Alpine Ash | | 30 | 2005 |
| Eucalyptus globoidea | White Stringybark | | <1 | 2005 |
| Eucalyptus globulus | Southern Blue Gum | | 23 | 2009 |
| Eucalyptus grandis | Flooded Gum | | <1 | 2005 |
| Eucalyptus gunnii | Cider gum | | 54 | 2009 |
| Eucalyptus johnstonii | Johnstons Gum | | 88 | 2005 |
| Eucalyptus leucoxydon | Yellow gum | | <1 | 1995 |
| Eucalyptus morrisbyi | Morrisbys Gum | Endangered | 89 | 2004 |
| Eucalyptus nitens | Shining gum | | <1 | 2005 |
| Eucalyptus nitida | Smithton peppermint | | 34 | 2007 |
| Eucalyptus nortonii | Long-leaved Box | | <1 | 1992 |
| Eucalyptus obliqua | Messmate | | 13 | 2099 |
| Eucalyptus ovata | Swamp Gum | | 15 | 2009 |
| Eucalyptus pauciflora | Snow Gum | | 11 | 2009 |
| Eucalyptus perriniana | Shinning Gum | | 14 | 2006 |
| Eucalyptus pseudoglobulus | Victorian Eurabbie | | <1 | 1951 |
| Eucalyptus pulchella | Narrow-leaved peppermint | | 80 | 2009 |
| Eucalyptus pulverulenta | Silver-leaved Mountain Gum | Vulnerable | 3 | 1984 |
| Eucalyptus regnans | Mountain Ash | | 29 | 2008 |
| Eucalyptus risdonii | Risdon Peppermint | | 100 | 2006 |
| Eucalyptus rodwayi | Swamp Peppermint | | 56 | 2006 |
| Eucalyptus rubida | Candlebark | | 11 | 2007 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|----------------------------------|-----------------------------|-------------|---|--------------------|
| <i>Eucalyptus sieberi</i> | Silvertop Ash | | 2 | 2007 |
| <i>Eucalyptus stricta</i> | Blue Mountains Mallee Ash | | 1 | 1978 |
| <i>Eucalyptus subcrenulata</i> | Tasmanian Alpine Yellow Gum | | 55 | 2006 |
| <i>Eucalyptus tenuiramis</i> | Silver peppermint | | 89 | 2009 |
| <i>Eucalyptus urnigera</i> | Urn gum | | 86 | 2006 |
| <i>Eucalyptus vernicosa</i> | Varnished Gum | | 68 | 2005 |
| <i>Eucalyptus viminalis</i> | Rough Barked Manna Gum | | 9 | 2009 |
| <i>Euryomyrtus ramosissima</i> | Euryomyrtus | | 18 | 2008 |
| <i>Kunzea ambigua</i> | Tick Bush | | 5 | 2008 |
| <i>Kunzea ericoides</i> | Kanuka | | 2 | 2007 |
| <i>Leptospermum glaucescens</i> | Tea tree | | 36 | 2099 |
| <i>Leptospermum grandiflorum</i> | Tea tree | | 70 | 2008 |
| <i>Leptospermum grandifolium</i> | Mountain Tea tree | | <1 | 1994 |
| <i>Leptospermum laevigatum</i> | Coastal Tea Tree | | 2 | 2007 |
| <i>Leptospermum lanigerum</i> | Silky Tea Tree | | 19 | 2009 |
| <i>Leptospermum nitidum</i> | Shining Tea tree | | 37 | 2008 |
| <i>Leptospermum riparium</i> | Tea tree | | 26 | 2000 |
| <i>Leptospermum rupestre</i> | Creeping Tea Tree | | 49 | 2007 |
| <i>Leptospermum scoparium</i> | Manuka | | 26 | 2099 |
| <i>Lophostemon confertus</i> | Brush Box | | <1 | 1993 |
| <i>Melaleuca armillaris</i> | Bracelet Honey-myrtle | | 1 | 2004 |
| <i>Melaleuca decussata</i> | Cross-leaved Honey-myrtle | | <1 | 2006 |
| <i>Melaleuca ericifolia</i> | Swamp Paperbark | | 3 | 2007 |
| <i>Melaleuca gibbosa</i> | Slender Honey-myrtle | | 11 | 2009 |
| <i>Melaleuca hypericifolia</i> | Hillock Bush | | <1 | |
| <i>Melaleuca paludicola</i> | River Bottlebrush | | 2 | 1984 |
| <i>Melaleuca pustulata</i> | Honey-myrtle | | 81 | 2008 |
| <i>Melaleuca squamea</i> | Swamp Honey-myrtle | | 28 | 2099 |
| <i>Melaleuca squarrosa</i> | Scented Paperbark | | 15 | 2099 |
| <i>Melaleuca wilsonii</i> | Violet Myrtle | | <1 | |
| <i>Syzygium australe</i> | Brush Cherry | | <1 | |
| <i>Thryptomene calycina</i> | Myrtle | | 3 | 2005 |
| <i>Thryptomene micrantha</i> | Myrtle | | 4 | 2008 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|-----------------------------|---|---|--------------------|
| Onagraceae | | Water Primrose and Willowherbs (13 out of 19 species in Australia) | | |
| <i>Epilobium billardierianum</i> | Willowherb | | 5 | 2007 |
| <i>Epilobium brunnescens</i> | Willowherb | | 50 | 2002 |
| <i>Epilobium curtisiae</i> | Willowherb | | 25 | 2002 |
| <i>Epilobium fugitivum</i> | Ben Lomond Willowherb | | 56 | 1996 |
| <i>Epilobium glabellum</i> | Willowherb | | 14 | 1985 |
| <i>Epilobium gunnianum</i> | Willowherb | | 10 | 2004 |
| <i>Epilobium hirtigerum</i> | Hairy Willowherb | | 3 | 2007 |
| <i>Epilobium pallidiflorum</i> | Willowherb | | <1 | 1996 |
| <i>Epilobium pedunculare</i> | Macquarie Island Willowherb | | 100 | 2007 |
| <i>Epilobium perpusillum</i> | Willowherb | | 12 | 1988 |
| <i>Epilobium sarmentaceum</i> | Mountain Willowherb | | 24 | 2007 |
| <i>Epilobium tasmanicum</i> | Tasmanian Willowherb | | 34 | 1991 |
| <i>Epilobium willisii</i> | Willowherb | | 41 | 1996 |
| Nymphaeales | | Hydatellid Herbs (2 out of 9 species in Australia) | | |
| Hydatellaceae | | | | |
| <i>Hydatella filamentosa</i> | Hydatella | | 55 | 2007 |
| <i>Trithuria submersa</i> | Trithuria | | 2 | 2007 |
| Oxalidales | | Christmas Bushes and Mararas (6 out of 37 species in Australia) | | |
| Cunoniaceae | | | | |
| <i>Anodopetalum biglandulosum</i> | Big-glanded Marara | | 36 | 2008 |
| <i>Bauera rubioides</i> | Dog Rose | | 21 | 2099 |
| <i>Eucryphia lucida</i> | | | 37 | 2008 |
| <i>Eucryphia milliganii</i> | | | 64 | 2006 |
| <i>Eucryphia moorei</i> | | | 2 | 1997 |
| <i>Vesselowskyia rubifolia</i> | Southern Marara | | 6 | 1999 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|------------------------|-------------|---|--------------------|
| Pinales | | | | |
| Araucariaceae | | | | |
| | Araucaria Pines | | (2 out of 7 species in Australia) | |
| Araucaria bidwillii | Bunya Pine | | 2 | |
| Araucaria heterophylla | Norfolk Island Pine | | 20 | 1984 |
| Cupressaceae | | | | |
| | Native Cypress | | (6 out of 23 species in Australia) | |
| Athrotaxis cupressoides | Pencil Pine | | 52 | 2007 |
| Athrotaxis selaginoides | King Billy Pine | | 54 | 2005 |
| Callitris monticola | Steelhead | | 17 | 1991 |
| Callitris oblonga | Tasmanian Cypress Pine | Vulnerable | 31 | 2006 |
| Callitris rhomboidea | Dune Cypress Pine | | 14 | 2009 |
| Diselma archeri | Cheshunt Pine | | 56 | 2007 |
| Podocarpaceae | | | | |
| | Podocarp Pines | | (6 out of 15 species in Australia) | |
| Lagarostrobos franklinii | Huon Pine | | 43 | 1999 |
| Microcachrys tetragona | Alpine Dwarf Pine | | 58 | 2007 |
| Pherosphaera hookeriana | Tasmanian Dwarf Pine | | 68 | 2007 |
| Phyllocladus aspleniifolius | Celery-topped Pine | | 39 | 2008 |
| Podocarpus elatus | Brown Pine | | <1 | 1985 |
| Podocarpus lawrencei | Mountain Plum Pine | | 25 | 2005 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---|-------------|---|--------------------|
| Poales | | | |
| Centrolepidaceae | | | |
| Centrolepid Sedges (13 out of 30 species in Australia) | | | |
| Aphelia gracilis | | 2 | |
| Centrolepis aristata | | 4 | 2003 |
| Centrolepis cephaloformis | | <1 | 1960 |
| Centrolepis fascicularis | | 8 | 2006 |
| Centrolepis glabra | | 3 | 1992 |
| Centrolepis monogyna | | 47 | 2008 |
| Centrolepis muscoides | | 71 | 2007 |
| Centrolepis pedderensis | Vulnerable | 71 | 2005 |
| Centrolepis polygyna | | <1 | 2006 |
| Centrolepis strigosa | | 5 | 2006 |
| Gaimardia amblyphylla | | 58 | 2008 |
| Gaimardia fitzgeraldii | | 74 | 2002 |
| Gaimardia setacea | | 57 | 2002 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------|-------------------|---|--------------------|
| Cyperaceae | | | |
| Sedges | | (114 out of 731 species in Australia) | |
| Baumea acuta | | 9 | 2008 |
| Baumea arthropphylla | | 14 | 2006 |
| Baumea articulata | | <1 | 1980 |
| Baumea gunnii | | 6 | 2008 |
| Baumea juncea | | 5 | 2007 |
| Baumea planifolia | | 4 | 2002 |
| Baumea rubiginosa | | 2 | 2004 |
| Baumea tetragona | | 15 | 2008 |
| Bolboschoenus caldwellii | | 3 | 2007 |
| Bolboschoenus medianus | | <1 | 2009 |
| Carex alsophila | | 3 | 1973 |
| Carex appressa | | 7 | 2009 |
| Carex archeri | | 41 | 2005 |
| Carex bichenoviana | | <1 | 1986 |
| Carex breviculmis | | 10 | 2009 |
| Carex canescens | | 15 | 1984 |
| Carex capillacea | | 7 | 2004 |
| Carex cataractae | | 76 | 2005 |
| Carex cephalotes | | 18 | 1982 |
| Carex chlorantha | | 10 | 1995 |
| Carex fascicularis | | 6 | 2006 |
| Carex flaviformis | | 83 | 2005 |
| Carex gaudichaudiana | | 12 | 2006 |
| Carex gunniana | | 18 | 2008 |
| Carex hypandra | | 18 | 1975 |
| Carex inversa | Knob Sedge | 3 | 2004 |
| Carex iynx | | 34 | 2007 |
| Carex longebrachiata | | 6 | 2009 |
| Carex polyantha | | 12 | 2005 |
| Carex pumila | | 7 | 1997 |
| Carex raleighii | | 19 | 1995 |
| Carex tasmanica | Curly Sedge | Vulnerable | 54 |
| Carex tereticaulis | | 5 | 2007 |
| Carex trifida | | 50 | 1993 |
| Carpha alpina | | 50 | 2007 |
| Carpha curvata | | 58 | 2000 |
| Carpha rodwayi | | 66 | 2005 |
| Caustis pentandra | Thick Twist Rush | 3 | 2008 |
| Chorizandra australis | | 17 | 1999 |
| Chorizandra cymbaria | | 3 | 1985 |
| Cyperus gunnii | | <1 | 2001 |
| Cyperus lucidus | | <1 | 2002 |
| Cyperus sanguinolentus | | 2 | 1984 |
| Eleocharis acuta | Sharp Spike Sedge | 5 | 2007 |
| Eleocharis gracilis | | 7 | 2006 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------------------|-------------|---|--------------------|
| <i>Eleocharis pusilla</i> | | 2 | 2007 |
| <i>Eleocharis sphacelata</i> | | 5 | 2005 |
| <i>Gahnia filum</i> | | 12 | 2008 |
| <i>Gahnia grandis</i> | | 37 | 2009 |
| <i>Gahnia microstachya</i> | | 11 | 2007 |
| <i>Gahnia radula</i> | | 8 | 2009 |
| <i>Gahnia rodwayi</i> | | 76 | 2006 |
| <i>Gahnia sieberiana</i> | | <1 | 2006 |
| <i>Gahnia trifida</i> | | 6 | 2007 |
| <i>Gymnoschoenus sphaerocephalus</i> | | 21 | 2009 |
| <i>Isolepis alpina</i> | | 68 | 2006 |
| <i>Isolepis aucklandica</i> | | 26 | 2007 |
| <i>Isolepis cernua</i> | | 7 | 2007 |
| <i>Isolepis crassiuscula</i> | | 26 | 2007 |
| <i>Isolepis fluitans</i> | | 11 | 2007 |
| <i>Isolepis gaudichaudiana</i> | | 5 | 1986 |
| <i>Isolepis habra</i> | | 5 | 1996 |
| <i>Isolepis hookeriana</i> | | 2 | 1996 |
| <i>Isolepis inundata</i> | | 6 | 2007 |
| <i>Isolepis montivaga</i> | | 28 | 2007 |
| <i>Isolepis platycarpa</i> | | 10 | 2006 |
| <i>Isolepis producta</i> | | 13 | 1998 |
| <i>Isolepis stellata</i> | | 1 | 2006 |
| <i>Isolepis subtilissima</i> | | 12 | 2000 |
| <i>Isolepis tasmanica</i> | | 57 | 1985 |
| <i>Isolepis wakefieldiana</i> | | 23 | 2001 |
| <i>Lepidosperma concavum</i> | | 12 | 2009 |
| <i>Lepidosperma curtisiae</i> | | 13 | 2007 |
| <i>Lepidosperma elatius</i> | | 17 | 2010 |
| <i>Lepidosperma ensiforme</i> | | 35 | 2009 |
| <i>Lepidosperma filiforme</i> | | 18 | 2060 |
| <i>Lepidosperma forsythii</i> | | 5 | 2005 |
| <i>Lepidosperma gladiatum</i> | | 6 | 2007 |
| <i>Lepidosperma globosum</i> | | 79 | 2006 |
| <i>Lepidosperma gunnii</i> | | 19 | 2008 |
| <i>Lepidosperma inops</i> | | 57 | 2009 |
| <i>Lepidosperma laterale</i> | | 5 | 2009 |
| <i>Lepidosperma longitudinale</i> | | 12 | 2007 |
| <i>Lepidosperma oldfieldii</i> | | 78 | 2055 |
| <i>Lepidosperma tortuosum</i> | | 9 | 2007 |
| <i>Lepidosperma viscidum</i> | | <1 | 1995 |
| <i>Oreobolus acutifolius</i> | | 47 | 1995 |
| <i>Oreobolus distichus</i> | | 15 | 2007 |
| <i>Oreobolus oligocephalus</i> | | 69 | 2000 |
| <i>Oreobolus oxycarpus</i> | | 45 | 1990 |
| <i>Oreobolus pumilio</i> | | 42 | 2007 |
| <i>Oreobolus tholicarpus</i> | | 43 | 2005 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------------------|-------------|---|--------------------|
| <i>Schoenoplectus pungens</i> | | 7 | 2007 |
| <i>Schoenoplectus tabernaemontani</i> | | <1 | 1999 |
| <i>Schoenus absconditus</i> | | 58 | 2004 |
| <i>Schoenus apogon</i> | | 9 | 2009 |
| <i>Schoenus biglumis</i> | | 29 | 1987 |
| <i>Schoenus brevifolius</i> | | <1 | 2005 |
| <i>Schoenus calyptratus</i> | | 21 | 2005 |
| <i>Schoenus carsei</i> | | 5 | 1998 |
| <i>Schoenus fluitans</i> | | 13 | 2002 |
| <i>Schoenus lepidosperma</i> | | 20 | 2008 |
| <i>Schoenus maschalinus</i> | | 7 | 2004 |
| <i>Schoenus nitens</i> | | 12 | 2008 |
| <i>Schoenus pygmaeus</i> | | 60 | 1986 |
| <i>Schoenus tesquorum</i> | | 11 | 2002 |
| <i>Schoenus turbinatus</i> | | 10 | 2001 |
| <i>Tetraria capillaris</i> | | 8 | 2008 |
| <i>Tricostularia pauciflora</i> | | 3 | 2005 |
| <i>Uncinia compacta</i> | | 31 | 2007 |
| <i>Uncinia divaricata</i> | | 100 | 1993 |
| <i>Uncinia elegans</i> | | 73 | 2006 |
| <i>Uncinia riparia</i> | | 32 | 1998 |
| <i>Uncinia tenella</i> | | 23 | 2004 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-------------------------|-----------------|-------------|---|--------------------|
| Juncaceae | Rushes | | (37 out of 70 species in Australia) | |
| Juncus amabilis | | | 6 | 2009 |
| Juncus antarcticus | | | 18 | 2007 |
| Juncus articulatus | Jointed Rush | | 8 | 2007 |
| Juncus astreptus | | | 56 | 2007 |
| Juncus australis | | | 15 | 2007 |
| Juncus bassianus | | | 28 | 2004 |
| Juncus bufonius | | | 4 | 2007 |
| Juncus caespiticius | | | 6 | 2006 |
| Juncus curtisiae | | | 48 | 2007 |
| Juncus falcatus | Falcate Rush | | 17 | 2006 |
| Juncus filicaulis | | | 12 | 2007 |
| Juncus fockei | | | 1 | 2006 |
| Juncus gregiflorus | | | 6 | 2007 |
| Juncus holoschoenus | Jointleaf Rush | | 5 | 2004 |
| Juncus kraussii | | | 7 | 2008 |
| Juncus pallidus | | | 10 | 2008 |
| Juncus pauciflorus | | | 11 | 2006 |
| Juncus planifolius | | | 7 | 2007 |
| Juncus prismatocarpus | | | 1 | 2000 |
| Juncus procerus | | | 15 | 2007 |
| Juncus ratkowskyanus | | | 100 | 1990 |
| Juncus revolutus | | | 12 | 2007 |
| Juncus sandwithii | | | 18 | 1992 |
| Juncus sarophorus | | | 11 | 2007 |
| Juncus scheuchzerioides | | | 50 | 2002 |
| Juncus subsecundus | | | 3 | 2007 |
| Juncus vaginatus | | | 6 | 2007 |
| Luzula acutifolia | | | 30 | 1996 |
| Luzula atrata | | | 17 | 1989 |
| Luzula australasica | | | 29 | 1998 |
| Luzula crinita | | | 100 | 2007 |
| Luzula densiflora | Dense Wood-rush | | 6 | 2007 |
| Luzula flaccida | | | 9 | 2005 |
| Luzula meridionalis | | | 5 | 2004 |
| Luzula modesta | | | 8 | 2007 |
| Luzula novae-cambriae | | | 22 | 1998 |
| Luzula poimena | | | 55 | 2004 |



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------|-------------|---|--------------------|
| Poaceae | | | |
| Grasses | | (148 out of 1057 species in Australia) | |
| Australian Bent | | 17 | 1990 |
| Agrostis australiensis | | 16 | 2002 |
| Agrostis bettyae | | 45 | 1996 |
| Agrostis diemenica | | 47 | 1996 |
| Agrostis joyceae | | 100 | 2007 |
| Agrostis magellanica | | 10 | 1981 |
| Agrostis muelleriana | | 43 | 2005 |
| Agrostis parviflora | | 29 | 2007 |
| Agrostis propinqua | | 30 | 1996 |
| Agrostis thompsoniae | | 18 | 2001 |
| Agrostis venusta | | 5 | 2002 |
| Amphibromus archeri | Vulnerable | 1 | 1980 |
| Amphibromus fluitans | | <1 | |
| Amphibromus macrorhinus | | 3 | 2004 |
| Amphibromus neesii | | 16 | 1996 |
| Amphibromus recurvatus | | 14 | 2002 |
| Amphibromus sinuatus | | 9 | 2009 |
| Anthoxanthum odoratum | | <1 | 1988 |
| Aristida benthamii | | <1 | 1988 |
| Aristida jerichoensis | | 30 | 2007 |
| Australopyrum pectinatum | | 6 | 1994 |
| Australopyrum velutinum | | 4 | 2007 |
| Austrodanthonia caespitosa | | 8 | 2004 |
| Austrodanthonia carphoides | | 46 | 1996 |
| Austrodanthonia diemenica | | 2 | 2007 |
| Austrodanthonia geniculata | | 12 | 2009 |
| Austrodanthonia induta | | 13 | 2005 |
| Austrodanthonia laevis | | 15 | 2007 |
| Austrodanthonia penicillata | | 10 | 2007 |
| Austrodanthonia pilosa | | 76 | 2009 |
| Austrodanthonia popinensis | Endangered | 3 | 2007 |
| Austrodanthonia racemosa | | 3 | 2007 |
| Austrodanthonia setacea | | 6 | 2007 |
| Austrodanthonia tenuior | | 10 | 2004 |
| Austrofestuca hookeriana | | 29 | 1999 |
| Austrofestuca littoralis | | 72 | 2008 |
| Austrostipa aphylla | | 3 | 2009 |
| Austrostipa bigeniculata | | 1 | 2005 |
| Austrostipa blackii | | 8 | 2007 |
| Austrostipa flavescens | | 6 | 2006 |
| Austrostipa mollis | | <1 | 1986 |
| Austrostipa nitida | | 4 | 2009 |
| Austrostipa nodosa | | 22 | 2007 |
| Austrostipa pubinodis | | 6 | 2007 |
| Austrostipa rudis | | <1 | 2009 |
| Austrostipa scabra | | | |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|---------------------------|----------------------|---|--------------------|
| Austrostipa semibarbata | | 11 | 2007 |
| Austrostipa stipoides | | 19 | 2007 |
| Austrostipa stuposa | | 32 | 2007 |
| Bothriochloa macra | | <1 | 1994 |
| Bromus arenarius | | <1 | 1966 |
| Cynodon dactylon | | <1 | 2008 |
| Deschampsia cespitosa | | 27 | 1993 |
| Deschampsia gracillima | | 100 | 1989 |
| Deyeuxia apsleyensis | | 86 | 2003 |
| Deyeuxia brachyathera | | 2 | 1994 |
| Deyeuxia carinata | | 13 | 1997 |
| Deyeuxia contracta | | 33 | 2007 |
| Deyeuxia decipiens | | 8 | 2001 |
| Deyeuxia densa | | 12 | 2005 |
| Deyeuxia frigida | | 12 | 1996 |
| Deyeuxia gunniana | | 6 | 1996 |
| Deyeuxia innominata | | 33 | 1996 |
| Deyeuxia minor | | 15 | 2007 |
| Deyeuxia monticola | | 27 | 2006 |
| Deyeuxia quadriseta | | 13 | 2007 |
| Deyeuxia rodwayi | | 9 | 2002 |
| Deyeuxia scaberula | | 8 | 2003 |
| Dichanthium sericeum | Silky Bluestem | <1 | 1982 |
| Dichanthium setosum | Vulnerable | 1 | 1988 |
| Dichelachne crinita | | 6 | 2007 |
| Dichelachne inaequiglumis | | 15 | 2007 |
| Dichelachne micrantha | | 2 | 2004 |
| Dichelachne rara | | 16 | 2009 |
| Dichelachne sieberiana | | 2 | 1988 |
| Distichlis distichophylla | | 8 | 2007 |
| Dryopoa dives | Giant Mountain Grass | 4 | 2009 |
| Echinochloa esculenta | | 5 | 2007 |
| Echinopogon ovatus | | 4 | 2007 |
| Ehrharta erecta | | 1 | 2007 |
| Elymus multiflorus | | 7 | 1986 |
| Elymus scaber | | 1 | 2001 |
| Eragrostis brownii | Brown's Lovegrass | <1 | 2006 |
| Festuca asperula | | <1 | 1973 |
| Festuca contracta | | 100 | 2007 |
| Festuca plebeia | | 81 | 2006 |
| Glyceria australis | Manna Grass | 8 | 2004 |
| Hemarthria uncinata | | 5 | 2007 |
| Hierochloa fraseri | | 52 | 2002 |
| Hierochloa rariflora | | 2 | 2003 |
| Hierochloa redolens | | 29 | 2007 |
| Hierochloa submutica | | 6 | 1967 |
| Imperata cylindrica | | <1 | 2006 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------------|-------------------------------|---|--------------------|
| Isachne globosa | | <1 | 2002 |
| Lachnagrostis billardierei | | 15 | 2000 |
| Lachnagrostis collicola | | 25 | 1985 |
| Lachnagrostis filiformis | | 4 | 2007 |
| Lachnagrostis lacunarum | | 31 | 1987 |
| Lachnagrostis meionectes | | 3 | 1983 |
| Lachnagrostis morrisii | | 55 | 1996 |
| Lachnagrostis punicea | | 4 | 2007 |
| Lachnagrostis robusta | | 13 | 2007 |
| Lachnagrostis scabra | | 14 | 1992 |
| Microlaena stipoides | | 6 | 2009 |
| Microlaena tasmanica | | 39 | 2001 |
| Notodanthonia gracilis | | 31 | 2007 |
| Notodanthonia semiannularis | | 8 | 2007 |
| Paspalum distichum | | <1 | 1989 |
| Paspalum scrobiculatum | Scrobic | <1 | 2000 |
| Pentapogon quadrifidus | | 19 | 2005 |
| Phragmites australis | | 2 | 2007 |
| Poa clelandii | | 8 | 2005 |
| Poa clivicola | | 21 | 2004 |
| Poa cookii | | 50 | 2002 |
| Poa costiniana | | 5 | 1989 |
| Poa fawcettiae | | <1 | 1985 |
| Poa foliosa | | 100 | 2007 |
| Poa gunnii | | 49 | 2007 |
| Poa halmaturina | | 3 | 1992 |
| Poa hiemata | | 5 | 1990 |
| Poa hookeri | | 39 | 2007 |
| Poa jugicola | | 25 | 1994 |
| Poa labillardierei | | 7 | 2009 |
| Poa litorosa | | 100 | 1993 |
| Poa mollis | | 56 | 2006 |
| Poa poiformis | | 12 | 2008 |
| Poa rodwayi | | 32 | 2009 |
| Poa sieberiana | | 5 | 2009 |
| Poa tenera | | 9 | 2008 |
| Puccinellia macquariensis | Macquarie Island Alkali Grass | 100 | 1998 |
| Puccinellia stricta | | 5 | 2006 |
| Rytidosperma australe | | 35 | 1996 |
| Rytidosperma dimidiatum | | 61 | 2007 |
| Rytidosperma fortunae-hibernae | | 57 | 2005 |
| Rytidosperma nitens | | 47 | 2005 |
| Rytidosperma nivicola | | 28 | 1996 |
| Rytidosperma nudiflorum | | 16 | 1996 |
| Rytidosperma pauciflorum | | 56 | 2002 |
| Saxipoa saxicola | | 25 | 2000 |
| Spinifex hirsutus | | <1 | 1983 |

Species are shown only if they belong to the families covered in ANHAT as shown in Appendix 1 and ANHAT holds records of the species from this region.



| Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|-----------------------------------|----------------------------|---|--------------------|
| <i>Spinifex sericeus</i> | | 3 | 2001 |
| <i>Sporobolus virginicus</i> | | <1 | 1982 |
| <i>Tetrarrhena acuminata</i> | | 21 | 2008 |
| <i>Tetrarrhena distichophylla</i> | | 29 | 2009 |
| <i>Tetrarrhena juncea</i> | | <1 | 1987 |
| <i>Tetrarrhena oreophila</i> | | 83 | 2002 |
| <i>Themeda triandra</i> | | 2 | 2009 |
| <i>Trisetum spicatum</i> | Bristle Grass | 14 | 1988 |
| <i>Zoysia macrantha</i> | | 4 | 2007 |
| Restionaceae | Restiona Sedges | (14 out of 146 species in Australia) | |
| <i>Apodasmia brownii</i> | | 13 | 2007 |
| <i>Baloskion australe</i> | | 32 | 2006 |
| <i>Baloskion tetraphyllum</i> | | 8 | 2004 |
| <i>Calorophus elongatus</i> | | 41 | 2004 |
| <i>Calorophus erostris</i> | | 50 | 2008 |
| <i>Chordifex hookeri</i> | | 49 | 2008 |
| <i>Chordifex monocephalus</i> | | 49 | 2006 |
| <i>Empodisma minus</i> | | 17 | 2099 |
| <i>Eurychorda complanata</i> | | 24 | 2008 |
| <i>Hypolaena fastigiata</i> | | 5 | 2008 |
| <i>Leptocarpus tenax</i> | | 15 | 2008 |
| <i>Lepyrodia muelleri</i> | | 8 | 2007 |
| <i>Sporodanthus tasmanicus</i> | | 34 | 2008 |
| <i>Winifredia sola</i> | | 53 | 2008 |
| Typhaceae | Cumbungi and Allies | (2 out of 2 species in Australia) | |
| <i>Typha domingensis</i> | Bulrush, Cumbungi | 1 | 2007 |
| <i>Typha orientalis</i> | | 1 | 2007 |
| Xyridaceae | Xyris Herbs | (5 out of 26 species in Australia) | |
| <i>Xyris gracilis</i> | | 6 | 2007 |
| <i>Xyris marginata</i> | | 45 | 2008 |
| <i>Xyris muelleri</i> | | 42 | 2008 |
| <i>Xyris operculata</i> | | 13 | 2008 |
| <i>Xyris tasmanica</i> | | 28 | 1999 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|--------------------------|--|--|---|--------------------|
| Proteales | | | | |
| Proteaceae | | | | |
| | Banksias, Grevilleas and Allies | (39 out of 1147 species in Australia) | | |
| Agastachys odorata | White waratah | | 52 | 2007 |
| Banksia marginata | Silver Banksia | | 13 | 2009 |
| Banksia serrata | Saw Banksia | | <1 | 2002 |
| Bellenden montana | Mountain rocket | | 49 | 2007 |
| Cenarrhenes nitida | Native Plum | | 38 | 2008 |
| Conospermum hookeri | Smoke-bush | | 30 | 2008 |
| Conospermum taxifolium | Smoke-bush | | 2 | 1990 |
| Grevillea australis | Alpine Grevillea | | 25 | 2004 |
| Grevillea linearifolia | Linear-leaf Grevillea | | <1 | |
| Grevillea oleoides | Grevillea | | 3 | |
| Grevillea repens | Creeping Grevillea | | 6 | |
| Grevillea robusta | Silky Oak | | <1 | 2000 |
| Grevillea rosmarinifolia | Grevillea | | 2 | 2007 |
| Grevillea sphacelata | Grevillea | | 2 | |
| Hakea dactyloides | Finger Hakea | | <1 | |
| Hakea decurrens | Hakea | | <1 | 2005 |
| Hakea epiglottis | Hakea | | 57 | 2007 |
| Hakea lissosperma | Needle Bush | | 48 | 2009 |
| Hakea megadenia | Hakea | | 75 | 2006 |
| Hakea microcarpa | Small Fruit Hakea | | 18 | 2009 |
| Hakea nodosa | Yellow Hakea | | 4 | 2008 |
| Hakea rostrata | Beaked Hakea | | <1 | 1982 |
| Hakea rugosa | Dwarf Hakea | | 1 | 1986 |
| Hakea sericea | Hakea | | <1 | 2005 |
| Hakea teretifolia | Dagger Hakea | | 12 | 2002 |
| Isopogon anethifolius | Isopogon | | 1 | 2004 |
| Lomatia fraseri | Lomatia | | <1 | 2002 |
| Lomatia polymorpha | Lomatia | | 56 | 2008 |
| Lomatia tinctoria | Guitar Plant | | 44 | 2009 |
| Orites acicularis | Yellow Bush | | 64 | 2007 |
| Orites diversifolius | Orites | | 71 | 2005 |
| Orites milliganii | Orites | | 57 | 1993 |
| Orites revolutus | Orites | | 59 | 2007 |
| Persoonia gunnii | Persoonia | | 59 | 2007 |
| Persoonia juniperina | Prickly Geebung | | 16 | 2009 |
| Persoonia moscalii | Persoonia | | 100 | 2005 |
| Persoonia muelleri | Persoonia | | 38 | 2004 |
| Telopea speciosissima | Waratah | | 4 | 2003 |
| Telopea truncata | Tasmanian Waratah | | 43 | 2008 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------|--|-----------------------|---|--------------------|
| Sapindales | | | | |
| Meliaceae | | | | |
| | Rosewoods and Mahoganies | | (1 out of 44 species in Australia) | |
| Melia azedarach | White Cedar | | <1 | 2001 |
| Rutaceae | | | | |
| | Boronia, Correa, Citrus, Phebalium, Philotheca, Zieria and Allies | | (28 out of 496 species in Australia) | |
| Acradenia euodiiformis | Yellow Satinheart | | 2 | |
| Acradenia frankliniae | Wirewood | | 10 | 2007 |
| Boronia anemonifolia | Boronia | | 7 | 2008 |
| Boronia citriodora | Lemon Boronia | | 50 | 2005 |
| Boronia elisabethiae | Boronia | | 44 | 2004 |
| Boronia gunnii | Gunn's Boronia | Vulnerable | 33 | 2004 |
| Boronia hippopala | Velvet Boronia | Vulnerable | 33 | 2006 |
| Boronia nana | Small Boronia | | 7 | 1996 |
| Boronia parviflora | Small Boronia | | 17 | 2008 |
| Boronia pilosa | Hairy Boronia | | 33 | 2008 |
| Boronia rhomboidea | Rhomboid Boronia | | 15 | 2006 |
| Boronia rozefeldsii | Boronia | | 100 | 2002 |
| Correa alba | White Correa | | 16 | 2007 |
| Correa backhouseana | Correa | | 6 | 1999 |
| Correa lawrenceana | Correa | | 13 | 2008 |
| Correa reflexa | Common Correa | | 6 | 2008 |
| Leionema bilobum | Notched Phebalium | | 8 | 2002 |
| Leionema montanum | Leionema | | 50 | 2001 |
| Leionema oldfieldii | Leionema | | 50 | 1996 |
| Nematolepis squamea | Nematolepis | | 28 | 2009 |
| Philotheca freyciana | Freycinet Waxflower | Critically endangered | 100 | 2003 |
| Philotheca myoporoides | Philotheca | | <1 | 1988 |
| Philotheca verrucosa | Philotheca | | 26 | 2008 |
| Philotheca virgata | Tasmanian Waxflower | | 24 | 2004 |
| Zieria arborescens | Stinkwood | | 23 | 2009 |
| Zieria cytisoides | Downy Zieria | | <1 | 1998 |
| Zieria floydii | Zieria | Endangered | 17 | 1992 |
| Zieria littoralis | Zieria | | 25 | 2007 |
| Sapindaceae | | | | |
| | Tamarind, Whitewood, Tuckeroo, Hop Bushes | | (2 out of 222 species in Australia) | |
| Dodonaea filiformis | Hop Bush | | 61 | 2006 |
| Dodonaea viscosa | Sticky Hop Bush | | 2 | 2009 |



| | Common Name | EPBC Status | Proportion of sampled range in the region (%) | Most recent record |
|------------------------|--|--|---|--------------------|
| Solanales | | | | |
| Convolvulaceae | | | | |
| | Bindweed, Morning Glory | (7 out of 132 species in Australia) | | |
| Calystegia soldanella | Bindweed | | 5 | 2009 |
| Convolvulus erubescens | Australian Bindweed | | 4 | 2004 |
| Cuscuta australis | Australian Dodder | | 2 | 1973 |
| Cuscuta tasmanica | Tasmanian Dodder | | 9 | 2006 |
| Dichondra repens | Kidney Weed | | 4 | 2009 |
| Wilsonia backhousei | Narrow-leaf Wilsonia | | 9 | 2007 |
| Wilsonia rotundifolia | Round-leaf Wilsonia | | 4 | 2008 |
| Solanaceae | | | | |
| | Native Gooseberries, Wild Tomatos and Boxthorns | (5 out of 210 species in Australia) | | |
| Cyphanthera tasmanica | Ray Flower | | 79 | 2006 |
| Solanum americanum | Glossy Nightshade | | <1 | 2006 |
| Solanum laciniatum | Kangaroo Apple | | 7 | 2007 |
| Solanum opacum | Green-berry Nightshade | | <1 | |
| Solanum vescum | Kangaroo Apple | | 9 | 2002 |



Appendix 1

List of all families covered by the Australian Natural Heritage Assessment Tool

These families of plants and animals are included in NRM region summaries where they occur in the region.

The groups covered across Australia include:

- all vertebrates
 - i.e. mammals, birds, reptiles, frogs and fish (4,200 species)
- 137 families of vascular plants (> 15,000 species) comprising the majority of Australia's named plant species
- A range of invertebrate groups including:
 - numerous insect families (> 16,600 species)
 - all land snails (> 2,300 species)
 - selected spider families (> 1,100 species)
 - rotifers – aquatic wheel animals (> 700 species)

Vertebrate fauna

Actinopterygii

Ambassidae
Apogonidae
Ariidae
Atherinidae
Batrachoididae
Belonidae
Blenniidae
Brachionichthyidae
Carangidae
Cepolidae
Chanidae
Clinidae
Clupeidae
Cynoglossidae
Diodontidae
Eleotridae
Elopidae
Engraulidae
Enoplosidae
Ephippidae
Galaxiidae
Gerreidae
Gobiidae
Gonorynchidae
Hemiramphidae

Ray-finned fish

Chanda Perches, Glassfishes, Perchlets
Big Eyes, Cardinal Fishes, Cardinalfishes, Gobbleguts, Soldier
Fork-tailed Catfishes
Hardyheads, Silversides, Whitebait
Bastard Stonefish, Frogfishes, Toadfishes
Long Toms, Longtoms, Needle Fishes, Needlefishes
Blennies, Fangblennies, Rockshippers
Handfishes
Amberjacks, Jacks, Pilotfishes, Pompanos, Queenfishes, Run
Bandfishes
Milkfish
Snake Blennies, Weedfishes
Herrings, Menhadens, Pilchards, Sardines, Shads, Sprats
Tongue Soles
Burrfishes, Porcupinefishes
Gudgeons, Sleepers
Ladyfish
Anchovies
Oldwife
Batfishes, Spadefishes
Galaxias, Salamanderfish
Mojarras, Silver Biddies, Silver-bellies, Silverbellies, Silverbiddi
Gobies
Beaked Salmon, Beaked Sandfish, Ratfish
Garfishes, Halfbeaks



Vertebrate fauna

| | |
|------------------|---|
| Kuhliidae | Flagtails |
| Kurtidae | Nurseryfishes |
| Kyphosidae | Drummers, Halfmoons, Knifefishes, Microcanthids, Nibblers, S |
| Labridae | Wrasses, Parrotfishes, Rockwhittings |
| Leiognathidae | Pony Fishes, Ponyfishes, Slipmouths |
| Lutjanidae | Fusilers, Hussars, Snappers, Tropical Snappers |
| Megalopidae | Tarpons |
| Melanotaeniidae | Rainbowfishes |
| Microdesmidae | Dartfishes, Wormfishes |
| Monacanthidae | Leatherjackets |
| Mugilidae | Mulletts |
| Muraenidae | Moray Eels |
| Ophichthidae | Snake Eels, Worm Eels |
| Osteoglossidae | Saratoga |
| Ostraciidae | Boxfishes, Coffe Fishes, Cowfishes, Trunkfishes, Turretfishes |
| Paralichthyidae | Sand Flounders |
| Percichthyidae | Australian Freshwater Basses, Australian Freshwater Blackfish |
| Platycephalidae | Flatheads |
| Pleuronectidae | Right Handed Flatfishes, Righteye Flounders |
| Plotosidae | Eel-Tailed Catfishes |
| Polynemidae | Tassel-fishes, Threadfin Salmons, Threadfins |
| Pomacentridae | Damselfishes |
| Pomatomidae | Bluefish, Tailor |
| Pristigasteridae | Ilishas, Pellonas, Pristigasterids |
| Pseudomugilidae | Blue Eyes, Blue-eyes |
| Retropinnidae | Southern Graylings, Southern Smelts |
| Scatophagidae | Butterfishes, Scats |
| Sciaenidae | Croakers, Drums, Jewfishes |
| Serranidae | Basslets, Perchlets, Rockcods, Wirrahs |
| Siganidae | Mi Mi, Rabbit Fish, Rabbitfishes, Spinefoot |
| Sillaginidae | Sillagos, Whittings |
| Soleidae | Soles |
| Sparidae | Breams, Porgies, Seabreams |
| Synbranchidae | One-gilled Eel, Swamp Eels |
| Syngnathidae | Pipefishes, Seahorses |
| Synodontidae | Lizardfishes |
| Terapontidae | Grunters |
| Tetraodontidae | Blowfishes, Pufferfishes, Puffers, Toadfishes, Tobys |
| Tetrarogidae | Fortesques, Sailback Scorpionfishes, Waspfish |
| Toxotidae | Archerfishes, Rifle Fish |
| Zanclidae | Moorish Idol |

Amphibia

| | |
|----------------|--------------------------------|
| Hylidae | Tree-Frogs |
| Microhylidae | Micro Tree-Frogs |
| Myobatrachidae | Myobatrachid or Southern Frogs |
| Ranidae | True Frogs |

Aves

| | |
|--------------|-----------------------------------|
| Acanthizidae | Scrubwrens, Thornbills and Allies |
| Accipitridae | Eagles, Hawks and Kites |

Amphibians

| |
|--------------------------------|
| Tree-Frogs |
| Micro Tree-Frogs |
| Myobatrachid or Southern Frogs |
| True Frogs |

Birds

| |
|-----------------------------------|
| Scrubwrens, Thornbills and Allies |
| Eagles, Hawks and Kites |



Vertebrate fauna

| | |
|------------------|--|
| Aegothelidae | Owlet Nightjars |
| Alaudidae | Larks |
| Alcedinidae | Kingfishers and Kookaburras |
| Anatidae | Swans, Ducks and Geese |
| Anhingidae | Darter |
| Anseranatidae | Magpie Goose |
| Apodidae | Swifts and Swiftlets |
| Ardeidae | Herons, Egrets and Bitterns |
| Artamidae | Butcherbirds, Currawongs and Woodswallows |
| Atrichornithidae | Scrub-birds |
| Burhinidae | Thicknees (Stone-Curlews) |
| Cacatuidae | Cockatoos |
| Campephagidae | Cuckoo-shrikes, Trillers and Minivets |
| Caprimulgidae | Nightjars |
| Casuariidae | Cassowary and Emus |
| Centropodidae | Pheasant Coucal |
| Charadriidae | Plovers, Dotterels and Lapwings |
| Ciconiidae | Black-necked Stork |
| Climacteridae | Treecreepers |
| Columbidae | Pigeons and Doves |
| Coraciidae | Dollarbird |
| Corcoracidae | Australasian Mud-nesters |
| Corvidae | Crows and Jays |
| Cuculidae | Cuckoos |
| Dicaeidae | Flower-peckers |
| Dicruridae | Fan-tails, Drongos and Monarchs |
| Diomedidae | Albatrosses |
| Estrildidae | Grass Finches |
| Eupetidae | Whipbirds, Quail-thrushes and Jewel-babblers |
| Falconidae | Falcons and Kestrels |
| Fregatidae | Frigatebirds |
| Glareolidae | Pratincoles |
| Gruidae | Cranes |
| Haematopodidae | Oystercatchers |
| Hirundinidae | Swallows and Martins |
| Hydrobatidae | Storm-petrels |
| Jacaniidae | Jacanas |
| Laridae | Gulls, Terns, Skuas and Jaegers |
| Maluridae | Fairy-wrens and Grasswrens |
| Megapodiidae | Mound-builders |
| Meliphagidae | Honeyeaters and Australian Chats |
| Menuridae | Lyrebirds |
| Meropidae | Rainbow Bee-eater |
| Motacillidae | Pipits and Wagtails |
| Muscicapidae | Old World Flycatchers and Thrushes |
| Nectariniidae | Sunbirds and Spiderhunters |
| Neosittidae | Australian Sitella |
| Oriolidae | Old World Orioles and Figbirds |
| Orthonychidae | Logrunners |
| Otididae | Australian Bustard |
| Pachycephalidae | Whistlers, Shrike-thrushes and Allies |



Vertebrate fauna

| | |
|-------------------|--|
| Paradisaeidae | Birds-of-Paradise |
| Pardalotidae | Pardalotes |
| Pedionomidae | Plains-wanderer |
| Pelecanidae | Pelicans |
| Petroicidae | Australasian Robins |
| Phaethonitidae | Tropicbirds |
| Phalacrocoracidae | Cormorants and Shags |
| Phasianidae | Quail, Fowl and Allies |
| Pittidae | Pittas |
| Podargidae | Frogmouths |
| Podicipedidae | Grebes |
| Pomatostomidae | Australasian Babblers |
| Procellariidae | Fulmars, Petrels, Prions and Shearwaters |
| Psittacidae | Parrots |
| Ptilonorhynchidae | Bowerbirds |
| Rallidae | Crakes, Moorhens, Rails and Allies |
| Recurvirostridae | Stilts and Avocet |
| Rostratulidae | Painted Snipe |
| Scolopacidae | Waders (Snipe, Godwits, Curlew, Sandpipers and Allies) |
| Spheniscidae | Penguins |
| Strigidae | Hawk-type Owls |
| Sturnidae | Starlings and Mynas |
| Sulidae | Boobies and Gannets |
| Sylviidae | Old World Warblers |
| Threskiornithidae | Ibis and Spoonbills |
| Turcidae | Button-quail |
| Tytonidae | Barn-type Owls |
| Zosteropidae | White-eyes (Silvereyes) |

Chondrichthyes

| | |
|------------|--|
| Alopiidae | Thresher Sharks |
| Dasyatidae | Stingrays |
| Lamnidae | Mackerel, White Pointer, Mako and Porbeagle Sharks |
| Pristidae | Sawfishes |

Mammalia

| | |
|-------------------|---|
| Acrobatidae | Feathertailed Gliders |
| Burramyidae | Pygmy-possums |
| Dasyuridae | Dasyurids (Quolls, Antechinus, Dunnarts and Allies) |
| Delphinidae | Dolphins, Orcas, Pilot Whales and Melon-headed Whales |
| Emballonuridae | Sheath-tailed Bats |
| Hipposideridae | Leaf-nosed Bats |
| Kogiidae | Dwarf and Pygmy Sperm Whales |
| Macropodidae | Wallabies, Kangaroos and Tree-kangaroos |
| Megadermatidae | False Vampire (Ghost) Bats |
| Molossidae | Free-tailed Bats |
| Muridae | Rats and Mice |
| Myrmecobiidae | Numbat |
| Notoryctidae | Marsupial Mole |
| Ornithorhynchidae | Platypus |
| Otariidae | Eared Seals |

Sharks and rays

Mammals



Vertebrate fauna

| | |
|------------------|---|
| Peramelidae | Bandicoots and Spiny Bandicoot |
| Petauridae | Wrist-winged Gliders, Striped Possum and Leadbeaters Possum |
| Phalangeridae | Brush-tail Possums, Cuscuses and Scaly-tailed Possum |
| Phascolarctidae | |
| Phocidae | True Seals |
| Phocoenidae | Porpoises |
| Physeteridae | Sperm Whales |
| Potoroidae | Potoroos, Bettongs and the Musky Rat Kangaroo |
| Pseudocheiridae | Ring-tailed Possums and Greater Glider |
| Pteropodidae | Fruit-bats and Allies |
| Rhinolophidae | Horseshoe Bats |
| Tachyglossidae | Echidna |
| Tarsipedidae | Honey Possum |
| Thylacinidae | Thylacine |
| Thylacomyidae | Bilbies |
| Vespertilionidae | Vespertilionid Bats |
| Vombatidae | |
| Ziphiidae | Beaked Whales |

Petromyzontida

| | |
|-----------------|----------|
| Petromyzontidae | Lampreys |
|-----------------|----------|

Reptilia

| | |
|-------------------|------------------------------|
| Acrochordidae | File snakes |
| Agamidae | Dragon lizards |
| Boidae | Pythons |
| Carettochelydidae | Pitted-shelled turtle |
| Chelidae | Side-necked Tortoises |
| Colubridae | Rear fang snakes |
| Crocodylidae | Crocodyles |
| Elapidae | Front fang snakes |
| Gekkonidae | Geckos |
| Pygopodidae | Legless lizards |
| Scincidae | Skinks |
| Typhlopidae | Worm-snakes and Blind-Snakes |
| Varanidae | Monitors |

Reptiles

Sarcopterygii

| | |
|----------------|------------------|
| Ceratodontidae | Lobe-finned fish |
|----------------|------------------|

Lungfishes



Invertebrate fauna

Arachnida

| | |
|------------------|---|
| Actinopodidae | Two-doored Trapdoor Spiders |
| Araneidae | Orb-weaving Spiders |
| Barychelidae | Brush-footed Trapdoor Spiders |
| Chthoniidae | |
| Ctenizidae | Trapdoor Spiders |
| Cyrtachenidiidae | Trapdoor Spiders |
| Dipluridae | Funnel-web-like Spiders |
| Garypidae | |
| Hexathelidae | Funnel-web Spiders |
| Idiopidae | Trapdoor Spiders |
| Lycosidae | Wolf Spiders |
| Migidae | Brushless-legged Trapdoor Spiders |
| Nemesiidae | Funnel-web Spiders |
| Sparassidae | Huntsmen Spiders (formerly Heteropodidae) |
| Theraphosidae | Bird-eating Trapdoor Spiders |

Bivalvia

| | |
|-------------|--------------------|
| Hyriidae | Freshwater Mussels |
| Sphaeriidae | Pea-shells |

Chilopoda

Scolopendridae

Bivalves

Centipedes

Diplopoda

Dalodesmidae
Metopidiotrichidae
Pachybolidae
Paradoxosomatidae

Millipedes

Eurotatoria

Adinetidae
Asplanchnidae
Atrochidae
Brachionidae
Collothecidae
Conochilidae
Dicranophoridae
Epiphanidae
Euchlanidae
Flosculariidae
Gastropodidae
Habrotrichidae
Hexarthridae
Ituridae
Lecanidae
Lepadellidae
Lindiidae
Microcodonidae
Mytilinidae
Notommatidae

Wheel Animals

Rotifers with Undifferentiated Cilia
Sac-like Wheel Animals
Sessile Wheel Animals
Sac-like Wheel Animal
Sessile Wheel Animals
Six-"armed" Wheel Animals
Tong-grabbing Wheel Animals
Shell-less Wheel Animals
Wheel Animals
Sessile-footed Wheel Animals
Shelled Wheel Animals
Rotifers with Paired Trochi
Gelatinous-colonial Wheel Animals
Shelled Wheel Animals
Shelled Wheel Animal
Wheel Animals
Wheel Animals
Wheel Animals
Wheel Animals
Pseudo-segmented Wheel Animals



Invertebrate fauna

| | |
|------------------|---|
| Philodinavidae | Rotifers with Simplified Trochi |
| Philodinidae | Rotifers with Paired Trochi |
| Proalidae | Worm-like Wheel Animals |
| Scaridiidae | Wheel Animals |
| Synchaetidae | Sac-, Bell- or Cone-shaped Wheel Animals |
| Testudinellidae | Free-swimming Round-shelled Wheel Animals |
| Tetrasiphonidae | Wheel Animal |
| Trichocercidae | Comma-like Wheel Animals |
| Trichotriidae | Wheel Animals |
| Trochosphaeridae | Globular Wheel Animals |

Gastropoda

Achatinellidae
Amphibolidae
Ancylidae
Assimineidae
Athoracophoridae
Bithyniidae
Bulimulidae
Camaenidae
Caryodidae
Cerastuidae
Charopidae
Corillidae
Cyclophoridae
Cystopeltidae
Diplommatinidae
Ellobiidae
Glacidorbidae
Helicarionidae
Helicinidae
Helicodiscidae
Hydrobiidae
Hydrocenidae
Hydrococcidae
Iravadiidae
Lymnaeidae
Megaspiridae
Neritidae
Planorbidae
Pomatiopsidae
Punctidae
Pupillidae
Pupinidae
Rathouisiidae
Rhytididae
Subulinidae
Succineidae
Thiaridae
Trochomorphidae
Truncatellidae

Snails

Conical-shelled Land Snails
Supra-littoral Globose "Land" Snails
Freshwater Limpets
Freshwater, Land and Saltmarsh Snails
Triangular Mantled Slugs
Operculate Freshwater Snails
Elongate High-spined Land Snails
Camaenid Land Snails
Giant Land Snails
Land Snails
Flattened and Turbinate Land Snails
Land Snail
Horny Circular Operculate Land Snails
Australian "Half" Slugs
Circular Operculate Land Snails
Primitive Supralittoral Pulmonate Snails
Freshwater Snails
Helicarionid Land Snails
Top-shaped Operculate Land Snails
Stenopylis Land Snail
Freshwater Operculate Snails
Calcareous Operculate Land Snails
Paucispiral Operculate Saltmarsh Snails
Estuarine and Marine Operculate Snails
Eutrophic Freshwater Snails
Coelocion Land Snail
Calcareous Operculate Marine and Estuarine Snails
Freshwater Snails
Salt-lake Snails
Translucent Frosted Land Snails
Pupa-shaped Shelled Land Snails
Glossy and Porcellaneous Operculate Land Snails
Carnivorous Primitive Slugs
Carnivorous Land Snails
Elongate Land Snails
Amber Land Snails
Freshwater Horny Operculate Snails
Native Slug
Supralittoral Calcareous Operculate Snails



Invertebrate fauna

Viviparidae Freshwater Subspirial Horny Operculate Snails

Insecta

Acanthosomatidae

Achilidae

Aenictopecheidae

Aeshnidae

Aleyrodidae

Anthocoridae

Aphididae

Aphylidae

Apidae

Apioceridae

Aradidae

Archipetalidae

Artheneidae

Asilidae

Atriplectididae

Austrocorduliidae

Austropetalidae

Baetidae

Belostomatidae

Berytidae

Blissidae

Bombyliidae

Buprestidae

Calocidae

Carabidae

Ceratocombidae

Cercopidae

Chlorocyphidae

Chorismagrionidae

Cicadidae

Cixiidae

Coccidae

Coenagrionidae

Colletidae

Colobathristidae

Conoesucidae

Cordulephyidae

Corduliidae

Coreidae

Corixidae

Cryptorhamphidae

Cydnidae

Cymidae

Delphacidae

Diaspididae

Dinidoridae

Diphlebiidae

Dipsocoridae

Insects

Shield Bugs

Aenictopecheidae

Hawker, Duskhawker and Emperor Damselflies

Whiteflies

Aphids

Social Bees

Flower-loving Flies

Flat bugs, Bark bugs

Tasmanian Redspot Dragonfly

Robber Flies

Caddisflies

Hawk, Mosquitohawk and Swiftwing Dragonflies

Redspot Dragonflies

Blue-winged Olive Mayflies, Small Mayflies

Stilt bugs

Chinch Bugs

Bee Flies

Jewel Beetles

Caddisflies

Ground Beetles

Jewel Damselflies

Pretty Relict Damselfly

Bluetail and Thin Damselflies

Short-tongued Bees

Caddisflies

Shutwing Dragonflies

Tigerhawk and Emerald Dragonflies

Waterboatmen

Burrower Bugs

Armoured Scales, Diaspids

Rockmaster Damselflies



Invertebrate fauna

| | |
|-------------------|---|
| Dytiscidae | Dytiscidine Water Beetles |
| Ecnomidae | Caseless Caddisflies |
| Enicocephalidae | |
| Eriococcidae | |
| Eurybrachyidae | |
| Flatidae | |
| Formicidae | Ants |
| Gelastocoridae | Toad Bugs |
| Geocoridae | |
| Gerridae | Water-striders |
| Glossosomatidae | Stone-case Caddisflies |
| Gomphidae | Dragon, Hunter and Vicetail Dragonflies |
| Gomphomacromiidae | Urly Dragonflies |
| Gyrinidae | Gyrinid Water Beetles |
| Halictidae | |
| Haliplidae | Haliplid Water Beetles |
| Helicophidae | Caddisflies |
| Helicopsychidae | Snail Case Caddisflies |
| Hemicorduliidae | Emerald and Swamp Emerald Dragonflies |
| Hemiphlebiidae | Ancient Greenling |
| Hesperiidae | Skippers, flats, awls & darts |
| Heterogastridae | |
| Hydrobiosidae | Caddisflies |
| Hydrometridae | Marsh Treaders, Water Measurers |
| Hydrophilidae | Water Beetle |
| Hydropsychidae | Net Spinning Caddisflies |
| Hydroptilidae | Micro Caddisflies |
| Hygrobiidae | Hygrobiid Water Beetles |
| Hyocephalidae | |
| Idiosolididae | |
| Isostictidae | Wiretail and Pin Damselflies |
| Kalotermitidae | Termite |
| Kokiriidae | Caddisflies |
| Largidae | |
| Leptoceridae | Long-horned Caddisflies |
| Leptophlebiidae | Prong Gilled Mayflies |
| Leptopodidae | Leptopodidae |
| Lestidae | Ringtail, Reedling and Dusky Spreadwing Damselflies |
| Lestoididae | Bluestreak Damselflies |
| Lestoniidae | |
| Libellulidae | Perchers, Archtails and other Dragonflies |
| Lindeniidae | Tiger Dragonflies |
| Lycaenidae | Blues, Coppers, Hairstreaks and Metalmarks |
| Lygaeidae | |
| Macromiidae | Cruiser Dragonflies |
| Mastotermitidae | Giant Northern Termite |
| Megachilidae | |
| Megapodagrionidae | Flatwing Damselflies |
| Membracidae | Horned Treehoppers |
| Mesoveliidae | |
| Miridae | Miridae |



Invertebrate fauna

| | |
|-------------------|--|
| Mydidae | Mydas Flies, Wasp-mimic Flies |
| Nabidae | |
| Naucoridae | |
| Nepidae | Water Scorpions |
| Nesameletidae | Mayflies |
| Ninidae | |
| Noteridae | Noterid Water Beetles |
| Notonectidae | Backswimmers |
| Nymphalidae | Browns, Nymphs, Danaines |
| Ochteridae | Velvety Shore Bugs |
| Oeconesidae | Caddisflies |
| Oniscigastridae | Mayflies |
| Oxycarenidae | |
| Oxygastridae | Orange Streamcruiser Dragonfly |
| Pachygronthidae | |
| Papilionidae | Swallowtails |
| Peloriidae | Moss Bugs |
| Pentatomidae | Stinkbugs |
| Petaluridae | Petaltail Dragonflies |
| Philopotamidae | Finger Net caddisflies, Silken Tube Spinners |
| Philorheithridae | Caddisflies |
| Pieridae | Whites and Yellows |
| Piesmatidae | |
| Plataspidae | |
| Plectrotarsidae | Caddisflies |
| Pleidae | Pygmy Backswimmers |
| Polycentropodidae | Caseless Caddisflies |
| Prosopistomatidae | Mayflies |
| Protoneuridae | Threadtail Damselflies |
| Pseudococcidae | Mealy Bugs |
| Pseudocorduliidae | Mistfly Dragonflies |
| Psyllidae | |
| Pyrrhocoridae | |
| Reduviidae | Assassin Bugs |
| Rhinotermitidae | Termite |
| Rhopalidae | |
| Rhyparochromidae | Seed Bugs |
| Schizopteridae | |
| Scutelleridae | |
| Stenopsychidae | Caddisflies |
| Stenotritidae | |
| Synlestidae | Whitetip and Needle Damselflies |
| Synthemistidae | Spottail and Tigertail Dragonflies |
| Tasimiidae | Tasimiidae |
| Telephlebiidae | Darner Dragonflies |
| Termitidae | Termite |
| Termopsidae | Termite |
| Tessaratomidae | |
| Tettigarctidae | |
| Thaumastocoridae | |
| Therevidae | Stiletto Flies |



Invertebrate fauna

| | |
|----------------|---------------------------------------|
| Tingidae | Lacebugs |
| Triozidae | |
| Urothemistidae | Basker, Pennant and Baron Dragonflies |
| Veliidae | Small Water Striders, Riffle Bugs |

Malacostraca

Crabs, lobsters, shrimps and woodlice

| | |
|------------------|-------------------------------------|
| Eusiridae | |
| Hadziidae | |
| Hyalidae | |
| Hypsimepodidae | Phreatoicidan Isopods |
| Melitidae | |
| Neoniphargidae | |
| Paracalliopiidae | |
| Paramelitidae | |
| Phreatoicidae | Phreatoicidan Isopods |
| Phreatoicopsinae | Phreatoicidan Isopods |
| Talitridae | sandfleas, sandhoppers, landhoppers |

Polychaeta

Bristleworms, Scale Worms, Fan Worms, Rag Worms

| | |
|--------------|------------------|
| Aphroditidae | sea mice |
| Cirratulidae | Cirratulid worms |
| Sigalionidae | scale worm |



Vascular flora

Cycadopsida

Cycadaceae
Zamiaceae

Cycads

Native Cycads
Native Cycads

Liliopsida

Alismataceae
Amaryllidaceae
Anarthriaceae
Aponogetonaceae
Araceae
Arecaceae
Asphodelaceae
Asteliaceae
Blandfordiaceae
Boryaceae
Burmanniaceae
Campynemataceae
Centrolepidaceae
Colchicaceae
Cyperaceae
Dioscoreaceae
Ecdeiocoleaceae
Eriocaulaceae
Flagellariaceae
Haemodoraceae
Hemerocallidaceae
Hydrocharitaceae
Hypoxidaceae
Iridaceae
Juncaceae
Juncaginaceae
Luzuriagaceae
Musaceae
Orchidaceae
Pandanaeae
Petermanniaceae
Philesiaceae
Philydraceae
Poaceae
Pontederiaceae
Potamogetonaceae
Restionaceae
Ripogonaceae
Ruppiaceae
Smilacaceae
Sparganiaceae
Stemonaceae
Taccaceae
Thismiaceae
Typhaceae
Xanthorrhoeaceae

Monocots

Water Plantains

Pond Lilies
Settlers Flax, Cunjevoi, Native Lilies
Palms
Native Leek
Palm-Lilies and Allies
Christmas Bells
Pincushion Lillies
Burmannias

Centrolepid Sedges
Nancies and Allies
Sedges
Native Yams

Pipeworts
Whip Vine
Bloodroots, Conostyles, Kangaroo Paws and their Allies

Swamp Lily, Eel Weed, Frogbit and Water Thyme
Native Lilies
Irises and Allies
Rushes
Water Ribbons
Orange and Turquoise Berries
Native Bananas
Orchids
Pandans and Allies

Frogmouths and Stream Lilies
Grasses
Water Hyacinths and Allies
Sea Grasses and Horned Pondweeds, Pondweed
Restiona Sedges

Lawyer Vines, Supplejack, Wombat Berry and Scrambling Lily
Floating Bur Reed
Stemona Climbers
Arrowroot and Allies
Fairy Lanterns
Cumbungi and Allies
Grass Trees, Lomandras and Allies



Vascular flora

Xyridaceae
Zingiberaceae

Xyris Herbs
Native Ginger and Allies

Magnoliopsida

Achariaceae
Aizoaceae
Akaniaceae
Amaranthaceae
Anacardiaceae
Aphanopetalaceae
Asteraceae
Atherospermataceae
Austrobaileyaceae
Balanopaceae
Balanophoraceae
Bombacaceae
Boraginaceae
Burseraceae
Calycanthaceae
Cardiopteridaceae
Caryophyllaceae
Casuarinaceae
Chenopodiaceae
Combretaceae
Connaraceae
Convolvulaceae
Cunoniaceae
Dichapetalaceae
Dilleniaceae
Droseraceae
Elaeagnaceae
Epacridaceae
Ericaceae
Erythroxylaceae
Euphorbiaceae
Eupomatiaceae
Fabaceae
Haloragaceae
Hernandiaceae
Himantandraceae
Hydatellaceae
Lauraceae
Leeaceae
Lentibulariaceae
Limeaceae
Linderniaceae
Lythraceae
Melastomataceae
Meliaceae
Menispermaceae
Menyanthaceae

Dicots

Pigfaces
Turnip Wood
Pigweed
Bollygum, Native Plum

Daisies and Allies

Austrobaileyas
Pimplebark
Balanophoras
Baobabs
Forget-me-not
Canarium and Garuga

Peripterygium Vines
Pin Cushion, Wort
She-oaks
Saltbushes and Allies
Plums, Bullwaddies, Mangroves and Bushwillow
Connara Climbers
Bindweed, Morning Glory
Christmas Bushes and Mararas
Dichapetals
Guinea Flowers
Sundews
Oleaster
Southern Heaths
Heathers and Rhododendrons
Erythroxylum
Spurges
Bolwarra
Peas
Raspworts and Milfoils
Hernandias and Valvantheras
Galbulimimas
Hydatellid Herbs
Laurels, Camphorwood and Australian Walnuts
Leeas
Bladderworts

Loosestrifes
Native Lasiandra and Allies
Rosewoods and Mahoganies
Snake, Round-leaf and Pearl Vines
Marshwort



Vascular flora

| | |
|------------------|--|
| Molluginaceae | Carpet Weed, Herb |
| Monimiaceae | Sassafras, Beech |
| Myrtaceae | Native Myrtles |
| Nelumbonaceae | Waterlilies |
| Nepenthaceae | Nepenthes Pitcher Plants |
| Nitrariaceae | |
| Nyctaginaceae | Tarvine, Pisonia and Birdlime Trees |
| Ochnaceae | Brackenridgea |
| Onagraceae | Water Primrose and Willowherbs |
| Opiliaceae | Oplilia Climbers |
| Orobanchaceae | |
| Phrymaceae | |
| Phyllanthaceae | |
| Phytolaccaceae | Introduced Monococcus |
| Picrodendraceae | |
| Portulacaceae | Purslane |
| Proteaceae | Banksias, Grevilleas and Allies |
| Putranjivaceae | |
| Rhizophoraceae | Mangroves |
| Rutaceae | Boronia, Correa, Citrus, Phebalium, Philotheca, Zieria and Allie |
| Sapindaceae | Tamarind, Whitewood, Tuckeroo, Hop Bushes |
| Scrophulariaceae | Figworts |
| Simaroubaceae | Native Plum and Allies |
| Solanaceae | Native Gooseberries, Wild Tomatos and Boxthorns |
| Sphenocleaceae | |
| Sterculiaceae | Kurrajongs and Bottletrees |
| Stylidiaceae | Triggerplants and Allies |
| Thymelaeaceae | Pimeleas |
| Tiliaceae | |
| Trimeniaceae | Trimenia |
| Vitaceae | |
| Winteraceae | Pepper Trees |
| Zygophyllaceae | Zygophyllaceae |
| Pinopsida | Pines and Allies |
| Araucariaceae | Araucaria Pines |
| Cupressaceae | Native Cypress |
| Podocarpaceae | Podocarp Pines |



Appendix 2 Acknowledgements

Thank you to the following organisations and individuals for providing species location data used in the Australian Natural Heritage Assessment Tool (ANHAT) to generate this Biodiversity Summary.

Council of Heads of Australian Faunal Collections (CHAFC)

ANHAT acknowledges the Council of Heads of Australian Fauna Collections (CHAFC) for supply of data from the following CHAFC member institutions:

- Australian Museum
- CSIRO Australian National Insect Collection
- Museum and Art Gallery of the Northern Territory
- Museum Victoria
- Queen Victoria Museum and Art Gallery (Launceston)
- Queensland Museum
- South Australian Museum
- Tasmanian Museum and Art Gallery (Hobart)
- Western Australian Museum

The taxonomic concepts used in this report reflect an ANHAT view of the data and not necessarily that of the CHAFC parent Museums.

Council of Heads of Australasian Herbaria (CHAH)

ANHAT acknowledges the Council of Heads of Australasian Herbaria Inc. (CHAH Inc.) and partners in Australia's Virtual Herbarium (AVH) for the supply of data from the AVH. The AVH is a collaborative project of Australian State, Territory and Commonwealth herbaria through CHAH Inc. and includes:

- Australian National Herbarium (CANB)
- National Herbarium of New South Wales (NSW) Botanic Gardens Trust
- Herbarium of the Northern Territory (DNA, NT)
- Queensland Herbarium (BRI)
- State Herbarium of South Australia (AD)
- Tasmanian Herbarium (HO)
- National Herbarium of Victoria (MEL) Royal Botanic Gardens Melbourne
- Western Australian Herbarium (PERTH).

The taxonomic concepts used in this report reflect an ANHAT view of the data and not necessarily that of the AVH parent Herbaria.



Other Government Organisations

- Commonwealth Department of Defence
- Commonwealth Department of Sustainability, Environment, Water, Population and Communities
- New South Wales - Department of Environment and Climate Change
- New South Wales - Department of Primary Industry, Forests NSW
- Northern Territory - Department of Natural Resources, Environment and the Arts
- Queensland - Environmental Protection Agency, WildNet
- South Australia - Department for Environment and Heritage (Biological Survey of South Australia Database)
- Tasmania - Department of Primary Industries and Water (Natural Values Atlas)
- Victoria - Department of Sustainability and Environment
- Western Australia - Department of Environment and Conservation

Non-government

- Birds Australia
- Mr Russell Sheil - Rotifers of Australia
- Mr David Crosby - Lepidoptera of Victoria