



Consultation Document on Listing Eligibility and Conservation Actions

***Calidris canutus* (red knot)**

You are invited to provide your views and supporting reasons related to:

- 1) the eligibility of *Calidris canutus* (red knot) for inclusion on the EPBC Act threatened species list in the Endangered category; and
- 2) the necessary conservation actions for the above species.

Evidence provided by experts, stakeholders and the general public are welcome. Responses can be provided by any interested person.

Anyone may nominate a native species, ecological community or threatening process for listing under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) or for a transfer of an item already on the list to a new listing category. The Threatened Species Scientific Committee (the Committee) undertakes the assessment of species to determine eligibility for inclusion in the list of threatened species and provides its recommendation to the Australian Government Minister for the Environment.

Draft information for your consideration of the eligibility of this species for listing as endangered starts at page 3 and information associated with potential conservation actions for this species starts at page 11. To assist with the Committee's assessment, the Committee has identified a series of specific questions on which it seeks your guidance at page 12.

Responses are to be provided in writing either by email to:
species.consultation@environment.gov.au

or by mail to:

The Director
Migratory Species Section
Wildlife, Heritage and Marine Division
Department of the Environment
PO Box 787
Canberra ACT 2601

Responses are required to be submitted by 4 December 2015

Contents of this information package	Page
General background information about listing threatened species	2
Information about this consultation process	2
Draft information about the red knot and its eligibility for listing	3
Conservation actions for the species	11
References cited	12
Collective list of questions – your views	13

General background information about listing threatened species

The Australian Government helps protect species at risk of extinction by listing them as threatened under Part 13 of the EPBC Act. Once listed under the EPBC Act, the species becomes a Matter of National Environmental Significance (MNES) and must be protected from significant impacts through the assessment and approval provisions of the EPBC Act. More information about threatened species is available on the department's website at: <http://www.environment.gov.au/biodiversity/threatened/index.html>.

Public nominations to list threatened species under the EPBC Act are received annually by the department. In order to determine if a species is eligible for listing as threatened under the EPBC Act, the Threatened Species Scientific Committee (the Committee) undertakes a rigorous scientific assessment of its status to determine if the species is eligible for listing against a set of criteria. These criteria are available on the Department's website at: <http://www.environment.gov.au/biodiversity/threatened/pubs/guidelines-species.pdf>.

As part of the assessment process, the Committee consults with the public and stakeholders to obtain specific details about the species, as well as advice on what conservation actions might be appropriate. Information provided through the consultation process is considered by the Committee in its assessment. The Committee provides its advice on the assessment (together with comments received) to the Minister regarding the eligibility of the species for listing under a particular category and what conservation actions might be appropriate. The Minister decides to add, or not to add, the species to the list of threatened species under the EPBC Act. More detailed information about the listing process is at: <http://www.environment.gov.au/biodiversity/threatened/nominations.html>.

To promote the recovery of listed threatened species and ecological communities, conservation advices and where required, recovery plans are made or adopted in accordance with Part 13 of the EPBC Act. Conservation advices provide guidance at the time of listing on known threats and priority recovery actions that can be undertaken at a local and regional level. Recovery plans describe key threats and identify specific recovery actions that can be undertaken to enable recovery activities to occur within a planned and logical national framework. Information about recovery plans is available on the department's website at: <http://www.environment.gov.au/biodiversity/threatened/recovery.html>.

Information about this consultation process

Responses to this consultation can be provided electronically or in hard copy to the contact addresses provided on Page 1. All responses received will be provided in full to the Committee and then to the Australian Government Minister for the Environment.

In providing comments, please provide references to published data where possible. Should the Committee use the information you provide in formulating its advice, the information will be attributed to you and referenced as a 'personal communication' unless you provide references or otherwise attribute this information (please specify if your organisation requires that this information is attributed to your organisation instead of yourself). The final advice by the Committee will be published on the department's website following the listing decision by the Minister.

Information provided through consultation may be subject to freedom of information legislation and court processes. It is also important to note that under the EPBC Act, the deliberations and recommendations of the Committee are confidential until the Minister has made a final decision on the nomination, unless otherwise determined by the Minister.

Calidris canutus

Red knot

Taxonomy

Conventionally accepted as *Calidris canutus*, Linnaeus 1758.

Other common names: knot, common knot, Iceland sandpiper, East Siberian sandpiper, lesser knot.

The species is polytypic, meaning more than one subspecies exists. Globally, the following six subspecies are recognised:

- *Calidris canutus canutus* (nominate subspecies) breeds in central Siberia;
- *C.c. piersmai* breeds in the New Siberian Islands;
- *C.c. rogersi* breeds on Chukotka Peninsula (north-eastern Siberia);
- *C.c. roselaari* breeds at Wrangel Island, Siberia, and north-west Alaska;
- *C.c. rufa* breeds in the Canadian Arctic, south of 75 °N; and
- *C.c. islandica* breeds on the islands of the Canadian high Arctic and northern Greenland (Bamford et al. 2008; Leyrer et al. 2014; Gill & Donsker 2015).

Two subspecies, *C. c. piersmai* and *C. c. rogersi*, regularly occur in Australia (Garnett et al. 2011). One other subspecies, *C. c. canutus*, is considered a vagrant in Australia (Garnett et al. 2011).

Species/Sub-species Information

Description

The red knot is a small to medium migratory shorebird. It has a length of 23–25 cm, a wingspan of 45–54 cm and a weight of 120 g. The species is robust, short-necked with a short straight bill, long wings extending beyond the tail and short legs (Higgins & Davies 1996). The red knot has a faint pale brow line. The upper body is brownish grey with fine dark streaks on the head and neck and the underbody is white with some light streaking. In breeding plumage, the upper body is boldly marked, contrasting with the mainly chestnut-red body (BirdLife Australia 2012).

The two subspecies *C. c. piersmai* and *C. c. rogersi* cannot be distinguished from each other in nonbreeding plumage (Rogers et al. 2010). However, the location in which the individual is present can help distinguish the two subspecies, with *C. c. piersmai* tending to overwinter almost exclusively in North-west Australia, and *C. c. rogersi* tending to overwinter in other parts of Australasia (del Hoyo et al. 2014).

Breeding plumage:

- *C. c. piersmai*: Deep brick-red underparts and reddish napes with black mantle and scapulars boldly marked by rufous fringes and panels within each feather. Many scapulars have narrow white tips but these are inconspicuous;
- *C. c. rogersi*: Paler, peachier underparts, and whitish napes with more extensive silvery variegation on the upperparts caused by a combination of broader grey-white tips to individual feathers, the presence of many scapulars with grey-white panels within the feather, and only a pale rufous tinge to other markings within the scapulars (Rogers et al. 2010).

Distribution

Global Distribution

The red knot (all six subspecies combined) has a global distribution and an extremely large range. The species breeds at a range of locations around the Arctic and, for the boreal winter,

migrates to non-breeding areas that extend to the southernmost parts of the Americas, Africa, Europe and Australasia (del Hoyo et al. 1996). For the species, the global extent of occurrence is estimated to be 1,600,000 km² (BirdLife International 2015).

Australian Distribution

The red knot breeds in the northern hemisphere and undertakes migrations along the East Asian-Australasian Flyway (EAAF) to spend the boreal winter in Australasia. The bulk of the population is considered to spend the non-breeding period in Australia (Bamford et al. 2008).

The red knot is common in all the main suitable habitats around the coast of Australia (Barrett et al. 2002), is less numerous in south-west Australia and is occasionally recorded inland in all regions (Higgins & Davies 1996). Very large numbers are regularly recorded in north-west Australia. In Queensland, the red knot migrates along the coast north of 19 °S, sometimes in large numbers. It is widespread along the coast south of Townsville, along the coasts of NSW and Victoria, and is a regular visitor, in small numbers, to the coasts of Tasmania. In South Australia, the species is found mostly from The Coorong, north and west to the Yorke Peninsula and Port Pirie. The Red Knot has also been recorded at Norfolk Island, Lord Howe Island, Macquarie Island, Kermadec Island, Chatham Islands, Auckland Islands and Campbell Islands (Higgins & Davies 1996). Red knots migrating to New Zealand may stage in Australia, particularly in the Gulf of Carpentaria (Bamford et al. 2008; Rogers et al. 2010; Garnett et al. 2011).

Relevant Biology/Ecology

Life History

The generation time of 7.8 years (Garnett et al., 2011) is derived from an age of first breeding of 2.0 years (Cramp et al. 1983), an adult survival of 68% (Boyd 1962) and a maximum longevity of 16.2 years and 19.2 years for *C. c. rogersi* and *C. c. piersmai* respectively (DOE 2015).

Breeding

The red knot does not breed in Australia. Red knots breed in north Siberia and Alaska during the austral winter (DEWHA 2009b). In June, the red knot lays 3-4 eggs and incubation lasts for around 21–22 days. On hatching, females depart the nest leaving the male to tend for young. Fledging occurs after 18–20 days. In one sample of 26 eggs, 54% hatched and 27% fledged (del Hoyo et al. 1996).

General Habitat

The red knot breeds on dry upland tundra in high Arctic areas. During the boreal summer, they nest on open vegetated tundra or stone ridges, often close to a clump of vegetation. Breeding density is normally around one pair per km² (del Hoyo et al. 1996).

During the non-breeding season in Australasia, the red knot mainly inhabit intertidal mudflats, sandflats and sandy beaches of sheltered coasts and sometimes on sandy ocean beaches or shallow pools on exposed rock platforms. They are occasionally seen on terrestrial saline wetlands near the coast and on sewage ponds and saltworks (Higgins & Davies 1996).

Feeding Habitat

The red knot usually forages in soft substrate near the water edge on intertidal mudflats or sandflats exposed by low tide. At high tide they may feed at nearby lakes, sewage ponds and floodwaters (Higgins & Davies 1996).

Roosting Habitat

The red knot roosts on sandy beaches, spits and islets, and mudflats (Higgins & Davies 1996). They have been seen roosting on an inland claypan near Roebuck Bay, north-west Western Australia (Collins et al. 2001). They like to roost in open areas far away from potential cover for predators, but close to feeding grounds (Rogers 2001). In hot conditions, shorebirds prefer to roost where a damp substrate lowers the local temperature and different roosts were used at night when birds chose safer, but more distant, roosts from foraging areas (Rogers et al. 2006).

Feeding

The red knot eats worms, bivalves, gastropods, crustaceans and echinoderms (Higgins & Davies 1996). In Australia, they predominantly forage on shellfish by being able to detect pore-water pressure differentials to locate hard, buried prey (Piersma et al. 1998). They have a large muscular gizzard for crushing bivalves which are swallowed whole (Piersma et al. 1993; van Gils et al. 2005; Rogers et al. 2010). In some circumstances they also visually locate prey items and sometimes take prey from the surface (Rogers 2001).

The red knot is diurnal and nocturnal. They forage in large, dense, often mixed-species flocks, with birds rapidly, intensively and methodically probing the wet mud as they walk quickly across the mudflats exposed by the falling tide (BirdLife Australia 2012). In non-breeding areas, feeding activity is regulated by the tide with birds closely following the tide-edge when foraging (Rogers 2001).

Migration Patterns

The red knot is migratory, breeding in the high Arctic and moving south to non-breeding areas to approximately 50 °S. They are capable of flying non-stop between north-eastern China and northern Australia and tend to use only a few staging areas (Bamford et al. 2008).

Departure from breeding grounds

The subspecies *C. c. rogersi* breeds in north-east Siberia, including around the Chukotka Peninsula and possibly farther west, and migrates mainly to Australia and New Zealand. Although the route of migration to Australia is not known it may move in a loop, migrating south across the west Pacific Ocean and north along the east Asian coast. The subspecies *C. c. piersmai* breeds in the New Siberian Islands and migrates along the coast of east Asia, with some birds reaching Australia and New Zealand (Higgins & Davies 1996).

Non-breeding season

In Australia, most red knots arrive on the north-west coast and the Gulf of Carpentaria from late August (Higgins & Davies 1996). They move south, mostly along coasts, with some inland records from September–November and arrive in south-west Australia from September (Higgins & Davies 1996). Information derived from banding and flagging programs suggests that the population that remains in north-west Australia is mostly the subspecies *C. c. piersmai*, although some may also occur in eastern Australia. The subspecies *C. c. rogersi* mainly occur in eastern Australia and New Zealand although some of these birds pass through north-west Australia on migration (Rogers et al. 2010).

During the non-breeding period, around 93% of the EAAF population of the red knot (subspecies *C.c. piersmai* and *C.c. rogersi*) occurs in Australia and New Zealand, with smaller numbers in China, Indonesia and other countries in southeast Asia (Bamford et al. 2008).

Return to breeding grounds

Red knots leave Tasmania from February–May and leave south-east mainland Australia from late February or late March to early April. Inland records suggest that some birds move overland on northern migration. They leave north-west Australia from late March to late April. Most

probably passing through the northern half of the Yellow Sea (Barter 2002) with large numbers seen in the Korean Peninsula in April and May. Some birds overwinter in Australia, mainly northern Australia (Higgins & Davies 1996).

Internationally, the Yellow Sea is extremely important as stopover habitat for red knot, with over 45% of the EAAF population using a single site at Bohai Bay, Yellow Sea during their migration (Rogers et al. 2010; Iwamura et al. 2013).

Threats

Migratory shorebirds, such as the red knot, are sensitive to certain development activities due to their high site fidelity, tendency to aggregate, very high energy demands, and need for habitat networks containing both roosting and foraging sites (DEWHA 2009a).

Threats to the global population of the red knot across its range include habitat loss and habitat degradation (e.g. through land reclamation, industrial use and urban expansion, changes to the water regime, invasive plants and environmental pollution), over-exploitation of shellfish, pollution/contamination impacts, disturbance, direct mortality (hunting), diseases, extreme weather events, and climate change impacts (DEWHA 2009b; BirdLife International 2015).

Habitat loss and habitat degradation

The red knot is threatened by wetland degradation in East Asia, where it stages on migration (Bamford et al. 2008). The red knot is specifically threatened at Bohai Bay, Yellow Sea where both subspecies (*C. c. piersmai* and *C. c. rogersi*) stage on the intertidal mudflats. Rogers et al. (2010) estimated that their study site area of 20 km of coastline in Bohai Bay was used by over 45% of the combined global population of adult *C. c. piersmai* and *C. c. rogersi*. Between 1994 and 2010, the reclamation of large areas (including intertidal mudflats) in the bay for two industrial projects caused the northward migrating red knot to become concentrated in an ever smaller remaining area. The northward migration numbers of *C. c. piersmai* and *C. c. rogersi* in this so far little affected area increased from 13% in 2007 to 62% in 2010 of the global populations (Yang et al. 2011). With the proposed continuation of land reclamation in Bohai Bay, it is predicted that shorebird densities in the remaining areas will increase to a point of collapse (Yang et al. 2011). Along with other major areas of tidal flat habitat in East Asia, the Bohai Bay tidal flats currently have no formal protection (Murray & Fuller 2015). Reclamation on intertidal mudflats is also a threat in other areas of the EAAF, for example Malaysia (Wei et al. 2006). In addition, intensive oil exploration and extraction, and reduction in river flows due to upstream water diversion, are other potentially significant threats in parts of China where this species is present in internationally significant numbers (Barter 2005; Barter et al. 1998).

In Australia, the loss of important habitat reduces the availability of foraging and roosting sites. This affects the ability of the birds to build up the energy stores required for successful migration and breeding. Some sites are important all year round for juveniles who may stay in Australia throughout the breeding season until they reach maturity. A variety of activities may cause habitat loss which include direct losses through land clearing, inundation, infilling or draining. Indirect loss may occur due to changes in water quality, hydrology or structural changes near roosting sites (DEWHA 2009b).

As most migratory shorebirds, such as the red knot, have specialized feeding techniques, they are particularly susceptible to slight changes in prey sources and foraging environments. Activities that cause habitat degradation include, but are not restricted to loss of marine or estuarine vegetation, which is likely to alter the dynamic equilibrium of sediment banks and mudflats; invasion of intertidal mudflats by weeds such as cordgrass; water pollution and changes to the water regime; changes to the hydrological regime; and exposure of acid sulphate soils, hence changing the chemical balance at the site (DEWHA 2009b).

The non-breeding grounds of the species in south-eastern Australia are threatened by habitat degradation, loss and human disturbance (Garnett et al. 2011), but those in the north are generally free of such disturbances (NTDoLRM 2012).

Pollution/contamination

Migratory shorebirds may be adversely affected by pollution, both on passage and in non-breeding areas (Harding et al. 2007; Wei et al. 2006).

Disturbance

Human disturbance can cause shorebirds to interrupt their feeding or roosting and may influence the area of otherwise suitable feeding habitat that is actually used. Disturbance from human recreation activities may force migratory shorebirds to increase the time devoted to vigilance and anti-predator behaviour and/or may compel the birds to move to alternative, less favourable feeding areas (Goss-Custard et al. 2006).

Disturbance can result from residential and recreational activities including fishing, boating, four wheel driving, walking dogs, noise and night lighting. While some disturbances may have a low impact, it is important to consider the combined effect of disturbances with other threats (DEWHA 2009b).

Diseases

The red knot is susceptible to avian influenza and so may be threatened by future outbreaks of the virus (Melville & Shortridge 2006).

Since, 1992, the viral disease testing of Charadriiformes from coastal northwest Australia has not detected any evidence of avian influenza virus excretion in the red knot or any other species tested. However, from serologic testing, there was evidence of past exposure to the virus in the sampled red knots and the exposure risk profile for this species had significantly higher values compared to other species (Curran et al. 2014).

Direct mortality

Direct mortality may result from the construction of wind farms located in migration or movement pathways, bird strike with aircraft, hunting, chemical spills and oil spills (DEWHA 2009b). Hunting is still a very serious problem for shorebirds in China, and the red knot has been identified as one of the species caught (Ming et al. 1998).

Climate change

Global warming and associated changes in sea level are likely to have a long-term impact on the breeding, staging and non-breeding grounds of migratory shorebirds (Harding et al. 2007). Rises in sea level could have a major impact on the red knot due to loss of intertidal habitat (Iwamura et al. 2013). Taking into account upshore movements of intertidal habitat, modelling indicates that, for both *C.c. piersmai* and *C.c. rogersi*, population flow could reduce by 15% with a 150 cm sea level rise (Iwamura et al. 2013).

Assessment of available information in relation to the EPBC Act Criteria and Regulations

Criterion 1. Population size reduction (reduction in total numbers)

Population reduction (measured over the longer of 10 years or 3 generations) based on any of A1 to A4

Critically Endangered
Very severe reduction

Endangered
Severe reduction

Vulnerable
Substantial reduction

A1	≥ 90%	≥ 70%	≥ 50%
A2, A3, A4	≥ 80%	≥ 50%	≥ 30%
A1	Population reduction observed, estimated, inferred or suspected in the past and the causes of the reduction are clearly reversible AND understood AND ceased.		(a) direct observation [except A3]
A2	Population reduction observed, estimated, inferred or suspected in the past where the causes of the reduction may not have ceased OR may not be understood OR may not be reversible.	<i>based on any of the followin</i>	(b) an index of abundance appropriate to the taxon
A3	Population reduction, projected or suspected to be met in the future (up to a maximum of 100 years) [(a) cannot be used for A3]		(c) a decline in area of occupancy, extent of occurrence and/or quality of habitat
A4	An observed, estimated, inferred, projected or suspected population reduction where the time period must include both the past and the future (up to a max. of 100 years in future), and where the causes of reduction may not have ceased OR may not be understood OR may not be reversible.		(d) actual or potential levels of exploitation
			(e) the effects of introduced taxa, hybridization, pathogens, pollutants, competitors or parasites

Evidence:

Eligible under Criterion 1 A2 (a) for listing as Endangered

The global population of red knot was previously estimated at 1,090,000 with the population in the EAAF at 220,000 (Bamford et al. 2008). Although numbers at several sites have declined, it is also thought that the previous estimate of red knots at Eighty Mile Beach, WA (80,700), based on extrapolation from aerial surveys, may have been inflated (Rogers et al. 2010). Rogers et al. (2010) presented revised estimates for Australian and New Zealand sites using the most recently available austral summer counts. Assuming around 93% of the EAAF population of the red knot occurs in Australia and New Zealand (Bamford et al. 2008), a revised population estimate for the EAAF, based on a sum of revised estimates for the Australian and New Zealand populations of 104,986 (Rogers et al. 2010), is approximately 112,000 individuals, of which 68,000 occur in Australia (Garnett et al., 2011).

In Australia, direct counts of red knots at key sites (e.g. Rogers et al. 2009) have shown a population decline of more than 30% over the last 20 years. For example, numbers in Victoria showed a decline in count data from 4,474 to 2,419 individuals (Wilson 2001). Numbers at Eighty-mile Beach declined by c.78% between 2000 and 2008 (Rogers et al. 2009), at Moreton Bay by 75% between 1993 and 2008 (Fuller et al. 2009) and by c.27% across 49 Australian sites between c.1983 and c.2007 (Garnett et al. 2011).

Numbers of red knots appear to have had a less severe decline elsewhere in the EAAF e.g. no clear trends in Japan between 1978 and 2008 (Amano et al. 2010).

A subsequent and more detailed assessment by a University of Queensland team (partly funded by the Department of the Environment under an Australian Research Council collaborative grant), puts the species into the endangered category (Fuller, pers. comm., 2014). Time series data from directly observed summer counts at a large number of sites across Australia indicate a severe population decline of 62.0% over 23 years (4.4% per year) which for this species is equal to three generations (Fuller, pers. comm. 2014).

In large part, the observed decline in red knot numbers across Australia stems from ongoing loss of intertidal mudflat habitat at key migration staging sites in the Yellow Sea (Murray et al., 2014). As such, qualification under criterion A2 rather than A1 seems warranted. However, threats are also occurring in Australia including coastal development and recreational activities causing disturbance.

The data presented above appear to demonstrate that the species is **eligible for listing as Endangered A2a** under this criterion. However, the purpose of this consultation document is to elicit additional information to better understand the species' status. This conclusion should therefore be considered to be tentative at this stage, as it may be changed as a result of responses to this consultation process.

Criterion 2. Geographic distribution as indicators for either extent of occurrence AND/OR area of occupancy			
	Critically Endangered Very restricted	Endangered Restricted	Vulnerable Limited
B1. Extent of occurrence (EOO)	< 100 km ²	< 5,000 km ²	< 20,000 km ²
B2. Area of occupancy (AOO)	< 10 km ²	< 500 km ²	< 2,000 km ²
AND at least 2 of the following 3 conditions indicating distribution is precarious for survival:			
(a) Severely fragmented OR Number of locations	= 1	≤ 5	≤ 10
(b) Continuing decline observed, estimated, inferred or projected in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) area, extent and/or quality of habitat; (iv) number of locations or subpopulations; (v) number of mature individuals			
(c) Extreme fluctuations in any of: (i) extent of occurrence; (ii) area of occupancy; (iii) number of locations or subpopulations; (iv) number of mature individuals			

Evidence:

Not eligible

The extent of occurrence in Australia is estimated to be 36 000 km² (stable) and area occupied 2 400 km² (stable; Garnett et al., 2011).

The data presented above appear to demonstrate the species is not eligible for listing under this criterion. However, the purpose of this consultation document is to elicit additional information to better understand the species' status. This conclusion should therefore be considered to be tentative at this stage, as it may be changed as a result of responses to this consultation process.

Criterion 3. Population size and decline			
	Critically Endangered Very low	Endangered Low	Vulnerable Limited
Estimated number of mature individuals	< 250	< 2,500	< 10,000
AND either (C1) or (C2) is true			
C1 An observed, estimated or projected continuing decline of at least (up to a max. of 100 years in future)	Very high rate 25% in 3 years or 1 generation (whichever is longer)	High rate 20% in 5 years or 2 generation (whichever is longer)	Substantial rate 10% in 10 years or 3 generations (whichever is longer)
C2 An observed, estimated, projected or inferred continuing decline AND its geographic distribution is precarious for its survival based on at least 1 of the following 3 conditions:			
(a) (i) Number of mature individuals in each subpopulation	≤ 50	≤ 250	≤ 1,000
(a) (ii) % of mature individuals in one subpopulation =	90 – 100%	95 – 100%	100%
(b) Extreme fluctuations in the number of			

Evidence:

Not eligible

The number of mature individuals in Australia was estimated at 68 000 in 2011 (Garnett et al., 2011), but has declined since. There are no current data available to allow assessment against this criterion.

The data presented above appear to demonstrate the species is not eligible for listing under this criterion. However, the purpose of this consultation document is to elicit additional information to better understand the species' status. This conclusion should therefore be considered to be tentative at this stage, as it may be changed as a result of responses to this consultation process.

Criterion 4. Number of mature individuals			
	Critically Endangered Extremely low	Endangered Very Low	Vulnerable Low
Number of mature individuals	< 50	< 250	< 1,000

Evidence:

Not eligible

The total number of mature individuals was estimated at 68 000 in 2011 (Garnett et al., 2011), but has declined since. The estimate is not considered extremely low, very low or low.

The data presented above appear to demonstrate the species is not eligible for listing under this criterion. However, the purpose of this consultation document is to elicit additional information to better understand the species' status. This conclusion should therefore be considered to be tentative at this stage, as it may be changed as a result of responses to this consultation process.

Criterion 5. Quantitative Analysis			
	Critically Endangered Immediate future	Endangered Near future	Vulnerable Medium-term future
Indicating the probability of extinction in the wild to be:	≥ 50% in 10 years or 3 generations, whichever is longer (100 years max.)	≥ 20% in 20 years or 5 generations, whichever is longer (100 years max.)	≥ 10% in 100 years

Evidence:

Not eligible

Population viability analysis has not been undertaken

The data presented above appear to demonstrate the species is not eligible for listing under this criterion. However, the purpose of this consultation document is to elicit additional information to better understand the species' status. This conclusion should therefore be considered to be

tentative at this stage, as it may be changed as a result of responses to this consultation process.

Conservation Actions

Recovery Plan

A decision about whether there should be a recovery plan for this species has not yet been determined. The purpose of this consultation document is to elicit additional information to help inform this decision.

Conservation and Management Actions

- Work with governments along the East Asian – Australasian Flyway to prevent destruction of key migratory staging sites.
- Protect important habitat in Australia.
- Support initiatives to improve habitat management at key sites.
- Maintain and improve protection of roosting and feeding sites in Australia.
- Advocate for the creation and restoration of foraging and roosting sites in Australia.
- Incorporate requirements for red knot into coastal planning and management.
- Manage important sites to identify, control and reduce the spread of invasive species.
- Manage disturbance at important sites when red knot are present – e.g. discourage or prohibit vehicle access, horse riding and dogs on beaches, implement temporary site closures.
- Monitor the progress of recovery, including the effectiveness of management actions and the need to adapt them if necessary.

Survey and Monitoring priorities

- Enhance existing migratory shorebird population monitoring programmes, particularly to improve coverage across northern Australia

Information and research priorities

- Undertake work to more precisely assess red knot life history, population size, distribution and ecological requirements.
- Improve knowledge about dependence of red knot on key migratory staging sites, and non-breeding sites in south-east Asia.
- Improve knowledge about threatening processes including the impacts of disturbance and hunting.

Collective list of questions – your views

1. Do you agree with the current taxonomic position of the Australian Faunal Directory and Birdlife Australia for this species (as identified in the draft conservation advice)
2. Can you provide any additional references, information or estimates on longevity, age of maturity, average life span and generation length?
3. Has the survey effort for this species been adequate to determine its national distribution and adult population size?
4. Do you accept the estimate provided in the nomination for the current population size of the species?
5. For any population with which you are familiar, do you agree with the population estimate provided? If not, are you able to provide a plausible estimate based on your own knowledge? If so, please provide in the form:
 - Lower bound (estimated minimum):
 - Upper bound (estimated maximum):
 - Best Estimate:
 - Estimated level of Confidence: %
6. Can you provide any additional data, not contained in the current nomination, on declines in population numbers over the past or next 10 years or 3 generations, whichever is the longer?
7. Is the distribution as described in the nomination valid? Can you provide an estimate of the current geographic distribution (extent of occurrence or area of occupancy in km²) of this species?
8. Has this geographic distribution declined and if so by how much and over what period of time?
9. Do you agree that the species is eligible for inclusion on the threatened species list, in the category listed in the nomination?
10. Do you agree that the threats listed are correct and that their effects on the species are significant?
11. To what degree are the identified threats likely to impact on the species in the future?
12. Can you provide additional or alternative information on threats, past, current or potential that may adversely affect this species at any stage of its life cycle?
13. In seeking to facilitate the recovery of this species, can you provide management advice for the following:
 - What individuals or organisations are currently, or need to be, involved in planning to abate threats and any other relevant planning issues?
 - What threats are impacting on different populations, how variable are the threats and what is the relative importance of the different populations?
 - What recovery actions are currently in place, and can you suggest other actions that would help recover the species? Please provide evidence and background information.
14. Can you provide additional data or information relevant to this assessment?

References cited in the advice

- Amano T., T. Székely, K. Koyama, H. Amano & W.J. Sutherland. (2010). A framework for monitoring the status of populations: an example from wader populations in the East Asian–Australasian flyway. *Biological Conservation* 143: 2238–2247.
- Bamford M., D. Watkins, W. Bancroft, G. Tischler & J. Wahl (2008). *Migratory Shorebirds of the East Asian - Australasian Flyway: Population estimates and internationally important sites*. [Online]. Canberra, ACT: Department of the Environment, Water, Heritage and the Arts, Wetlands International-Oceania. Available from: <http://www.environment.gov.au/biodiversity/migratory/publications/shorebirds-east-asia.html>.
- Barrett, G., A. Silcocks, S. Barry, R. Poulter & R. Cunningham (2002). *Australian Bird Atlas 1998-2001 Main Report To Environment Australia*. Melbourne: Birds Australia.
- Barter, M.A. (2002). Shorebirds of the Yellow Sea: Importance, Threats and Conservation Status. Wetlands International Global Series No. 8, International Wader Studies 12. Canberra, ACT: Wetlands International.
- Barter, M.A. (2005). Yellow Sea-driven priorities for Australian shorebird researchers. In: Straw, P., ed. Status and Conservation of Shorebirds in the East Asian-Australasian Flyway. Proceedings of the Australasian Shorebirds Conference 13-15 December 2003, Canberra, Australia. Sydney, NSW: Wetlands International Global Series 18, International Wader Studies 17.
- Barter, M.A., D. Tonkinson, J.Z. Lu, S.Y. Zhu, Y. Kong, T.H. Wang, Z.W. Li & X.M. Meng (1998). Shorebird numbers in the Huang He (Yellow River) Delta during the 1997 northward migration. *Stilt*. 33:15-26.
- BirdLife Australia (2012). Species factsheet: Red Knot *Calidris canutus*. <http://www.birdlife.org.au/bird-profile/red-knot> (Accessed 19/05/2015).
- BirdLife International (2015). Species factsheet: *Calidris canutus*. Downloaded from <http://www.birdlife.org> on 18/05/2015.
- Boyd, H. (1962) Mortality and fertility of European Charadrii. *Ibis* 104: 368-387.
- Collins, P., A. Boyle, C. Minton & R. Jessop (2001). The importance of inland claypans for waders in Roebuck Bay, Broome, NW Australia. *Stilt*. 38:4-8.
- Cramp, S., K.E.L. Simmons, D.C. Brooks, N.J. Collar, E. Dunn, R. Gillmor, P.A.D. Hollom, R. Hudson, E.M. Nicholson, M.A. Ogilvie, P.J.S. Olney, C.S. Roselaar, K.H. Voous, D.I.M. Wallace, J. Wattel, & M.G. Wilson (Eds) (1983). *Handbook of the Birds of Europe, the Middle East and North Africa: the Birds of the Western Palearctic: 3. Waders to Gulls*. Oxford, UK: Oxford University Press.
- Curran, J.M., T.M. Ellis & I.D. Robertson. (2014). Surveillance of Charadriiformes in Northern Australia shows species variations in exposure to Avian Influenza Virus and suggests negligible virus prevalence. *Avian Diseases* 58: 199-204.
- del Hoyo, J., A. Elliott & J. Sargatal, eds. (1996). *Handbook of the Birds of the World. Volume 3, Hoatzin to Auks*. Barcelona: Lynx Edicions.
- del Hoyo, J., N. J. Collar, D. A. Christie, A. Elliott, A. & L. D. C. Fishpool. (2014). *HBW and BirdLife International Illustrated Checklist of the Birds of the World*. Barcelona: Lynx Editions and BirdLife International.
- Department of the Environment (DOE) (2015). Australian Bird and Bat Banding Scheme Database. (Accessed 04/09/2015).

Department of the Environment, Water, Heritage and the Arts (DEWHA) (2009a). *Draft - Significant impact guidelines for 36 migratory shorebird species - Migratory species: EPBC Act policy statement 3.21* <http://www.environment.gov.au/resource/draft-significant-impact-guidelines-36-migratory-shorebird-species-migratory-species-epbc> (Accessed 04/05/2015).

Department of the Environment, Water, Heritage and the Arts (DEWHA) (2009b). *Draft - Significant impact guidelines for 36 migratory shorebird species - Migratory species: Background paper to EPBC Act policy statement 3.21* <http://www.environment.gov.au/resource/draft-significant-impact-guidelines-36-migratory-shorebird-species-migratory-species-epbc> (Accessed 04/05/2015).

Fuller, R.A. (2014). Personal Communication by email, 14 July 2014. University of Queensland.

Fuller, R.A., H.B. Wilson, B.E. Kendall & H.P. Possingham. (2009). Monitoring shorebirds using counts by the Queensland Wader Study Group. A report to the Queensland Wader Study Group and the Department of Environment and Resource Management. Brisbane, Australia.

Garnett, S., J. Szabo & G. Dutson (2011). *The Action Plan for Australian Birds 2010*. CSIRO Publishing.

Gill, F & D Donsker (Eds). (2015). IOC World Bird List (v 5.2). doi : 10.14344/IOC.ML.5.2. <http://www.worldbirdnames.org/> (Accessed 18.05.2015).

Goss-Custard, J.D., P. Triple., F. Sueur & A.D. West. (2006). Critical thresholds of disturbance by people and raptors in foraging wading birds. *Biological Conservation* 127: 88-97.

Harding, S.B., J.R. Wilson & D.W. Geering (2007). Threats to shorebirds and conservation actions. In: Geering, A., L. Agnew & S. Harding, eds. *Shorebirds of Australia*. Page(s) 197-213. Melbourne, Victoria: CSIRO Publishing.

Higgins, P.J. & S.J.J.F. Davies, eds (1996). *Handbook of Australian, New Zealand and Antarctic Birds. Volume Three - Snipe to Pigeons*. Melbourne, Victoria: Oxford University Press.

Iwamura, T., H.P. Possingham, I. Chades, C. Minton, N.J. Murray, D.I. Rogers, E.A. Treml & R.A. Fuller (2013). Migratory connectivity magnifies the consequences of habitat loss from sea-level rise for shorebird populations. *Proceedings of the Royal Society B: Biological Sciences*.

Leyrer, J., N. van Nieuwenhove, N. Crockford & S. Delany. (2014). Proposals for Concerted and Cooperative Action for Consideration by CMS COP 11, November 2014: Far Eastern Curlew *Numenius madagascariensis*, Bar-tailed Godwit *Limosa lapponica*, Great Knot *Calidris tenuirostris*, Red Knot *Calidris canutus*. http://www.cms.int/sites/default/files/document/COP11_Inf_44_Proposals_for_Concerted_and_Cooperative_Action_Bird_Species_for_Consideration_by_COP11_0.pdf (Accessed 18/05/2015).

Melville, D.S., & K.F. Shortridge. (2006). Migratory waterbirds and avian influenza in the East Asian-Australasian Flyway with particular reference to the 2003-2004 H5N1 outbreak. In: Boere, G.; Galbraith, C., Stroud, D. (ed.), *Waterbirds around the world*, pp. 432-438. The Stationary Office, Edinburgh, UK.

Ming, M., L. Jianjian, T. Chengjia, S. Pingyue & H. Wei (1998). The contribution of shorebirds to the catches of hunters in the Shanghai area, China, during 1997-1998. *Stilt*. 33:32-36.

Murray, N.J. & R.A. Fuller (2015). Protecting stopover habitat for migratory shorebirds in East Asia. *J. of Ornithology* DOI 10.1007/s10336-015-1225-2

- Piersma, T., A. Koolhaas & A. Dekinga (1993). Interactions between stomach structure and diet choice in shorebirds. *Auk* 110: 552–564.
- Piersma, T., R. van Aelst, K. Kurk, H. Berkhoudt & L.R.M. Maas (1998). A new pressure sensory mechanism for prey detection in birds: the use of principles of seabed dynamics? *Proceedings of the Royal Society of London. Series B. Biological Sciences* 265: 1377–1383.
- Rogers, D. (2001). *Conservation and ecology of migratory shorebirds in Roebuck Bay, north-western Australia*. Wetlands Unit, Environment Australia.
- Rogers, D.I., P.F. Battley, T. Piersma, J.A. van Gils & K.G. Rogers. (2006). High-tide habitat choice: insights from modelling roost selection by shorebirds around a tropical bay. *Animal Behaviour* 72: 563-575.
- Rogers, D., C.Hassell, J. Oldland, R. Clemens, A. Boyle & K. Rogers (2009). *Monitoring Yellow Sea migrants in Australia (MYSMA): north-western Australian shorebird surveys and workshops, December 2008*.
- Rogers D.I., H-Y. Yang, C.J. Hassell, A.N. Boyle, K.G. Rogers, B. Chen, Z-W. Zhang & T. Piersma (2010). Red Knots (*Calidris canutus piersmai* and *C. c. rogersi*) depend on a small threatened staging area in Bohai Bay, China. *Emu* 110: 307–315.
- Northern Territory Department of Land Resource Management (NTDoLRM). (2012). Threatened Species of the Northern Territory: Red Knot *Calidris canutus*. http://www.lrm.nt.gov.au/__data/assets/pdf_file/0016/143116/Red-Knot_VU_FINAL.pdf (Accessed 19/05/2015).
- van Gils, J.A., S.R. de Rooij, J. van der Meer, A. Dekinga, T. Piersma & R. Drent (2005). Digestive bottleneck affects foraging decisions in red knots (*Calidris canutus*). I. Prey choice. *Journal of Animal Ecology* 74: 105–119.
- Wei, D.L.Z., Y.C. Aik, L.K. Chye, K. Kumar, L.A. Tiah, Y. Chong & C.W. Mun (2006). Shorebird survey of the Malaysian coast November 2004-April 2005. *Stilt*. 49:7-18.
- Wilson, J.R. (2001). The January and February 2001 Victoria wader count. *Stilt*. 40:55-64.
- Yang, H-Y., B. Chen, M. Barter, T. Piersma, C-F Zhou, F-S. Li & Z-W Zhang. (2011). Impacts of tidal land reclamation in Bohai Bay, China: ongoing losses of critical Yellow Sea waterbird staging and wintering sites. *Bird Conservation International* 21:241–259.